

CIRUGÍA ESPAÑOLA



www.elsevier.es/cirugia

Video of the month

Laparoscopic approach to accessory wandering spleen, an uncommon association



Abordaje laparoscópico del bazo errante accesorio, una asociación poco frecuente

Ester Ferrer-Inaebnit^a, Juan José Segura-Sampedro ^{a,b,c,*}, Cristina Pineño-Flores ^{a,c}, Rafael Morales-Soriano ^a

A 46-year-old woman, with intermittent colicky abdominal pain, was diagnosed of an intraperitoneal tumor in an abdominal tomography (Fig. 1), oriented as a gastrointestinal stromal tumor.

An exploratory laparoscopy with two trocars was performed, finding a mass with a long vascular pedicle depending on the splenocolic ligament (Fig. 1). It was performed a section of the vascular pedicle by Ligasure at




Fig. 1 – Abdominal contrast-enhanced computed tomography: $32 \times 32 \times 34$ mm homogeneus hypervascular well-defined solid tumor with a small central hypodensity (red arrow) located in the intraperitoneal left posterior abdominal void, with an apparent intraperitoneal location. The dependency organ is not defined, only broad contact is seen with adjacent intestinal loops but without thickening or stenosing them.

^a General and Digestive Surgery Service: Hospital Universitario Son Espases, Palma de Mallorca, Spain

^b Faculty of Medicine, Universidad de las Islas Baleares, Palma de Mallorca, Spain

^c Health Research Institute of the Balearic Islands (IdISBa), Palma de Mallorca, Spain

^{*} Corresponding author.

the infrasplenic level, with extraction of the specimen by Pfannestiel incision.

She was discharged the day after and has been asymptomatic since then.

The histology report was accessory spleen of 3.5 cm with a long vascular pedicle.

DIAGNOSTIC: Accessory wandering spleen.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:https://doi.org/10.1016/j.ciresp.2022.06.017.