



IMAGE OF THE MONTH

Endoscopic closure of a duodenal perforation caused by early migration of a biliary plastic stent with an over-the-scope-clip

Cierre endoscópico con un clip de una perforación duodenal causada por la migración temprana de un stent biliar de plástico

Sónia Bernardo*, Carlos Freitas, João Lopes, Carlos Noronha Ferreira, Rui Tato Marinho

Serviço de Gastroenterologia e Hepatologia, Hospital de Santa Maria, Centro Hospitalar Universitário Lisboa Norte, Lisbon, Portugal

A 52-year-old woman underwent elective endoscopic retrograde cholangiopancreatography for treatment of an infra-hilar main bile duct (MBD) stenosis and fistula from the left hepatic duct due to a cholecystectomy performed one month prior. After dilatation with a 10Fr Savary bougie, a plastic stent (10Fr, 9 cm) was placed across the stenotic segment with the proximal tip in the right hepatic duct. Eight hours later, the patient was admitted with abdominal pain and vomiting. Physical examination was positive for a painful abdomen with no tenderness. Blood tests revealed: C-reactive protein – 44.6 mg/dl and procalcitonin – 4.66 ng/ml. Urgent abdominal computed tomography showed pneumoretroperitoneum, the proximal end of the stent at the distal MBD and the distal end in the retroperitoneum, with presence of fluid and extra-luminal gas (Fig. 1). Urgent esophagogastroduodenoscopy confirmed stent migration to the second portion of duodenum and penetration through the duodenal wall. The plastic stent was removed using rat-tooth forceps, revealing a fistula orifice measuring 3 mm in the duodenal wall (Fig. 2a and b). The perforation was closed with an over-the-scope-clip (OTSC)



Figure 1 Urgent abdominal computed tomography showing pneumoretroperitoneum, the proximal end of the stent at the distal MBD and the distal end in the retroperitoneum, with presence of fluid and extra-luminal gas bubbles.

and the final opacification showed no extravasation of contrast (Fig. 3). However, there was a late failure of OTSC with need of surgical drainage of retroperitoneal abscesses and fistula close. Plastic stent migration with duodenal perforation is a rare but serious condition (3–5%), occurring usually within 2 weeks after the procedure.¹ To our knowledge only 2 cases of duodenal closure using an OTSC were described.^{2,3}

* Corresponding author.

E-mail address: soniamcb@hotmail.com (S. Bernardo).




Figure 2 (a) Urgent esophagogastroduodenoscopy showing stent migration to the second portion of duodenum and penetration through the duodenal wall. (b) Fistula orifice in the duodenal wall after stent extraction.



Figure 3 Final opacification showing no extravasation of contrast after perforation closure with an over-the-scope-clip.

Author contributions

All authors contributed equally to this report.

Conflicts of interest and source funding

None declared.

References

1. Kim HS, Moon HJ, Lee NY, Lee HT, Woo MS, Lee JW, et al. Endoscopic management of duodenal perforations caused by migrated biliary plastic stents. *Endosc Int Open*. 2019;7: E792–5.
2. Le Mouel JP, Hakim S, Thiebault H. Duodenal perforation caused by early migration of a biliary plastic stent: closure with over-the-scope clip. *Clin Gastroenterol Hepatol*. 2019;17: e6–7.
3. Roses LL, Ramirez AG, Seco AL, Blanco ES, Alonso DI, Avila S, et al. Clip closure of a duodenal perforation secondary to a biliary stent. *Gastrointest Endosc*. 2000;51:487–9.