




# Allergología et immunopathología

Sociedad Española de Inmunología Clínica,  
Alergología y Asma Pediátrica

[www.elsevier.es/ai](http://www.elsevier.es/ai)



## ORIGINAL ARTICLE

# Imbalance of serum IL-10 and TGF- $\beta$ in patients with pollen food syndrome

S. Ma, J. Yin \*

Department of Allergy, Peking Union Medical College Hospital, Peking Union Medical College & Chinese Academy of Medical Sciences, Beijing 100730, China

Received 10 October 2012; accepted 2 January 2013

Available online 6 May 2013

### KEYWORDS

Food allergy;  
Pollinosis;  
Pollen food  
syndrome;  
IL-10;  
TGF- $\beta$

### Abstract

**Background:** Pollen food syndrome is one of the main causes of food allergies in adults. However, the intrinsic immunological mechanisms remain unclear.

**Methods:** Forty pollinosis sufferers [23 with a food allergy (PSFA) and 17 without a food allergy (PS)] and 17 non-atopic healthy controls were included in this study. The PSFA group was subdivided into an oral allergy syndrome group, a systemic reaction group, and an anaphylactic reaction group according to their symptoms after eating the suspected foods. Serum IL-10 and TGF- $\beta$  levels of all participants were determined by ELISA. Clinical characteristics of the patients were also evaluated.

**Results:** There were no significant differences in age, sex, pollen-associated symptoms, duration of respiratory disease, and positive parental history of atopy between the PSFA and PS groups. Compared to healthy controls, serum IL-10 levels of both the PSFA group and PS group were significantly lower ( $p \leq 0.01$ ), but TGF  $\beta$  levels were significantly higher in the PSFA group ( $35.3 \pm 5.6$  ng/ml vs.  $31.2 \pm 6.6$  ng/ml, respectively;  $p = 0.037$ ). Within the PSFA group, IL-10 levels in the anaphylactic reaction subgroup were significantly lower compared to oral allergy syndrome subgroup ( $1.87 \pm 0.47$  pg/ml vs.  $1.40 \pm 0.30$  pg/ml, respectively;  $p = 0.027$ ). More severe food allergy symptoms were associated with lower serum IL-10 levels. In contrast, the highest serum levels of TGF- $\beta$  were found in patients from the anaphylactic reaction subgroup.

**Conclusions:** With the exception of a defect in regulatory cells represented by the reduction of IL-10, other potential immunological mechanisms (e.g., Th17 or IL-23 together with TGF- $\beta$ ) may be involved in the development of pollen food syndrome.

© 2012 SEICAP. Published by Elsevier España, S.L. All rights reserved.

**Abbreviations:** PS, pollinosis sufferer without food allergy; PSFA, pollinosis sufferer with food allergy.

\* Corresponding author.

E-mail address: [doctoryinja@163.com](mailto:doctoryinja@163.com) (J. Yin).

## Introduction

The prevalence of food allergy is increasing throughout the world. Food-induced allergic reactions range from mild discomfort restricted to the oral cavity, to systemic and severe symptoms, such as anaphylaxis. Foods are the most common triggers of severe anaphylaxis.<sup>1</sup>

There are typically two types of food allergies: one is primary sensitisation, which initiates in the gastrointestinal tract, and the other is secondary sensitisation, which is a result of cross-reactivity to inhalant allergens. Primary sensitisation is most frequently seen in children and is usually the first manifestation of the atopic march (ranging from food allergy-related eczema/urticaria to allergic rhinitis and then asthma). Pollen is the most common cross-reactive inhalant allergen involved in plant food allergies.

Pollen-related food allergy, also called 'pollen food syndrome', is very common, and even represents the predominant type of food allergy in some countries.<sup>2,3</sup> It was first described by Ortolani et al. in 1988 who noticed that the symptoms were limited to oral allergy reactions.<sup>4</sup> Thereafter, it became known that oral discomfort after consumption of fruits or vegetables may be just a premonition to a variety of generalised symptoms, including life-threatening anaphylaxis.

The extensive presence of pan-allergens (e.g., PR-10s, nsLTPs, profilins, and others) from taxonomically-related or unrelated species sharing highly preserved structures and functions accounts for the high incidence of cross-reactivity between pollens and plant foods.<sup>5</sup> An accurate prevalence of pollen food syndrome is difficult to estimate, because many people who suffer from mild discomfort after eating the implicated food may simply avoid it instead of reporting the symptoms to a physician. According to previous reports, approximately 40–70% of pollinosis sufferers develop a relevant plant food allergy.<sup>6–9</sup> These findings suggest that a substantial number of patients who are allergic to pollen do not develop a food allergy later in life. There may also be some genetic or immunological differences between patients with hay fever with and without pollen food syndrome. Cuesta-Herranz et al. previously compared the clinical and laboratory characteristics of pollen-allergic patients with and without a food allergy. They found that pollen-allergic patients with an associated plant food allergy had a higher frequency of asthma, higher frequency of polysensitisation, and a different pattern of specific IgEs to several kinds of pollen allergens. In addition, patients sensitised to pollen from *Platanus*, *Platago*, *Artemisia*, *Betula*, *Parietaria*, and *Salsola* were predisposed to the development of a plant food allergy.<sup>6</sup> Peach was found to have the highest frequency of allergy among the relevant plant foods in their study.

Recently, the close relationship between regulatory T cells and oral tolerance and food allergy has been well characterised.<sup>10–13</sup> There are two types of regulatory T cells: natural regulatory T cells (nTreg), which are generated in the thymus expressing high level of Foxp3, and inducible T cells (iTreg), which are converted from naïve T cells in the periphery. The latter includes at least two subtypes: IL-10-secreting type 1 regulatory T cells (Tr1)<sup>14</sup> and TGF- $\beta$ -producing Th3 regulatory T cells.<sup>15</sup> IL-10 and TGF- $\beta$  are the

main effector molecules of regulatory T cells and play an important role in the maintenance of immune homeostasis throughout the entire body.

This study aimed to determine serum levels of IL-10 and TGF $\beta$  in pollinosis patients with and without a food allergy and to investigate their potential relationship with the development of pollen food syndrome.

## Materials and methods

### Subjects

Forty patients suffering from *Artemisia* pollinosis and/or birch pollinosis [23 with a food allergy (PSFA) and 17 without a food allergy (PS)], and 17 non-atopic healthy controls were included in this study. PSFA patients were subdivided into an oral allergy syndrome group ( $n=8$ ), systemic reaction group ( $n=6$ ), and anaphylactic reaction group ( $n=9$ ) according to the severity of symptoms after consuming the suspected foods. Clinical characteristics of the patients, including age, sex, pollen-associated symptoms, food-related symptoms, duration of respiratory disease, and positive parental history of atopy, were also evaluated.

Anaphylactic reaction, namely anaphylaxis, was defined as rapid-onset, multisystem, severe, or even life-threatening reactions occurring upon allergen exposure based on World Allergy Organization (WAO) guidelines.<sup>16</sup> Systemic reactions showing symptoms that were restricted to only one system, such as respiratory, gastrointestinal, or skin/mucosa, did not meet the diagnosis criteria of anaphylaxis.

This research protocol was approved by the Ethical Committee of Peking Union Medical College Hospital. Written informed consent was obtained from each participant.

### Inclusion criteria

Inclusion criteria included seasonal symptoms of allergic rhinitis and/or asthma; confirmed sensitisation to birch and/or mugwort pollen by a serum-specific IgE test and skin prick test; plant food allergy diagnosed on the basis of compelling history, including repetitive adverse reactions to certain plant foods and no occurrence of relevant discomfort after complete avoidance of the same food; and a positive serum-specific IgE test against the suspected food.

### Exclusion criteria

All patients included in this study were generally healthy. Patients with serious cardiac and vascular diseases, liver problems, renal dysfunction, mental disorders, tumours, autoimmune diseases, and infectious diseases as well as other types of atopic disease, such as eczema or chronic urticaria, were excluded. Patients who had ever received immunotherapy before recruitment onto the study were also excluded. No glucocorticoids or analogues were allowed one month prior to blood collection.

**Table 1** Clinical characteristics of patients included in this study.

Characteristics	PS (n = 17)	PSFA (n = 23)	p value
<i>Male no. (%)</i>	9 (52.9)	11 (47.8)	0.749
<i>Age, mean ± SD</i>	27.2 ± 14.1	24.3 ± 13.3	0.497
<i>Positive family history of atopy, no. (%)</i>	6 (35.3)	13 (56.5)	0.184
<i>Total IgE (CAP) (kUA/L), median (range)</i>	341 (69–1817)	275 (26.5–2322)	0.594
<i>Duration of respiratory disease (month) median (range)</i>			
Rhinitis	72 (12–360)	48 (24–360)	0.658
Asthma	18 (1–360)	36 (2–84)	0.720
<i>Seasonal respiratory symptoms</i>			0.822
Rhinitis	9 (52.9)	13 (56.5)	
Rhinitis and asthma	8 (47.1)	10 (43.5)	
<i>Pollen involved</i>			
Artemisia	15 (88.2)	17 (73.9%)	0.263
Birch	8 (47.1)	16 (69.6%)	0.151

## Sample preparation

Four millilitres of peripheral venous blood were collected from each subject. Serum was isolated by centrifuging the samples at 3900 rpm for 5 min after allowing them to stand at room temperature for 30 min. After centrifugation the samples were frozen at –80 °C for further analysis.

## Serum analysis

Serum IL-10 and TGF-β levels of all participants were detected using a commercial ELISA kit (eBioscience, USA). The detection limitation was 0.39 pg/ml for IL-10 and 63 pg/ml for TGF-β. The sensitivity of the two kits were 0.05 pg/ml and 15 pg/ml, respectively. Total IgE and specific IgE levels were detected by ImmunoCAP system (Thermo Fisher, Uppsala, Sweden).

## Statistical analysis

The statistical software SPSS 16.0 (SPSS, Inc., Chicago, IL, USA) was used to analyse the data. Frequencies of the qualitative variables were compared by Pearson's chi-squared test or Fisher's exact test. The comparison of quantitative variables was carried out using the Student's *t*-test in case of normally distributed data and the Mann–Whitney *U*-test in case of non-normally distributed data. A *p* value < 0.05 was considered statistically significant.

## Results

All 40 patients with hay fever were from northern China. Birch and Artemisia pollen was the most common trigger of pollinosis in spring and late summer/early autumn, respectively. The clinical characteristics of the patients are shown in Table 1. No significant difference was found in sex, age, family history of atopy, serum total IgE levels, duration of respiratory disease, or pollen-related respiratory symptoms between the PS and PSFA groups. The average age of healthy controls was 27 ± 15.1 years, which included 11 females

and six males. There were no differences in the age and sex ratios between healthy controls (HC) and PS or PSFA patients.

Among the total 40 pollinosis patients, seven were only allergic to birch, 16 were only allergic to Artemisia, and 17 were allergic to both. Among the 23 pollinosis patients with a food allergy, six were allergic to birch pollen, seven were allergic to Artemisia pollen, and 11 were allergic to both. The frequencies of Artemisia and birch pollen allergies in the PS group were equal to those in the PSFA group. Among the PS group, nine patients were recruited within the allergy seasons and eight were recruited in the non-allergy seasons. Among the PSFA group, 12 were recruited within the allergy seasons and 11 were recruited in the non-allergy seasons. There was no difference in the percentage of patients recruited within allergy or non-allergy seasons in both groups (Pearson's  $\chi^2$ , *p* = 0.962).

All patients in the PSFA group were allergic to peach (100%), while some of them were also allergic to other plant foods with less frequency, such as apple (69.6%); pear (30.4%); Chinese date (30.4%); apricot (26.1%); mango (21.7%); and hazelnut (21.7%). The sensitisation profiles of the 23 pollinosis sufferers with a plant food allergy are shown in Table 2. All six patients who were only allergic to birch pollen were sensitised to rBet v 1 and its homologue, rPru p 1. Among the seven patients only allergic to Artemisia pollen, all were sensitised to nArt v 3 and rPru p 3. Although a positive IgE response to Birch pollen was identified in three of the patients, none of them were sensitised to the relevant food allergen rPru p 1. The 10 patients allergic to both birch and Artemisia pollen showed a mixed sensitisation profile. Only four patients were simultaneously sensitised to rBet v 1 and nArt v 3 as well as their respective homologues rPru p 1 and rPru p 3. Two patients were sensitised to rBet v 2, nArt v 3, and rPru p 3. Two were only sensitised to rBet v 1 and rPru p 1, and the last two were sensitised to rBet v 1, nArt v 3 and rPru p 3, but did not show positive response to rPru p 1, the homologue of rBet v 1.

Using an IL-10 ELISA kit with high sensitivity, the serum IL-10 levels of only one patient in the PSFA group were lower than the detection limit for the assay. Because the OD value

**Table 2** Basic information and sensitisation profiles of pollinosis sufferers with a plant food allergy.


Pollen	No.	Sex	Age	Specific IgE (kU/L)								
				t3	rBet v 1	rBet v 2	w6	nArt V 1	nArt V 3	peach	rPru p 1	rPru p 3
Birch	1	Female	42	100	100	0	0.18	0	0	11.3	21.8	0
	2	Male	42	33.4	29.8	0.62	0.1	0	0	12.5	49.4	0.98
	3	Female	28	100	100	0	0	0	0.14	13.2	32.8	0.15
	4	Female	44	89.9	70.9	0	0	0	0	8.25	32.6	0
	5	Female	41	24.3	18	0.24	0	0	0	3.26	13.2	0.53
	6	Female	45	77.1	77.7	0	0	0	0	1.76	23.2	0
Birch + Artemisia	7	Male	30	16.1	20.6	0.02	11.9	0.81	16.1	7.32	8.82	1.27
	8	Male	15	2.5	0	3.55	100	80.7	100	24.2	0	10.2
	9	Male	9	19.9	20.1	0	4.18	4.15	0	4.08	8.01	0
	10	Female	49	53.5	47.7	0	2.34	0	6.76	10.1	19.5	2.43
	11	Female	19	11.6	0	10.1	71.2	22.1	68.8	3.69	0	0.73
	12	Male	9	39.7	38.9	0.14	50.8	18.9	27.6	25.2	19.7	9.02
	13	Female	16	0.77	0.55	0	39.8	17.9	34.6	3.24	0.25	4.54
	14	Male	8	2.42	1.62	0.04	100	100	100	96.3	0.07	88.9
	15	Male	13	90.6	74.8	6.4	100	75.8	100	18.8	9.98	13.4
	16	Male	19	24.6	21.7	0.01	36.5	38.2	0.07	4.7	10.7	0
Artemisia	17	Male	15	0.04	0.01	0	4.03	0.06	6.89	2.44	0	1.95
	18	Male	25	0.82	0.02	1.08	21.6	5.5	25	2.01	0.01	0.59
	19	Female	23	0	0	0	13.1	0.33	14.2	9.46	0	9.27
	20	Female	19	1.26	0.01	2.02	72	14.5	100	21.2	0	9.04
	21	Female	17	0.03	0	0	35.06	0	32.58	18.4	0	21.4
	22	Male	10	25.2	0.62	9.3	100	10	100	50.2	0.03	10.2
	23	Female	23	0.07	0.01	0.01	89.3	50.4	87.5	2.33	0	1.42

for this sample was very close to the detection limit, we also calculated the value using the formula deduced from the standard curve. The serum levels of TGF- $\beta$  were detectable in all patients and healthy controls.


No differences in serum levels of IL-10 or TGF- $\beta$  were found among patients allergic to different types of pollen (Figs. 1 and 2) or patients recruited at different times (Fig. 3). In both the PS and PSFA groups, no significant

differences were found in serum levels of IL-10 and TGF- $\beta$  between patients with and without asthmatic symptoms ( $p > 0.05$ ).


Compared to healthy controls, the serum IL-10 levels of both the PSFA and PS groups were significantly lower ( $p \leq 0.01$ ), but TGF- $\beta$  was significantly higher only in the PSFA group ( $35.3 \pm 5.6$  ng/ml vs.  $31.2 \pm 6.6$  ng/ml, respectively;  $p = 0.037$ ). No significant difference was found in IL-10




**Figure 1** Comparison of serum levels of (A) TGF- $\beta$  and (B) IL-10 among patients allergic to different types of pollen (within a total of 40 pollinosis patients).



**Figure 2** Comparison of serum levels of (A) TGF- $\beta$  and (B) IL-10 among patients allergic to different types of pollen (within 23 pollinosis patients with a food allergy).



**Figure 3** Comparison of serum levels of (A) TGF- $\beta$  and (B) IL-10 among patients recruited within or out of pollen seasons.



**Figure 4** (A) Serum levels of IL-10 and TGF- $\beta$  in healthy controls and pollinosis patients with or without a food allergy and (B) serum levels of IL-10 and TGF- $\beta$  of patients with varying severity of food allergy (OAS = oral allergy syndrome; SS = systemic symptom; AR = anaphylactic reaction).

levels between the PSFA and PS groups ( $1.66 \pm 0.60$  pg/ml vs.  $1.29 \pm 0.50$  pg/ml, respectively;  $p = 0.07$ ), but serum TGF- $\beta$  levels in the PSFA group were markedly higher than those in the PS group ( $35.3 \pm 5.6$  ng/ml vs.  $30.6 \pm 4.8$  ng/ml, respectively;  $p = 0.008$ ). Within the PSFA group, IL-10 levels in the anaphylactic reaction subgroup were significantly lower compared to the oral allergy syndrome subgroup ( $1.87 \pm 0.47$  pg/ml vs.  $1.40 \pm 0.30$  pg/ml, respectively;  $p = 0.027$ ). We found that more severe food allergy symptoms corresponded to lower serum IL-10 levels. However, in contrast, the highest serum TGF- $\beta$  levels were observed in the anaphylactic reaction subgroup (Fig. 4).

## Discussion

Pollen food allergy syndrome is a main cause of adult food allergies in many countries, and food-associated symptoms are often serious and perennial.<sup>17</sup> Therefore, this condition substantially affects the quality of life of these sufferers. This study investigated the intrinsic mechanisms of pollen food syndrome, which may help identify a useful preventive and therapeutic approach for this burdensome disorder. Peach is one of the most common allergenic fruits involved in pollen food syndrome, and its component allergens, Pru p 1, Pru p 3, and Pru p 4, are all cross-reactive and belong to the PR-10, nsLTP, and Profilin families, respectively. Thus, peach was chosen to be a model food in this study to analyse the potential cross-reactivity between pollens and foods. As shown in Table 2, birch pollinosis patients with a peach allergy were usually sensitised to rBet v 1 and its homologous allergen rPru p 1, and patients allergic to mugwort pollen and peach were simultaneously sensitised to nArt v 3 and its homologue rPru p 3. Moreover, patients allergic to both birch and mugwort pollen displayed mixed sensitisation patterns of peach.

Our data showed that no differences in age, sex, family history of atopy, serum total IgE levels, ratio of patients with asthmatic symptoms, duration of pollen-related respiratory disease, or type of allergenic pollen were found between the PS and PSFA groups in this study. However, serum levels of IL-10 and TGF- $\beta$  were significantly different between the groups. Based on the severity of the food allergy symptoms, the levels of the two cytokines varied inversely; that is, patients with more severe symptoms had higher TGF- $\beta$  levels and lower IL-10 levels. Serum IL-10 and TGF- $\beta$  levels are the most important effector molecules of regulatory T cells, which play a crucial role in maintaining the immunological balance throughout the body.

Although pollen food syndrome is different from a type 1 food allergy, which is primarily sensitised in the gut, the more important triggering phase of both types of food allergy occurs in the gastrointestinal tract. The immune environment of the gut is also crucial for the development and subsequent clinical manifestations of pollen-related food-induced disorders.

Normally, the gastrointestinal tract remains in a state of immunological homeostasis. Immunocytes and immune factors present in the gastrointestinal tract react correctly to the pathogens and tolerate food antigens and commensal bacteria well. All types of regulatory T cells are involved in the maintenance of immune balance of the gut.

Foxp3+ nTregs generated in the thymus can migrate to other tissues, including the gastrointestinal tract, to exert suppressive function with the help of a spectrum of homing receptors, such as gut-homing integrin and  $\alpha 4\beta 7$ . The iTregs generated in the gut, such as Tr1 and Th3T cells, could also enter the circulation and reach other organs to help maintain immune tolerance in those organs. This phenomenon is called 'oral tolerance', which has been shown to have the potential to prevent and treat autoimmune disease.<sup>13</sup>

TGF- $\beta$ , a powerful immunosuppressive cytokine essential for the establishment of oral tolerance, is abundantly secreted by gut immunocytes upon oral antigen exposure. It can induce the expression of Foxp3, which is the key gene controlling the development and functioning of nTregs,<sup>18</sup> and can convert the conventional CD4+CD25 $-$  T cells to CD4+CD25 $+$  regulatory T cells.<sup>19,20</sup> Moreover, TGF- $\beta$  can inhibit the expression of the lineage-specific transcription factors T-bet and GATA-3 thereby blocking the differentiation of Th1 and Th2 cells, respectively.<sup>21</sup> The contribution of TGF- $\beta$  in maintaining the homeostasis of the gut is also displayed in its capacity to promote IgA class switching.<sup>22</sup>

In addition, TGF- $\beta$  is a profibrotic cytokine that is crucial for airway wound repair. However, excessive activation of TGF- $\beta$  can promote airway fibrosis and ultimately leads to airway remodelling.<sup>23</sup> Protein and mRNA gene expression of TGF- $\beta$  are generally increased in asthmatic patients. In our study, the duration of respiratory diseases and number of patients with asthma in the PS group was comparable to that in the PSFA group. Therefore, the impact of these factors on the findings of this study was excluded from the analysis. No difference was found in TGF- $\beta$  levels between patients with only seasonal rhinitis or patients with both rhinitis and asthma, which could be due to the fact that all patients appeared healthy and only presented pollen-related symptoms during limited periods of the year. The actual duration of symptoms and inflammatory status of the airway is quite shorter than that of the patients with persistent rhinitis and/or asthma.

IL-10 is another multifunctional modulatory cytokine.<sup>24</sup> IL-10 can downregulate the expression of the IgE receptor and its signal molecules, including Syk, Fyn, and Akt, which alleviates the inflammatory reaction caused by IgE. It can also induce the differentiation of naïve T cells to Tr1 cells and maintain the expression of Foxp3 in nTregs. In addition, IL-10 is capable of promoting the generation of IgG4, which is always thought to be a blocking antibody that represents a state of tolerance.

Previous studies have suggested that gene polymorphisms and peripheral expression of IL-10 are associated with food allergies.<sup>25,26</sup> We previously found that a single nucleotide polymorphism in the *IL-10* gene was associated with the occurrence of anaphylaxis (data not published). Moreover, in a mouse model of food allergy, Frossard et al. showed that exogenous supplementation of IL-10 decreased food-induced anaphylaxis and prevented IgE-type sensitisation to common food allergens.<sup>27</sup> Taken together, these data support the finding that a reduction of serum IL-10 levels is related to the occurrence and development of pollen-related food allergies.

The most interesting finding of our study was that the serum TGF- $\beta$  levels in pollinosis sufferers with a food allergy were abnormally elevated, and the highest levels of

TGF- $\beta$  appeared in the group with the most serious symptoms. Previous studies have presented controversial conclusions with respect to the role of TGF- $\beta$  in anaphylaxis. Okamoto et al. found that oral administration of high dose TGF- $\beta$  can suppress a food-induced IgE response and systemic anaphylaxis,<sup>28</sup> whereas an earlier study conducted by Kim et al. showed that TGF- $\beta$  was key for activating mast cells and the development of anaphylaxis.<sup>29</sup> Therefore, it is likely that an optimal level of TGF- $\beta$  may be beneficial, whereas high and low levels could be harmful. However, TGF- $\beta$  could also be pathogenic in some inflammatory diseases, such as autoimmune diseases and colitis, when combined with pro-inflammatory cytokines (e.g., IL-6 or IL-23) by promoting the generation of Th17 cells.<sup>30,31</sup> The combination of TGF- $\beta$  and IL-6 can also prevent the generation of Foxp3 cells,<sup>32</sup> which might explain why higher serum TGF- $\beta$  levels were associated with more severe food allergy symptoms. In addition, IL-23 can restrain regulatory T cell activity and suppress IL-10 function, thereby breaking the immunological homeostasis of the gut.<sup>33,34</sup> Thus, further assessment of IL-6 or IL-23 levels might help us better understand the pathogenesis of food pollen allergies.

This was a preliminary study with some limitations. First, the study sample was relatively small. Second, patients allergic to different pollens in different seasons were recruited, and therefore only a few people were monosensitised to only one type of pollen. However, we carefully compared the serum levels of IL-10 and TGF- $\beta$  among patients allergic to different types of pollen and between patients enrolled during different allergy seasons, and no differences were found. Thus, the results of our study were objective and convincing.

In summary, our data indicate that with the exception of a defect in regulatory cells represented by the reduction of IL-10, there may be potential intrinsic mechanisms (e.g., Th17 or IL-23 together with TGF- $\beta$ ) that are involved in the development of pollen food allergies.

## Ethical disclosures

**Protection of human subjects and animals in research.** Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this investigation.

**Patients' data protection.** Confidentiality of data. The authors declare that they have followed the protocols of their work centre on the publication of patient data and that all the patients included in the study have received sufficient information and have given their informed consent in writing to participate in that study.

**Right to privacy and informed consent.** Right to privacy and informed consent. The authors have obtained the informed consent of the patients and/or subjects mentioned in the article. The author for correspondence is in possession of this document.

## Conflict of interests

The authors have no conflict of interests.

## References

- Huang F, Chawla K, Jarvinen KM, Nowak-Wegrzyn A. Anaphylaxis in a New York City pediatric emergency department: Triggers, treatments, and outcomes. *J Allergy Clin Immunol*. 2012;129:162–8.
- Skypala IJ, Calderon MA, Leeds AR, Emery P, Till SJ, Durham SR. Development and validation of a structured questionnaire for the diagnosis of oral allergy syndrome in subjects with seasonal allergic rhinitis during the UK birch pollen season. *Clin Exp Allergy*. 2011;41:1001–11.
- Asero R, Antonicelli L, Arena A, Bommarito L, Caruso B, Crivellaro M, et al. EpidemAAITO: features of food allergy in Italian adults attending allergy clinics: a multi-centre study. *Clin Exp Allergy*. 2009;39:547–55.
- Ortolani C, Ispano M, Pastorello E, Bigi A, Ansaldi R. The oral allergy syndrome. *Ann Allergy*. 1988;61:47–52.
- Hauser M, Roulias A, Ferreira F, Egger M. Panallergens and their impact on the allergic patient. *Allergy Asthma Clin Immunol*. 2010;6:1.
- Cuesta-Herranz J, Barber D, Blanco C, Cistero-Bahíma A, Crespo JF, Fernández-Rivas M, et al. Differences among pollen-allergic patients with and without plant food allergy. *Int Arch Allergy Immunol*. 2010;153:182–92.
- Amlot PL, Kemeny DM, Zachary C, Parkes P, Lessof MH. Oral allergy syndrome (OAS): symptoms of IgE-mediated hypersensitivity to foods. *Clin Allergy*. 1987;17:33–42.
- Dreborg S, Foucard T. Allergy to apple, carrot and potato in children with birch pollen allergy. *Allergy*. 1983;38:167–72.
- Cuesta-Herranz J, Lázaro M, Figueredo E, Igea JM, Umpiérrez A, De-Las-Heras M. Allergy to plant-derived fresh foods in a birch- and ragweed-free area. *Clin Exp Allergy*. 2000;30:1411–6.
- Battaglia M, Gianfrani C, Gregori S, Roncarolo MG. IL-10-producing T regulatory type 1 cells and oral tolerance. *Ann NY Acad Sci*. 2004;1029:142–53.
- Chehade M, Mayer L. Oral tolerance and its relation to food hypersensitivities. *J Allergy Clin Immunol*. 2005;115(3-12):13.
- Burks AW, Laubach S, Jones SM. Oral tolerance, food allergy, and immunotherapy: implications for future treatment. *J Allergy Clin Immunol*. 2008;121:1344–50.
- Tsuji NM, Kosaka A. Oral tolerance: intestinal homeostasis and antigen-specific regulatory T cells. *Trends Immunol*. 2008;29:532–40.
- Wu K, Bi Y, Sun K, Wang C. IL-10-producing type 1 regulatory T cells and allergy. *Cell Mol Immunol*. 2007;4:269–75.
- Weiner HL. Induction and mechanism of action of transforming growth factor-beta-secreting Th3 regulatory cells. *Immunol Rev*. 2001;182:207–14.
- Simons FE, Arduoso LR, Bilò MB, Dimov V, Ebisawa M, El-Gamal YM, et al. 2012 Update: World Allergy Organization Guidelines for the assessment and management of anaphylaxis. *Curr Opin Allergy Clin Immunol*. 2012;12:389–99.
- Geroldinger-Simic M, Zelniker T, Aberer W, Ebner C, Egger C, Greiderer A, et al. Birch pollen-related food allergy: clinical aspects and the role of allergen-specific IgE and IgG4 antibodies. *J Allergy Clin Immunol*. 2011;127:616–22.
- Fontenot JD, Gavin MA, Rudensky AY. Foxp3 programs the development and function of CD4+CD25+ regulatory T cells. *Nat Immunol*. 2003;4:330–6.
- Pyzik M, Piccirillo CA. TGF-beta1 modulates Foxp3 expression and regulatory activity in distinct CD4+ T cell subsets. *J Leukoc Biol*. 2007;82:335–46.
- Zheng SG, Wang JH, Gray JD, Soucier H, Horwitz DA. Natural and induced CD4+CD25+ cells educate CD4+CD25– cells to develop suppressive activity: the role of IL-2, TGF-beta, and IL-10. *J Immunol*. 2004;172:5213–21.

21. Wan YY, Flavell RA. Regulatory T cells, transforming growth factor-beta, and immune suppression. *Proc Am Thorac Soc*. 2007;4:271–6.
22. Macpherson AJ, Slack E. The functional interactions of commensal bacteria with intestinal secretory IgA. *Curr Opin Gastroenterol*. 2007;23:673–8.
23. Halwani R, Al-Muhsen S, Al-Jahdali H, Hamid Q. Role of transforming growth factor-beta in airway remodeling in asthma. *Am J Respir Cell Mol Biol*. 2011;44:127–33.
24. Sabat R, Grutz G, Warszawska K, Kirsch S, Witte E, Wolk K. Biology of interleukin-10. *Cytokine Growth Factor Rev*. 2010;21:331–44.
25. Campos Alberto EJ, Shimojo N, Suzuki Y, Mashimo Y, Arima T, Matsuura T, et al. IL-10 gene polymorphism, but not TGF-beta1 gene polymorphisms, is associated with food allergy in a Japanese population. *Pediatr Allergy Immunol*. 2008;19:716–21.
26. Krogulska A, Borowiec M, Polakowska E, Dynowski J, Mlynarski W, Wasowska-Królikowska K, et al. FOXP3, IL-10, and TGF-beta genes expression in children with IgE-dependent food allergy. *J Clin Immunol*. 2011;31:205–15.
27. Frossard CP, Steidler L, Eigenmann PA. Oral administration of an IL-10-secreting *Lactococcus lactis* strain prevents food-induced IgE sensitization. *J Allergy Clin Immunol*. 2007;119:952–9.
28. Okamoto A, Kawamura T, Kanbe K, Kanamaru Y, Ogawa H, Okumura K, et al. Suppression of serum IgE response and systemic anaphylaxis in a food allergy model by orally administered high-dose TGF-beta. *Int Immunopharmacol*. 2005;17:705–12.
29. Kim HM, Lee YM. Role of TGF-beta 1 on the IgE-dependent anaphylaxis reaction. *J Immunol*. 1999;162:4960–5.
30. Feng T, Qin H, Wang L, Benveniste EN, Elson CO, Cong Y, et al. Th17 cells induce colitis and promote Th1 cell responses through IL-17 induction of innate IL-12 and IL-23 production. *J Immunol*. 2011;186:6313–8.
31. Pernis AB. Th17 cells in rheumatoid arthritis and systemic lupus erythematosus. *J Intern Med*. 2009;265:644–52.
32. Bettelli E, Carrier Y, Gao W, Korn T, Strom TB, Oukka M, et al. Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature*. 2006;441:235–8.
33. Izcue A, Hue S, Buonocore S, Arancibia-Cárcamo CV, Ahern PP, Iwakura Y, et al. Interleukin-23 restrains regulatory T cell activity to drive T cell-dependent colitis. *Immunity*. 2008;28:559–70.
34. Liu Z, Feng BS, Yang SB, Chen X, Su J, Yang PC. Interleukin (IL)-23 suppresses IL-10 in inflammatory bowel disease. *J Biol Chem*. 2012;287:3591–7.