



ORIGINAL ARTICLE

Insect deterrent and nematicidal screening of microfungi from Mexico and anti-aphid compounds from *Gliomastix masseei*

Ana L. Ruiz-Jiménez^a, Azucena González-Coloma^b, Mari Fe Andrés-Yeves^b, Esaú Ruiz-Sánchez^c, Gabriela Heredia^d, Sergio R. Peraza-Sánchez^a, Irma L. Medina-Baizabal^a, Manuela Reyes-Estebanez^e, Blondy Canto-Canché^a, Marcela Gamboa-Angulo^{a,*}



CrossMark

^a Unidad de Biotecnología, Centro de Investigación Científica de Yucatán, AC, Mérida, Yucatán, Mexico

^b Departamento de Ciencias Agrícolas-CCMA, CSIC, Madrid, Spain

^c Instituto Tecnológico de Conkal, Conkal, Yucatán, Mexico

^d Departamento de Biología de Suelos, Instituto de Ecología, Xalapa, Veracruz, Mexico

^e Departamento de Microbiología Ambiental y Biotecnología, Universidad Autónoma de Campeche, Campeche, Mexico

Received 20 October 2015; accepted 12 August 2016

Available online 1 March 2017

KEYWORDS

Clonostachys rosea;
Fatty acids;
Gliomastix masseei;
Myzus persicae;
Rhopalosiphum padi

Abstract Fungal metabolites are promising alternatives for the development of biorational pesticides. In this sense, microfungi from tropical regions are valuable sources of natural compounds for pest management. With the aim of broadening the search for new eco-friendly products to manage plant pests, this study was carried out to evaluate the biological activity of 23 tropical fungal extracts on three species of phytophagous insects and a plant parasitic nematode. In addition, the active principles of the most effective extract were identified. The insect deterrent activity of fungal extracts was evaluated on the settling of aphids *Myzus persicae* and *Rhopalosiphum padi*, and on the feeding of lepidoptera larva *Spodoptera littoralis*; the nematostatic activity was evaluated on the mobility of *Meloidogyne javanica*. Active metabolites from *Gliomastix masseei* were identified by GC-MS techniques and by comparison with commercial standards. Results showed seven extracts with strong effect on the settling of *M. persicae* and *R. padi* (settling inhibition >80%). The calculated median of effective concentration (EC_{50}) values ranged from 8 to 38 $\mu\text{g}/\text{cm}^2$ for the extracts of *Clonostachys rosea* and *G. masseei*, respectively. Bioassay-guided separation of the ethyl acetate extract of *G. masseei*

* Corresponding author.

E-mail address: mmarcela@cicy.mx (M. Gamboa-Angulo).

revealed the presence of fatty acids and their derivatives, where methyl 9-octadecenoate was the most active compound with EC₅₀ values of 16 µg and 35 µg/cm² for *M. persicae* and *R. padi*, respectively. Extracts of *C. rosea* and *G. masseei* could be a promising option in the control of pest aphids in agriculture.

© 2016 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Clonostachys rosea;
Ácidos grasos;
Gliomastix masseei;
Myzus persicae;
Rhopalosiphum padi

Evaluación de la actividad disuasiva de insectos y nematotóxica de microhongos de México y compuestos anti-áfidos de *Gliomastix masseei*

Resumen Los metabolitos fúngicos son agentes prometedores para el desarrollo de plaguicidas biorracionales. En este sentido, los hongos microscópicos de zonas tropicales representan una valiosa fuente de compuestos naturales para el manejo de plagas. Con la finalidad de ampliar la investigación en productos amigables con el medio ambiente, en este estudio se evaluó la actividad biológica de 23 extractos de hongos sobre 3 especies de insectos fitófagos y un nematodo fitoparásito; además se identificaron los componentes del extracto más activo. El efecto disuasivo de los extractos fúngicos se evaluó en el asentamiento de *Myzus persicae* y *Rhopalosiphum padi*, así como en la alimentación de *Spodoptera littoralis*; la actividad nematostática se evaluó sobre la movilidad de *Meloidogyne javanica*. Los metabolitos activos de *Gliomastix masseei* se identificaron por cromatografía de gases-espectrometría de masas y por comparación con muestras comerciales. Los resultados mostraron 7 extractos con fuerte efecto en la inhibición del asentamiento (> 80%) de *M. persicae* y *R. padi*. Los valores de la concentración efectiva media (CE₅₀) estuvieron en el rango de 8 a 38 µg/cm² para los extractos de *Clonostachys rosea* y *G. masseei*, respectivamente. El extracto de acetato de etilo de *G. masseei* se fraccionó por un proceso biodirigido y reveló la presencia de ácidos grasos y sus derivados, donde el 9-octadecenoato de metilo fue el más activo, con una CE₅₀ de 16 µg/cm² para *M. persicae* y 35 µg/cm² para *R. padi*. Los extractos fúngicos de *C. rosea* y *G. masseei* pueden ser una alternativa promisoria en el control de áfidos que son plaga en la agricultura.

© 2016 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The negative impact of synthetic pesticides on the environment and human health has highlighted the use of natural products as a promising alternative for biorational plant pest management⁷. Fungal secondary metabolites are considered a valuable source of molecules with a wide variety of biological applications. Various reports have pointed out the potential of these compounds for the control of insects and nematodes. One of the first detailed works on the insecticidal properties of fungal metabolites was published by Cole and Rolinson⁹. More recently, Asaff et al.⁴ showed that dipicolinic and oxalic acids, isolated from *Paecilomyces fumosoroseus*, have strong insecticidal effects on nymphs of *Bemisia* species. Likewise, Gaich and Mulzer¹² showed that the sesquiterpenoid, penifulvin A (*Penicillium griseofulvum*) has antifeedant effects on *Spodoptera frugiperda*. In addition, nematotoxic activity has been reported in a wide variety of fungal metabolites, such as cytochalasin B, enniatin B, moniliformin, verrucarin A²⁰, trichodermin^{36,37}, and butyric acid from *Paecilomyces lilacinus* 6029³³.

Our research group has undertaken a long-term project on bioprospection of mycodiversity and its biotechnological application. In this sense, the search for microfungi in the tropical regions of Mexico has yielded a fungal collection whose mycelial extracts have been evaluated on several pathogenic bacteria and fungi, as well as the plant parasitic nematode *Meloidogyne incognita*^{13,29}. None of these native fungi have been assessed against phytophagous insects. Therefore, in the present work, with the aim of broadening the search for new and more eco-friendly agents to control plant pests, twelve saprophytic fungi (Table 1) were selected, based on their biological profile. The fungi were cultured and their organic extracts were evaluated on the root-knot nematode *Meloidogyne javanica* (Neal) Chitwood (Nematoda: Meloidognidae), the polyphagous Lepidoptera *Spodoptera littoralis* Boisduval (Lepidoptera: Noctuidae), and aphids *Myzus persicae* Sulzer (Hemiptera: Aphididae) and *Rhopalosiphum padi* L. (Hemiptera: Aphididae). In addition, one of the most active extracts was partitioned and its active constituents were identified through bioassay-guided fractionation.

Table 1 List of Mexican microfungi tested (organic extracts) on insect deterrent and nematostatic activities

Fungal strains	Key	Order	Site
<i>Cladosporium cladosporioides</i> (Fresen.) G.A. de Vries	XH1E	Capnodiales	1
<i>Clonostachys rosea</i> (Link) Schroers, Samuels, Seifert & W. Gams	TH27	Hypocreales	1
<i>Corynespora cassiicola</i> (Berk. & M.A. Curtis) C.T. Wei	MR01	Pleosporales	1
<i>Cylindrium elongatum</i> Bonord	MR45	Hypocreales	2
<i>Cylindrocarpon congoense</i> J.A. Mey	XH8A	Hypocreales	1
<i>Fusarium</i> sp.	TA54	Hypocreales	1
<i>Gliomastix masseei</i> (Sacc.) Matsush.*	CICY029	Hypocreales	1
<i>Perelegamyces parviechinulatus</i> W.B. Kendr. & R.F. Castañeda	GH25	Incertae sedis	3
<i>Phialophora verrucosa</i> Medlar	MR54	Chaetothyriales	1
<i>Spadicoides</i> sp.	MR52	Sordariales	1
<i>Verticillium</i> sp.	TH28	Hypocreales	1
<i>Volutella</i> sp.	TH22	Hypocreales	1

Site: 1: Yucatán, 2: Veracruz, 3: Tabasco.

* Previously reported as *Gliomastix murorum* MR26²⁸.

Materials and methods

Saprophytic fungi

Fungal strains were obtained from the culture collection of the Unidad de Biotecnología of the Centro de Investigación Científica de Yucatán (**Table 1**) as previously reported by Gamboa-Angulo et al.¹³ and Reyes-Estebanez et al.²⁹.

Culture and extraction of fungi

Fungi were grown in fermented rice (FR) for 40 days at $25 \pm 2^\circ\text{C}$ and in light-dark photoperiod of 12:12 h. Fungal cultures were lyophilized, fragmented and extracted with ethyl acetate three times (50 ml \times 100 g FR) and methanol (50 ml \times 100 g FR, at approximately 50 °C). The solvents were eliminated by reduced pressure in a rotary evaporator. Ethyl acetate (EA) and methanol (ME) extracts were kept in darkness at 4 °C until use²⁹.

Partition and chromatographic fractionation of ethyl acetate extract of *Gliomastix masseei*

Ethyl acetate extract (100 mg) of *G. masseei* was re-suspended in acetonitrile (5 ml) and partitioned with hexane three times (2:1, 1:1, 1:1, v/v). Solvents were eliminated under reduced pressure to obtain fractions of low (ARGM-1A, 21%) and medium (ARGM-1B, 5%) polarity, and a precipitate. Both fractions were sub-fractionated by silica gel chromatographic column and eluted with hexane, ethyl acetate, acetone, and methanol mixtures. Extract and fractions were analyzed by thin layer chromatography (TLC) on silica gel (60F₂₅₄ plates) and detected under UV light and by spraying phosphomolybdic acid as revealing agent.

Preparation of methyl derivatives

Methyl derivatives of fatty acids were prepared in the laboratory. Potassium carbonate (50 mg) re-suspended in acetone was added to the sample (100 mg) and, after 15 min

of continuous stirring, methyl iodide was added (150 µl) and the mixture was left for one night at room temperature. Methylated products were recovered by extraction with ethyl acetate¹⁷.

Analytical methods

Gas chromatography-mass spectrometry (GC-MS) analyses were performed on an Agilent Technologies 6890N chromatograph coupled to an Agilent Technologies 5975B mass selective detector, with Ultra 1 column [methylsiloxane, 25 m long, 0.32 mm i.d., 0.52 µm film thickness, helium at flow rate = 1.0 ml/min, $T_1 = 150^\circ\text{C}$, gradient = 10 °C min, $T_2 = 280^\circ\text{C}$ (20 min)]. Each extract was analyzed using 0.4 µl of sample at 2% (w/v). Most components were identified by matching their mass spectra to those of the corresponding compounds in the database of the equipment (NIST 05) and/or by comparison with commercial standards (Sigma-Aldrich).

Bioassays

Settling inhibition of *M. persicae* and *R. padi*

Evaluations of settling inhibition (SI) were performed on adults of *M. persicae* and *R. padi*. Aphids were grown on pepper (*Capsicum annuum* L.) and barley (*Hordeum vulgare* L.), respectively. Colonies were kept in a growth chamber at 60–70% room humidity, photoperiod of 16:8 h (L:D), and temperatures of $21 \pm 2^\circ\text{C}$.

Bioassays were conducted in plastic boxes (3 cm \times 3 cm \times 1.5 cm) lined at the bottom with 2% agar. Two leaf pieces (ca. 1 cm²) cut from the appropriate host plant were placed on the agar. One piece was coated with 100 µg extract diluted in 10 µl of either acetone or methanol (treated = T) and the other leaf piece was coated only with 10 µl of the respective solvent (control = C). Twenty replicates (plastic boxes) per extract were evaluated, each replicate containing ten aphids. The percentage of aphids settled on each piece of leaf was recorded 24 h after exposure. A settling inhibition index (%SI) was calculated for each extract, %SI = [1 – (%T/%C)] \times 100, where %T and %C

are the percentage of aphids settled on the treated and control leaf pieces, respectively⁶. The dose-response experiments for the most active extracts were carried out with the following serial dilutions: 200, 100, 50, 20, and 4 µg/cm².

Feeding inhibition of *S. littoralis*

S. littoralis larvae were reared on an artificial diet as described by Poitout and Bues²⁶. The colony was kept in a growth chamber at 60–70% room humidity, photoperiod of 16:8 h (L:D), and temperature of 20±2 °C. For the bioassay, sixth instar larvae of *S. littoralis* were used. Two larvae were placed in a Petri dish (9 cm diameter × 1 cm) containing agar (2%). Four wells were prepared and four discs (1 cm²) of fresh bell-pepper leaves were deposited in the wells. Two of the discs were coated with 100 µg of the extracts dissolved in 10 µl acetone or methanol (T), and the other two discs were coated with the solvent only (C). Larvae were starved for 3 h prior to the experiments and subsequently placed on the leaf disc and allowed to consume approximately 75% of the disc area before evaluation. Each treatment was conducted in quintuplicate and repeated two times for the active extracts. The uneaten disc areas were measured with a digital program (Image J Version 1.37r, 2006, <http://rsb.info.nih.gov/ij/>), and the percentage of feeding inhibition was calculated by the equation %FR = [1 – (T/C)] × 100, where T and C are the consumption of treated and control leaf discs, respectively⁶.

Mobility inhibition of *M. javanica*

Egg masses of *M. javanica* were obtained from roots of tomato plants (*Solanum lycopersicum* L.) grown under greenhouse conditions. The eggs were incubated in distilled water at 25 °C for 24 h to obtain second-stage juveniles (J2). Bioassay with J2 was performed in 96-well plates. Fungal extracts were dissolved in DMSO with 0.6% Tween 20 (20 µg/µl). Subsequently, 5 µl of the extracts were deposited in each well and 100–150 J2 in 95 µl of water were added to each well. The well plates were incubated at 25 °C for 72 h. Inhibition of mobility (IM) of J2 was obtained as previously described by Hernández-Carlos et al.¹⁶.

Data analysis

The effect of an extract is considered strong when SI > 70% for aphids¹⁵, FR > 80% for *S. littoralis*¹⁰ and IM > 90% after 48 h exposure for *Meloidogyne* spp.² Effective concentrations (EC₅₀) were calculated by the probit analysis using the Statistical Analysis System (SAS), version 8.1 for Windows. EC₅₀ values were considered significantly different if confidence intervals did not overlap.

Molecular identification of *G. masseei*

Molecular identification of the fungal strain MR-36 was performed as described by Moo-Koh et al.²². The mycelial mass of fungal culture was obtained and the genomic DNA was extracted using the Kit ZR Fungal/Bacterial DNA MiniPrep™. Nuclear ribosomal DNA internal transcribed spacer (ITS) regions (ITS1-5.8S rDNA ITS2) were amplified and sequenced using primers ITS1 and ITS4³⁵. The amplified products

were visualized by electrophoresis in 2% agarose gel (m/v) (Sigma®) and the products of amplification were sequenced in Macrogen (www.macrogen.com). Sequences were edited to eliminate the noise at ends, minus strands were converted into reverse complement, and both strands were aligned. A query of 500 bp was used to compare with those in the database of the Gene Bank of the National Center for Biotechnology Information (<http://blast.ncbi.nlm.gov>), using the BLAST software, and deposited in GenBank (accession number KRO69085).

Results

The results of settling inhibition, feeding reduction, and nematotoxic activities are shown in Table 2. None of the extracts inhibited the mobility of *M. javanica* (MI < 50%) or the feeding of *S. littoralis* (FR < 70%). In contrast, the EA extracts of *G. masseei*, *Perelegamyces parviechinulatus* and *Verticillium* sp. TH28, and the ME extracts of *Clonostachys rosea* and *Corynespora cassiicola*, *Phialophora verrucosa*, and *Spadicoides* sp. inhibited the settling of *M. persicae* and *R. padi* (SI > 80%; Table 2). The extracts of *C. rosea* and *G. masseei* caused strong settling inhibition of both aphid species. The dose-response evaluations for these extracts are shown in Table 3. For *M. persicae*, the ME extract of *C. rosea* was the most active, with an EC₅₀ value of 8 µg/cm²; whereas for *R. padi*, the EA extract of *G. masseei* was the most active with an EC₅₀ value of 19 µg/cm².

Molecular identification by using ITS1 and ITS4 primers resulted in a DNA band of ~650 bp. After edition, 561 bp were used as query in BLAST, finding the highest homology with *G. masseei* (first hit accession AB540554.1). Query cover about 97% of known *G. masseei* ITS fragment, with an E value of 0.0 and 99% of identity.

The chemical composition of the EA extract of *G. masseei* was determined by a bioassay-guided fractionation. The hexane (ARGM-1A) and acetonitrile (ARGM-1B) fractions had strong effects on settling inhibition of *M. persicae* (>85%) and *R. padi* (SI > 70%) (Table 3). Analysis of fraction ARGM-1A and its methyl derivative (ARGM-4) by GC-MS and by comparison with commercial standards (free and methyl esters) revealed the presence of four fatty acids as major components. Fatty acids included hexadecanoic acid (16:0), octadecanoic acid (18:0), 9-octadecenoic acid (18:1) and 9,12-octadecadienoic acid (18:2) (Table 4). The most abundant components were hexadecanoic acid (74%) and 9,12-octadecadienoic acid (16.8%). In addition, ARGM-1A fraction was methylated and the product, ARGM-4, also had an effect on the settling of both aphid species.

Purification of the second fraction ARGM-1B by silica gel columns yielded a complex mixture of compounds which included ergosterol. This compound was isolated and identified by comparison with a commercial sample (E6510-10G, Sigma-Aldrich). The subfractions were not active on aphid settling.

The compounds isolated in this work and commercial standards were evaluated on both aphid species (Table 5). Methyl hexadecanoate had a low effect on the settling of *R. padi* and was not tested on *M. persicae*. Methyl 9-octadecenoate strongly inhibited the settling of both aphid species (SI > 70%); likewise, methyl 9,12-octadecadienoate

Table 2 *In vitro* evaluation of the organic extracts from microfungi on *Meloidogyne javanica* (100 µg/ml), *Myzus persicae*, *Rhopalosiphum padi*, and *Spodoptera littoralis* (100 µg/cm²)

Fungal strains	Type of extract	M. j. %IM ^a	S. l. %FR ^b	Myzus persicae			Rhopalosiphum padi		
				%C	%T	%SI ^c	%C	%T	%SI ^c
<i>Cladosporium cladosporioides</i>	EA	9.1±2.1	20.2±16.2	64.9±4.0	35.1±4.0	45.1±7.0	77.5±2.7	22.5±2.7	67.5±5.3
	ME	0.7±0.6	47.3±17.0	83.9±3.0	16.2±3.0	77.4±4.9	81.6±3.3	18.4±3.3	73.5±5.2
<i>Clonostachys rosea</i>	EA	9.8±2.7	5.8±3.7	83.5±2.5	16.5±2.5	78.2±3.7	69.4±3.4	30.6±3.4	51.4±6.9
	ME ^d	13.3±1.6	47.5±17.0	94.4±1.2	5.7±1.2	93.3±1.4	86.6±2.0	13.4±2.0	81.5±3.3
<i>Corynespora cassiicola</i>	EA	8.5±1.0	28.4±16.0	76.5±2.8	23.5±2.8	65.7±5.3	68.8±3.1	31.3±3.1	50.4±6.7
	ME	3.8±0.1	33.8±15.0	92.2±1.7	7.8±1.7	90.8±2.0	79.6±2.5	20.5±2.5	72.0±3.9
<i>Cylindrium elongatum</i>	EA	2.7±0.6	43.8±17.0	76.7±4.5	23.3±4.5	65.0±7.7	66.1±4.0	33.9±4.1	44.5±7.7
	ME	7.3±0.6	12.3±8.0	73.2±3.1	26.8±3.1	58.1±6.6	75.8±3.5	24.2±3.5	62.6±6.1
<i>Cylindrocarpon congoense</i>	EA	5.6±0.1	41.9±19.4	74.1±4.2	25.9±4.2	56.2±8.3	76.6±2.7	23.4±2.7	66.3±4.9
	ME	2.6±1.1	53.2±16.8	84.6±2.5	15.4±2.5	77.5±4.2	83.1±4.0	16.9±4.0	76.0±6.3
<i>Fusarium</i> sp.	EA	12.4±1.5	13.1±8.3	71.3±2.9	28.7±2.9	55.3±6.0	75.1±3.7	24.9±3.7	61.6±7.2
	ME	33.4±3.8	16.1±8.6	80.3±2.3	19.7±2.3	73.4±3.9	77.0±2.7	23.0±2.7	63.2±4.9
<i>Gliomastix masseei</i>	EA ^d	8.2±0.7	20.6±10.0	93.7±1.7	6.3±1.7	91.7±2.5	93.3±1.3	6.7±1.3	91.9±1.7
	ME	0.3±0.7	23.4±10.6	76.8±2.7	23.2±2.7	66.9±4.5	84.5±4.0	15.5±4.0	74.4±6.6
<i>Perelegamyces parviechinulatus</i>	EA	8.1±1.8	18.1±9.4	87.3±3.0	12.7±3.0	82.3±4.9	81.3±3.4	18.7±3.4	72.5±5.8
	ME	3.9±1.5	13.9±6.4	77.4±3.4	22.6±3.4	65.2±6.8	86.1±5.2	13.9±5.2	77.7±8.8
<i>Phialophora verrucosa</i>	EA	4.4±0.7	21±8.1	66.8±4.7	33.2±4.7	49.1±7.6	67.4±3.4	32.6±3.4	48.3±6.9
	ME	0.7±0.3	7.6±4.8	90.7±1.8	9.3±1.8	88.9±2.3	68.0±4.3	32.0±4.3	48.7±7.8
<i>Spadicoides</i> sp.	EA	20.0±1.6	19.2±8.8	77.3±4.9	22.7±4.9	64.7±8.6	71.6±5.8	21.1±2.4	64.1±6.7
	ME	3.2±0.3	0	90.1±2.7	10.0±2.7	87.5±3.9	82.4±4.5	17.6±4.5	72.5±8.1
<i>Verticillium</i> sp.	EA ^d	3.2±0.7	25.3±12.6	95.0±1.1	5.0±1.1	93.7±1.3	81.0±3.8	19.0±3.8	71.9±6.8
	ME	3.0±1.2	61.3±11.3	83.9±4.6	16.1±4.6	75.0±8.0	66.4±5.6	33.6±5.6	47.7±9.0
<i>Volutella</i> sp.	EA	nt	nt	nt	nt	nt	nt	nt	nt
	ME	14.4±2.1	19.2±13.0	83.6±3.9	16.4±3.9	74.4±7.6	68.5±4.4	31.5±4.4	49.3±7.8
Fermented rice (blank)	EA	15.0±1.2	27.1±9.7	63.6±5.14	36.4±5.1	43.0±8.2	68.8±3.2	31.2±3.2	48.4±7.3
	ME	-0.5±0.5	13.8±6.4	64.8±3.6	35.2±3.6	46.5±5.8	54.7±1.9	45.3±1.9	18.5±4.7

EA: ethyl acetate extract, ME: methanol extract, M. j.: *Meloidogyne javanica*, S. l.: *Spodoptera littoralis*, nt: not tested.^a %IM: percent inhibition of mobility; n = 100–150 for *Meloidogyne javanica*.^b Percentage of feeding reduction (FR) = [1 – (T/C)] × 100; where T and C are the consumption of treated and control leaf discs, respectively; n = 5.^c Settling inhibition in percentage (SI) = [1 – (%T/%C)] × 100; aphids settled on treated leaf disc (%T) and control leaf disc (%C) after 24 h. n = 20 replicates for aphids (d: n = 40 replicates).

Table 3 Dose-response evaluation of active fungal extracts from *Clonostachys rosea* and *Gliomastix masseei* on settling inhibition of *Myzus persicae* and *Rhopalosiphum padi*

Fungal strain/solvent	Concentration μg/cm ²	<i>Myzus persicae</i>				<i>Rhopalosiphum padi</i>			
		%C	%T	%SI ^a	EC ₅₀ (CL) μg/cm ²	%C	%T	%SI ^a	EC ₅₀ (CL) μg/cm ²
<i>Clonostachys rosea</i> <i>MeOH extract</i>	200	96.7 ± 2.1	3.3 ± 2.1	95.1 ± 3.4	8 (6, 10)	87.9 ± 2.1	12.1 ± 2.1	84.6 ± 2.9	38 (25, 48)
	100 ^b	94.4 ± 1.2	5.7 ± 1.2	93.3 ± 1.4		86.6 ± 2.0	13.4 ± 2.0	81.5 ± 3.3	
	50	88.0 ± 3.5	12.0 ± 3.5	83.8 ± 5.4		72.2 ± 4.0	27.8 ± 4.0	54.3 ± 7.9	
	20	79.5 ± 3.8	20.5 ± 3.8	70.0 ± 6.5					
	4	57.7 ± 4.6	42.3 ± 4.6	33.7 ± 7.2					
<i>Gliomastix masseei</i> <i>Ethyl acetate extract</i>	200	96.8 ± 1.6	3.2 ± 1.6	96.0 ± 2.0	15 (13, 18)	92.6 ± 2.0	7.4 ± 2.0	91.0 ± 2.5	19 (16, 23)
	100 ^b	93.7 ± 1.7	6.3 ± 1.7	91.7 ± 2.5		93.3 ± 1.3	6.7 ± 1.3	91.9 ± 1.7	
	50	82.0 ± 2.9	18.0 ± 2.9	74.6 ± 5.3		77.3 ± 3.7	22.7 ± 3.7	66.6 ± 6.5	
	20	68.3 ± 4.5	31.7 ± 4.5	50.9 ± 7.7		60.7 ± 2.7	39.3 ± 2.7	34.3 ± 5.6	
	4	49.2 ± 6.2	50.8 ± 6.2	22.6 ± 8.7		57.5 ± 2.2	42.5 ± 2.2	26.5 ± 5.2	
ARGM-1A	50	90.6 ± 2.5	9.4 ± 2.5	87.7 ± 3.8		78.9 ± 2.6	21.1 ± 2.6	70.8 ± 4.6	
ARGM-1B	50	93.9 ± 2.0	6.1 ± 2.0	92.5 ± 2.5		83.3 ± 3.1	16.7 ± 3.1	76.5 ± 5.3	

^a SI: settling inhibition in percentage ± standard error; SI = [1 - (%T/%C)] × 100; where %T and %C are the percentage of aphids settled on the treated and control leaf pieces, respectively, after 24 h; n = 20 replicates, 40 replicates.

EC₅₀ (CL): effective concentration (confidence limits) for settling inhibition. EC₅₀ values are considered significantly different if confidence limits do not overlap.

nt = not treated.

Fraction ARG-1A: hexane fraction after partition of EA extract.

Fraction ARG-1B: acetonitrile fraction after partition of EA extract.

Table 4 Analysis of the chemical constituents of *Gliomastix masseei* hexane fraction (ARGM-1A) and its methylated derivative (ARGM-4) by GC-MS techniques

Compound assignment	Retention time (min)	% Relative abundance
<i>Fraction ARGM-1A</i>		
Methyl hexadecanoate	9.88	15.0
Ethyl hexadecanoate	10.88	3.0
Methyl 9,12-octadecadienoate	11.82	17.8
Methyl 9-octadecenoate	11.91	9.3
Ethyl 9,12-octadecadienoate	12.46	5.4
<i>ARGM-4</i>		
Methyl hexadecanoate	9.58	75.0
Methyl 9,12-octadecadienoate	11.49	3.0
Methyl 9-octadecenoate	11.59	16.8
Methyl octadecanoate	11.86	4.0

only inhibited the settling of *M. persicae* ($SI > 70\%$). Ergosterol showed no activity. Dose-response evaluation of both active pure esters showed that methyl 9-octadecenoate was the most active compound, with an EC_{50} value of $16 \mu\text{g}/\text{cm}^2$ for *M. persicae* and $35 \mu\text{g}/\text{cm}^2$ for *R. padi* (Table 5).

Discussion

In the present work, the evaluation of organic extracts of microfungi showed that the ME and EA extracts had different effects on the plant pest targets. The extracts had very low or no effect on *S. littoralis* or *M. javanica*. However, various extracts displayed a strong effect on the settling of aphids *M. persicae* and *R. padi*. To our knowledge, the fungal extracts evaluated in the present work had not been previously assessed on aphid settling. The results showed that 30% of the evaluated fungi produced metabolites that deterred aphids. The most active were the EA extract of *G. masseei* (EC_{50} , $15\text{--}19 \mu\text{g}/\text{cm}^2$) and the ME extract of *C. rosea* (EC_{50} , $8\text{--}38 \mu\text{g}/\text{cm}^2$).

The molecular analysis identified fungus *G. masseei*. Minor discrepancies between the query and retrieved sequences were observed, however identity was supported by the congruency of molecular identification with traditional taxonomy performed according to Kiyuna et al.¹⁹. Thus, the ITS fragment was enough to identify the fungus and discrepancies can be attributed to usual polymorphism among strains. The fungus *G. masseei* is a cosmopolitan species that has antimicrobial properties²⁸; however, no studies on identification of secondary metabolites produced by this species have been reported. The primary fractionation (ARGM-1A and 1B) of the extract revealed that *G. masseei* produces metabolites that affect aphid settling. These metabolites were detected in the low and medium polarity fractions. In the low polarity fraction, fatty acids and their derivatives are the most abundant components. Although hexadecanoic acid (75%) was the major constituent, its effect on the settling of *M. persicae* was produced only by their derivatives methyl and ethyl esters ($5.5\text{--}8.7 \mu\text{g}/\text{cm}^2$), in agreement with previous reports³². Surprisingly, these compounds showed no effect on the

settling of *R. padi*. Other components of the low polarity fraction were the methyl esters of 9,12-octadecadienoic acid (17.8%) and 9-octadecenoic acid (9.3%). In this case, previous studies have shown that the acid forms of these compounds have anti-aphid properties⁸. In the present study, methyl 9-octadecenoate was the most active compound with an EC_{50} of 16 and $35 \mu\text{g}/\text{cm}^2$, for *M. persicae* and *R. padi*, respectively (Table 5). Methyl 9,12-octadecadienoate, in contrast, only inhibited the settling of *M. persicae* ($EC_{50} = 20 \mu\text{g}/\text{cm}^2$), but not of *R. padi*.

During the purification of the extract, we observed that the settling inhibition activity remained in low and medium polarity fractions for *M. persicae*, but decreased for *R. padi*, where SI values decreased from crude extract (91.9) to 70 and 76%, respectively. An opposite trend was observed for the natural methylated ARGM-4 fraction ($SI = 84\%$). We then verified the activity of the isolated methyl ester mixture and an artificial mixture of commercial standards of the major constituents present in the fraction ARGM-1A. The mixture was prepared with the methyl esters of the hexadecanoic, 9,12-octadecadienoic, and 9-octadecenoic acids (27:6:1). The methyl 9-octadecenoate and natural mixture of hexane fraction inhibited the settling of both aphids ($SI > 70\%$); however, 9,12-octadecadienoate and the artificial mixture only inhibited the settling of *M. persicae*, in the same manner as the saturated methyl hexadecanoate. These data showed that on esterified fatty acids, the pattern for anti-aphid activity is different in comparison with that of the free unsaturated fatty acids⁸. The difference between the effects on the settling of *R. padi* caused by the artificial and natural mixtures can be attributed to the presence of other minor unknown constituents in the latter.

The biological activity of fatty acids and their derivatives on pest insects have been previously reported²⁷. For example, 9,12-octadecadienoic acid has shown activity on *Aedes aegypti*, *Callosobruchus maculatus*, *Lymantria dispar*, *Liposcelis bostrychophila*, and *Malacosoma disstria*; 9-octadecenoic acid on *Anopheles aegypti*, *Anopheles stephensi*; 9,12-octadecadienoic, hexadecanoic and octadecanoic acids on *S. frugiperda*^{1,18,25}. Fatty acids can be extracted from plants and from fungi. In this sense, various studies have found that different species of fungi, such as *Aspergillus ustus*, *Cladosporium* sp., *Curvularia* sp., *Penicillium* sp., and *Rhizopus oryzae* produce this type of compounds^{23,24,31}.

The TLC and GC-MS analyses of acetonitrile fraction of *G. masseei* showed a complex mixture of compounds, where traces of ergosterol, a component of the fungal membrane, were observed. Ergosterol is commonly oxidized in the presence of light into ergosterol endoperoxide. There are various reports on the strong antimicrobial activity of ergosterol endoperoxide, but no data on its effect against insects has been published^{5,30}. Although ergosterol endoperoxide was not detected in the *G. masseei* extracts, its effect was evaluated on aphid settling, where this compound slightly inhibited the settling of *M. persicae* and *R. padi* (Table 5).

It is important to note that the activity of the full extract of *G. masseei*, compared to that of the fractions, subfractions and commercial standards, suggests that the anti-aphid effect was most likely due to the synergistic effect of the known and unknown minor components; however, further investigation is required in this regard.

Table 5 Effects of methylated fraction (ARGM-4) from *Gliomastix masseei*, and methyl esters from commercial fatty acids on settling of *Myzus persicae* and *Rhopalosiphum padi* and dose-response evaluation of the most active methyl esters

Sample	Concentration μg/cm ²	<i>Myzus persicae</i>				χ^2	<i>Rhopalosiphum padi</i>				χ^2
		%C	%T	%SI ^a	EC ₅₀ (CL) μg/cm ²		%C	%T	%SI ^a	EC ₅₀ (CL) μg/cm ²	
ARGM-4	50	82.8 ± 2.6	17.2 ± 2.6	76.7 ± 4.0			87.6 ± 2.3	12.4 ± 2.3	84.1 ± 3.3		
Methyl hexadecanoate	50	nt	nt	nt			73.8 ± 2.5	26.2 ± 2.5	61.8 ± 4.4		
Methyl 9-octadecenoate	50	87.3 ± 2.6	12.6 ± 2.6	83.5 ± 3.7	16 (15, 17)	2.0	79.4 ± 2.7	20.5 ± 2.7	71.4 ± 4.4	35 (29, 41)	2.13
	25	73.9 ± 7.0	26.0 ± 7.0	66.9 ± 8.6			59.0 ± 3.8	40.9 ± 3.8	34.0 ± 6.5		
	12.5	54.6 ± 7.3	45.3 ± 7.3	38.8 ± 9.0			53.5 ± 4.3	46.4 ± 4.3	23.4 ± 6.7		
Methyl 9,12- octadecadienoate	50	82.0 ± 2.3	18 ± 2.3	76.2 ± 3.4	20 (18, 22)	1.6	nt	nt	nt		
	25	68.7 ± 5.6	31.3 ± 5.6	52.7 ± 9.0							
	12.5	62.2 ± 6.3	37.7 ± 6.3	41.0 ± 9.0							
Artificial mixture*	50	77.5 ± 4.2	19.7 ± 3.3	71.2 ± 5.8			70.6 ± 2.8	29.4 ± 2.8	54.5 ± 5.5		
Ergosterol	50	68.6 ± 3.0	31.4 ± 3.0	51.8 ± 6.5			69.3 ± 3.2	30.7 ± 3.2	51.5 ± 6.9		
Ergosterol endoperoxide	50	74.6 ± 3.0	25.4 ± 3.0	64.4 ± 5.6			64.1 ± 3.6	35.9 ± 3.6	38.6 ± 8.2		

ARGM-4: methylated ARGM-1A fraction.

* Artificial mixture: methyl hexadecanoate, methyl 9-octadecenoate, and methyl 9,12-octadecadienoate (27:6:1).

^a Settling inhibition in percentage ± standard error; SI = [1 - (%T/%C)] × 100, where %T and %C are the percentage of aphids settled on the treated and control leaf pieces, respectively, after 24 h; n = 20 replicates.

EC₅₀ (CL): effective concentration (confidence limits) for settling inhibition. EC₅₀ values are considered significantly different if confidence limits do not overlap.

nt = not tested.

The fungus *C. rosea* is a cosmopolitan species which has shown a wide range of biological activities, in particular as a biocontrol agent against phytopathogenic fungi and nematodes^{21,34}. The activity in this case has been attributed to secondary metabolites, such as glioclidine A-E, verticillin A and 11'-deoxy-verticillin A, Sch52900, and Sch52901¹¹. In previous works, we have found that the EA extract of *C. rosea* is active against human pathogenic bacteria, plant pathogenic fungi, and plant parasitic nematodes^{13,14}. Surprisingly, in the present work, the EA extract of *C. rosea* showed no activity on *M. javanica*.

It is also pertinent to mention that the effect of the most active fungal extracts evaluated in the present work was similar to that of other natural products isolated from microorganisms or plants³.

In summary, the settling inhibition capacity of the fungal extracts screened in this work highlights the potential of these compounds for biorational plant protection. ME extracts of *C. rosea* and EA extracts of *G. masseei* showed the highest anti-settling activity on *M. persicae* and *R. padi*. Furthermore, we observed that ester derivatives of hexadecanoic, 9-octadecenoic and 9,12-octadecadienoic acids were some of the active principles and the most abundant components in the EA extract of *G. masseei*, which may have a synergistic effect with other unknown minor constituents. This work also contributes with insights into the biological activity of natural products derived from micromycetes isolated from the tropical regions of Mexico.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that no patient data are included in this article.

Right to privacy and informed consent. The authors declare that no patient data are included in this article.

Conflict of interest

The authors declare that there is no conflict of interests regarding the publication of this article.

Acknowledgments

This research was supported by Conacyt (Grants 47549 and 2009/CB131256, respectively); our thanks also go to Conacyt for the postgraduate scholarship to A.L.R.J. (Conacyt No. 228272). We thank Daniela Hurtado-Cantillo, Miguel Tzec-Sima and Felicia Moo-Koh their assistance in the laboratory.

References

1. Adebowale KO, Adedire CO. Chemical composition and insecticidal properties of the underutilized *Jatropha curcas* seed oil. Afr J Biotechnol. 2006;5:901–6.
2. Ahmad F, Rather MA, Siddiqui MA. Nematicidal activity of leaf extracts from *Lantana camara* L. against *Meloidogyne incognita* (Kofoid and White) Chitwood and its use to manage roots infection of *Solanum melongena* L. Braz Arch Biol Technol. 2010;53:543–8.
3. Argandoña VH, Rovirosa J, San-Martín A, Riquelme A, Díaz-Marrero AR, Cueto M, Darias J, Santana O, Guadaño A, González-Coloma A. Antifeedant effects of marine halogenated monoterpenes. J Agric Food Chem. 2002;50:7029–33.
4. Asaff A, Cerdá-García-Rojas C, de la Torre M. Isolation of dipicolinic acid as an insecticidal toxin from *Paecilomyces fumosoroseus*. Appl Microbiol Biotechnol. 2005;68:542–7.
5. Budhiraja A, Nepali K, Sapra S, Gupta S, Kumar S, Dhar KL. Bioactive metabolites from an endophytic fungus of *Aspergillus* species isolated from seeds of *Gloriosa superba* Linn. Med Chem Res. 2013;22:323–9.
6. Burgueño-Tapia E, Castillo L, González-Coloma A, Joseph-Nathan P. Antifeedant and phytotoxic activity of the sesquiterpene p-benzoquinone and some of its derivatives. J Chem Ecol. 2008;34:766–71.
7. Cantrell CL, Dayan FE, Duke SO. Natural products as sources for new pesticides. J Nat Prod. 2012;75:1231–42.
8. Castillo L, Díaz M, González-Coloma A, González A, Alonso-Paz E, Bassagoda MJB, Rossini C. *Clytostoma callistegioides* (Bignoniacae) wax extract with activity on aphid settling. Phytochemistry. 2010;71:2052–7.
9. Cole M, Rolinson GN. Microbial metabolites with insecticidal properties. Appl Microbiol. 1972;24:660–2.
10. Di Toto Blessing L, Álvarez Colom O, Popich S, Neske A, Bardón A. Antifeedant and toxic effects of acetogenins from *Annona montana* on *Spodoptera frugiperda*. J Pest Sci. 2010;83:307–10.
11. Dong JY, He HP, Shen YM, Zhang KQ. Nematicidal epipolysulfanyl-dioxopiperazines from *Gliocladium roseum*. J Nat Prod. 2005;68:1510–3.
12. Gaich T, Mulzer J. Total synthesis of (–)-penifulvin A, an insecticide with a dioxafenestrane skeleton. J Am Chem Soc. 2009;131:452–3.
13. Gamboa-Angulo M, De la Rosa-García SC, Heredia-Abarca G, Medina-Baizabal IL. Antimicrobial screening of tropical microfungi isolated from sinkholes located in the Yucatan peninsula, Mexico. Afr J Microbiol Res. 2012;6:2305–12.
14. Gamboa-Angulo M, Molina-Salinas GM, Chan-Bacab M, Peraza-Sánchez SR, Heredia G, De la Rosa-Garcia SC, Reyes-Estebanez M. Antimycobacterial and antileishmanial effects of microfungi isolated from tropical regions in Mexico. Parasitol Res. 2013;112:559–66.
15. García M, González-Coloma A, Donadel OJ, Ardanaz CE, Tonn CE, Sosa ME. Insecticidal effects of *Flourensia oolepis* Blake (Asteraceae) essential oil. Biochem Syst Ecol. 2007;35:181–7.
16. Hernández-Carlos B, González-Coloma A, Orozco-Valencia AU, Ramírez-Mares MV, Andrés-Yeves MF, Joseph-Natan P. Bioactive saponins from *Microsechium helleri* and *Sicyos bulbosus*. Phytochemistry. 2011;72:743–51.
17. Juvalle K, Stefan K, Wiese M. Synthesis and biological evaluation of flavones and benzoflavones as inhibitors of BCRP/ABCG2. Eur J Med Chem. 2013;67:115–26.
18. Kannanathan K, Senthilkumar A, Venkatesulu V, Chandrasekaran M. Larvicidal activity of fatty acid methyl esters of *Vitex* species against *Culex quinquefasciatus*. Parasitol Res. 2008;103:999–1001.
19. Kiyuna T, An KD, Kigawa R, Sano C, Miura S, Sugiyama J. Molecular assessment of fungi in “black spots” that deface murals in the Takamatsuzuka and Kitara Tumuli in Japan: *Acremonium* sect. *Gliomastix* including *Acremonium tumulicola* sp.nov. and *Acremonium felinum* comb. nov. Mycoscience. 2011;52:1–17.
20. Li GH, Zhang KQ, Xu JP, Dong JY, Liu YJ. Nematicidal substances from fungi. Recent Pat Biotechnol. 2007;1:212–33.

21. Li J, Yang J, Huang X, Zhang KQ. Purification and characterization of an extracellular serine protease from *Clonostachys rosea* and its potential as a pathogenic factor. *Process Biochem.* 2006;41:925–9.
22. Moo-Koh FA, Alejo JC, Reyes-Ramírez A, Tun-Suárez JM, Sandoval-Luna R, Ramírez-Pool JA. *In vitro* activity of an aqueous extract of *Bonellia flammnea* against phytopathogenic fungi. *Agrociencia.* 2014;48:833–45.
23. Oda Y, Yajima Y, Kinoshita M, Ohnishi M. Differences of *Rhizopus oryzae* strains in organic acid synthesis and fatty acid composition. *Food Microbiol.* 2003;20:371–5.
24. Oleinikova GK, Slinkina NN, Afifyatullov Sh. Nonpolar compounds and free fatty acids from marine fungi *Aspergillus ustus*. *Chem Nat Compd.* 2011;47:806–7.
25. Pérez-Gutiérrez S, Zavala-Sánchez MA, González-Chávez MM, Cárdenas-Ortega NC, Ramos-Lopez MA. Bioactivity of *Carica papaya* (Caricaceae) against *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Molecules.* 2011;16:7502–9.
26. Poitout S, Bues R. Elevage de plusieurs espèces de Lépidoptères Noctuidae sur milieu artificiel riche et sur milieu artificiel simplifié. *Ann Zool Ecol Anim.* 1970;2:79–91.
27. Ramos-Lopez MA, González-Chávez MM, Cárdenas-Ortega NC, Zavala-Sánchez MA, Pérez S. Activity of the main fatty acids components of the hexane leaf extract of *Ricinus communis* against *Spodoptera frugiperda*. *Afr J Biotechnol.* 2012;11:4274–8.
28. Reyes Estebanez M, Heredia Abarca G, Gamboa Angulo MM. Perfil biológico de hongos anamórficos del sureste de México. *Rev Mex Micol.* 2008;28:49–56.
29. Reyes-Estebanez M, Herrera-Parra E, Cristóbal-Alejo J, Heredia-Abarca G, Canto-Canché B, Medina-Baizabal I, Gamboa-Angulo M. Antimicrobial and nematicidal screening of anamorphic fungi isolated from plant debris of tropical areas in Mexico. *Afr J Microbiol Res.* 2011;5:1083–9.
30. Rhee YH, Jeong SJ, Lee HJ, Lee HJ, Koh W, Jung JH, Kim SH, Kim SH. Inhibition of STAT3 signaling and induction of SHP1 mediate antiangiogenic and antitumor activities of ergosterol peroxide in U266 multiple myeloma cells. *BMC Cancer.* 2012;12: 28–38.
31. Rues L, Häggblom MM, García Zapata EJ, Dighton J. Fatty acids of fungi and nematodes—possible biomarkers in the soil food chain? *Soil Biol Biochem.* 2002;34:745–56.
32. Santana O, Reina M, Fraga BM, Sanz J, González-Coloma A. Antifeedant activity of fatty acid esters and phytosterols from *Echium wildpretii*. *Chem Biodivers.* 2012;9:567–76.
33. Sharma A, Sharma S, Dalela M. Nematicidal activity of *Paecilomyces lilacinus* 6029 cultured on Karanja cake medium. *Microb Pathog.* 2014;75:16–20.
34. Viccini G, Mannich M, Capalbo DMF, Valdebenito-Sanhueza R, Mitchell DA. Spore production in solid-state fermentation of rice by *Clonostachys rosea*, a biopesticide for gray mold of strawberries. *Process Biochem.* 2007;42:275–8.
35. White TJ, Bruns T, Lee S, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, editors. *PCR protocols: a guide to methods and applications.* San Diego, CA: Academic Press; 1990. p. 315–22.
36. Yang ZS, Li GH, Zhao PJ, Zheng X, Luo SL, Li L, Niu XM, Zhang KQ. Nematicidal activity of *Trichoderma* spp. and isolation of an active compound. *World J Microbiol Biotechnol.* 2010;26:2297–302.
37. Yang ZS, Yu Z, Lei L, Xia Z, Shao L, Zhang KQ, Li GH. Nematicidal effect of volatiles produced by *Trichoderma* sp. *J Asia-Pac Entomol.* 2012;15:647–50.