

Original article

Plant extracts to control *Alternaria alternata* in Murcott tanger fruits

Daniel Diego Costa Carvalho^a, Eduardo Alves^{a,*}, Renato Barbosa Camargos^a,
Denilson Ferreira Oliveira^b, José Roberto Soares Scolforo^c, Douglas Antônio de Carvalho^d,
Tereza Raquel Sâmia Batista^e

^a Departamento de Fitopatologia, Universidade Federal de Lavras, Lavras, Minas Gerais, Brazil

^b Departamento de Química, Universidade Federal de Lavras, Lavras, Minas Gerais, Brazil

^c Departamento de Ciências Florestais, Universidade Federal de Lavras, Lavras, Minas Gerais, Brazil

^d Departamento de Biologia, Universidade Federal de Lavras, Lavras, Minas Gerais, Brazil

^e Centro Universitário de Lavras, Lavras, Minas Gerais, Brazil

ARTICLE INFO

Article history:

Received 22 February 2011

Accepted 3 May 2011

Available online 13 May 2011

Keywords:

Alternaria brown spot

Citrus

Postharvest

Plant disease control

Natural products

Scanning electron microscopy

ABSTRACT

Background: *Alternaria alternata* causes the *Alternaria* brown spot disease (ABS) in many tangerines and their hybrids worldwide. Plant extracts offer an alternative method for controlling this disease, which control is based on chemical fungicides.

Aims: To identify plant species with antifungal properties against *A. alternata*, the causal agent of the ABS.

Methods: Plant extracts prepared from leaves, barks, flowers, and stalks collected from 105 plant species in the State of Minas Gerais, Brazil, were tested for activity against the fungus *A. alternata* in vitro and in vivo.

Results: The most promising extract was obtained from *Anadenanthera colubrina*, which reduced the disease on Murcott tanger fruits to levels obtained with commercial fungicides. *Artemisia annua*, *Cariniana estrellensis*, *Ficus carica*, and *Ruta graveolens* presented moderate in vitro antifungal activity, but no effects were observed on the disease when the extracts were applied to fruits inoculated with the fungus. Besides, *A. colubrina* was the most active extract against *A. alternata* in the in vitro assay.

Conclusions: The results obtained in the in vitro and in vivo assays suggested that the fungal growth test, which uses 96-well polypropylene plates, seems to be appropriate for selecting potential plant species for testing new methods to control ABS.

© 2011 Revista Iberoamericana de Micología. Published by Elsevier España, S.L. All rights reserved.

Extractos vegetales para el control de *Alternaria alternata* en frutos de tanger Murcott

RESUMEN

Palabras clave:

Mancha marrón de *Alternaria*

Cítricos

Postcosecha

Control de enfermedades vegetales

Productos naturales

Microscopia electrónica de barrido

Antecedentes: *Alternaria alternata* causa la mancha marrón en muchas mandarinas y en sus híbridos en todo el mundo. Extractos de plantas proporcionan un método alternativo para controlar esta enfermedad cuyo control se basa en fungicidas químicos.

Objetivos: Identificar las especies de plantas con propiedades antifúngicas contra *A. alternata*, el agente causal de la mancha marrón.

Métodos: Extractos de plantas preparados a partir de hojas, corteza, flores y tallos recogidos de 105 especies de plantas en el Estado de Minas Gerais, Brasil, fueron utilizados para estudiar su actividad contra el hongo *A. alternata* in vitro e in vivo.

Resultados: El extracto más prometedor se obtuvo de *Anadenanthera colubrina*, que redujo la enfermedad en las frutas de tanger Murcott a los niveles obtenidos con fungicidas comerciales. *Artemisia annua*, *Cariniana estrellensis*, *Ficus carica* y *Ruta graveolens* presentaron moderada actividad antifúngica in vitro, pero no se observaron efectos sobre la enfermedad cuando los extractos fueron aplicados a los frutos inoculados con el hongo. Además, *A. colubrina* fue el más activo contra *A. alternata* en el ensayo in vitro.

* Corresponding author.

E-mail address: ealves@dfp.ufla.br (E. Alves).

Conclusiones: Los resultados obtenidos en los ensayos *in vitro* e *in vivo* sugieren que el método de crecimiento de hongos, que utiliza placas de 96 pozos de polipropileno, parece apropiado para la selección de especies potenciales para testar nuevos métodos de control de la mancha marrón.

© 2011 Revista Iberoamericana de Micología. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Alternaria brown spot (ABS) is an important disease of many tangerines and their hybrids in humid and semiarid areas worldwide.²³ In Brazil, ABS was first found in the State of Rio de Janeiro and then became widespread in the States of São Paulo and Minas Gerais, the main citrus growing areas in Brazil.^{16,17} The disease has caused economical losses to Murcott tangor (probable *Citrus sinensis* [L.] Osb. x *C. reticulata* Blanco hybrid) producers.¹⁵ The causal agent of this disease is the fungus *Alternaria alternata* (Fr:Fr) Keissl f. sp. *citri*, which can attack both leaves and fruits, reducing the yield. Specifically on fruits, which are very susceptible to infections according to Vicent et al.,²⁵ lesions varying from small dark necrotic spots to large sunken pockmarks turn them unmarketable, further increasing the producers losses.²⁴

In order to produce fruits with good external quality in areas affected by ABS, the application of foliar fungicides is usually necessary to reduce the losses caused by this disease. When the environmental conditions are adequate to the development of the fungus, up to 15 fungicide applications might be necessary to control ABS.²³ Thus, the production costs are increased and contamination of foods and environment with toxic substances also occur.¹¹ Consequently, the development of more efficient, cheaper, and less toxic products active against *A. alternata* is necessary.^{3,10}

One of the potentially useful alternatives to expensive and possibly toxic fungicides could be the use of plant extracts. They have presented very promising results on the control of plant pathogens.²² For example, Zimmu (*Allium cepa* L. x *Allium sativum* L.) extracts efficiently reduced the mycelial growth of *Alternaria solani* Sorauer.⁹ Tegengne et al.²¹ conducted a screening program from which they selected plants with antifungal properties and the extract of *Agapanthus africanus* (L.) Hoffm was shown to be the most potent against *Mycosphaerella pinodes*, the causal agent of black spot in pea.

In the present study our goal was to identify plant species with antifungal properties against *Alternaria alternata*, the causal agent of the ABS disease in many tangerines and their hybrids worldwide. Therefore, we tested 105 plant species and 18 were selected for additional assays.

Materials and methods

Plant extracts

The 105 plant species were collected in Minas Gerais State, Brazil. They were identified by comparison with specimens available in the Herbarium ESAL-Universidade Federal de Lavras, during the year of 2005 (Table 1). Leaves, barks, flowers, and stalks were oven-dried for 48 h at 40 °C. After grounding plant materials to pieces smaller than 2 mm, part (20 g) of each sample was immersed into methanol (50 ml) for 48 h at room temperature (15 to 30 °C). The resulting mixtures were filtrated through cotton wool plugs and the residues were extracted twice with more methanol. The three liquid phases obtained from each plant material were combined and concentrated to dryness in a rotary evaporator. The resulting 126 dry extracts were freeze-dried and stored at –10 °C for future use in the experiments.

Fungus isolation and growth conditions

Fruits exhibiting symptoms of ABS were submitted to the pathogen isolation protocol.⁵ Thus, pieces (10–25 mm²) of washed fruit peels from Murcott tangor were subsequently immersed into 70% ethanol (30–60 s), 2% sodium hypochlorite (30–60 s) and distilled water (2 x 30 s). Four fragments were placed in a Petri dish containing potato-dextrose-agar (PDA). After seven days at 25 °C, under a 12 h photoperiod, 9 mm agar plugs of the medium containing the fungus mycelium were transferred to new PDA plates.¹⁸ Then, *A. alternata* (Fr:Fr) Keissl f. sp. *citri* was isolated and identified as described by Carvalho et al.⁵

Production of conidia

Fungal plugs (9 mm diameter) from the culture medium were transferred to PDA plates, which were maintained for seven days at 25 °C under constant illumination provided by Philips daylight fluorescent lamps (20W, TLT, 75RS). The conidia were removed from the PDA plates by adding 10 ml of 1% (v/v) Tween 80. The resulting mixture was filtered using sterilized cheesecloth.

Fungal growth assay

Dried plant extracts (2 mg) were subsequently dissolved in 500 µl of 1% (v/v) Tween 80 and mixed with 100 µl of an *A. alternata* conidial suspension at 2.6–3.0 x 10⁵ conidia/ml. Twenty microliters of the resulting conidia suspension were poured into each 300 µl well of a 96-well polypropylene plate containing 130 µl of PDA with terramycin-oxitetraclacin chlorhydrate 500 mg (Pfizer, 0.55 mg/ml PDA). After three days at 25 °C, under a 12 h photoperiod, plant extracts that prevented fungal growth were considered active. This experiment was carried out with four replicates, using 1% (v/v) Tween 80 as negative control. Aqueous 3.5 mg/ml Dacobre WP (chlorotalonil 250 g/kg and copper oxychloride 504 g/kg, Iharabras S.A. Chemicals Industries) solution in 1% (v/v) Tween 80, and aqueous 0.16 mg/ml Amistar 500 WG (Azoxistrobin 500 g/kg, Syngenta Crop Protection) solution in 1% (v/v) Tween 80, were used as positive controls.

Scanning electron microscopy

Fungal disks treated with the most active extracts (*A. colubrina*, *R. graveolens*, and *A. annua*), and the positive (Dacobre WP, Amistar 500 WG) and the negative controls (1% Tween 80) during the fungal growth assay, were removed from the polypropylene plates and submitted to the procedure described by Bozzola and Russel⁴ with few adaptations. Each disk was fixed in a modified Karnovsky solution (2.5% glutaraldehyde, 2% paraformaldehyde in a 0.05 M sodium cacodylate buffer at pH 7.2 containing 0.001 M CaCl₂) for 48 h. The disks were washed three times with the same buffer for 30 min, post-fixed for 2 h in a 1% osmium tetroxide solution in 0.05 M sodium cacodylate buffer at pH 7.2, and washed three times with distilled water. They were then dehydrated in a gradient series of acetone solutions (25, 50, 75, 90 and 100%) and dried with carbon dioxide in a critical point dryer (Bal-tec CPD 030). Finally, the disks were mounted on aluminum stubs with double-sided tape

Table 1

Plant species collected in Minas Gerais State (Brazil) and used in the present study.

<i>Achillea millefolium</i> L. ^{a,d}	<i>Dendropanax cuneatus</i> (DC.) Decne & Planchon ^b	<i>Origanum vulgare</i> L. ^a
<i>Ageratum conyzoides</i> L. ^a	<i>Digitalis lanata</i> Ehrh. ^a	<i>Petiveria alliacea</i> L. ^a
<i>Albizia polycephala</i> (Benth.) Killip ^a	<i>Eclipta alba</i> (L.) Hassk ^a	<i>Piper tuberculatum</i> Jacq. ^a
<i>Allophylus edulis</i> (A. St.-Hil.) Radlk. ^{a,b}	<i>Equisetum arvense</i> L. ^d	<i>Plantago lanceolata</i> L. ^a
<i>Amaioua guianensis</i> Aublet ^b	<i>Eugenia florida</i> DC. ^a	<i>Plantago major</i> L. ^a
<i>Anadenanthera colubrina</i> (Vell.) Brenan ^b	<i>Euphorbia tirucalli</i> L. ^d	<i>Porophyllum ruderale</i> (Jack.) Cass. ^a
<i>Annona cacans</i> Warm. ^b	<i>Ficus carica</i> L. ^a	<i>Protium heptaphyllum</i> (Aublet) Marchand ^a
<i>Annona squamosa</i> L. ^a	<i>Ficus trigona</i> L. ^{f,b}	<i>Psidium guajava</i> L. ^a
<i>Artemisia absinthium</i> L. ^a	<i>Foeniculum vulgare</i> Miller ^{a,d}	<i>Pteridium aquilinum</i> L. ^a
<i>Artemisia annua</i> L. ^a	<i>Ginkgo biloba</i> L. ^a	<i>Punica granatum</i> L. ^a
<i>Artemisia vulgaris</i> L. ^a	<i>Glechoma hederacea</i> L. ^a	<i>Rhamnidium elaeocarpum</i> Reissek ^b
<i>Baccharis trimera</i> L. ^a	<i>Guazuma ulmifolia</i> Lam. ^b	<i>Ricinus communis</i> L. ^a
<i>Bathysa meridionalis</i> Smith & Downs ^b	<i>Hedera helix</i> L. ^a	<i>Rosamarinus officinalis</i> L. ^a
<i>Brugmansia suaveolens</i> (Willd.) Bercht. & Presl. ^b	<i>Hypericum perforatum</i> L. ^a	<i>Ruta graveolens</i> L. ^{a,c}
<i>Cabralea canjerana</i> (Vell.) Mart. ^{a,b}	<i>Ixora warmingii</i> Müll. Arg. ^{a,b}	<i>Salvia officinalis</i> L. ^a
<i>Calendula officinalis</i> L. ^{a,c}	<i>Jatropha curcas</i> L. ^{a,c}	<i>Sambucus nigra</i> L. ^{a,c}
<i>Callisthene major</i> Mart ^b	<i>Justicia pectoralis</i> Vault. ^a	<i>Schinus terebinthifolius</i> Raddi ^a
<i>Calyptanthus clusiifolia</i> (Miq.) O. Berg ^{a,b}	<i>Laurus nobilis</i> L. ^a	<i>Solanum argenteum</i> Dunal ^{a,b}
<i>Cariniana estrellensis</i> (Raddi) Kuntze ^{a,b}	<i>Lavandula officinalis</i> Chaich ^a	<i>Sonchus oleraceus</i> L. ^a
<i>Cariniana legalis</i> (Mart.) Kuntze ^a	<i>Leonurus sibiricus</i> L. ^a	<i>Styrax pohlilii</i> A.DC. ^a
<i>Celtis iguanaea</i> (Jacquin) Sargent ^{a,b}	<i>Malva sylvestris</i> L. ^a	<i>Symphytum officinale</i> L. ^a
<i>Centella asiatica</i> (L.) Urban ^a	<i>Mangifera indica</i> L. ^a	<i>Tagetes</i> spp. L. ^{a,c}
<i>Chenopodium ambrosioides</i> L. ^a	<i>Melissa officinalis</i> L. ^a	<i>Taraxacum officinale</i> Cass. ^a
<i>Citrus aurantium</i> L. ^a	<i>Mentha arvensis</i> L. ^a	<i>Terminalia brasiliensis</i> Camb. ^{a,b}
<i>Coffea arabica</i> L. ^a	<i>Mentha longifolia</i> (L.) Hudson ^a	<i>Tetradenia riparia</i> (Hoechst) NE. Br ^a
<i>Coix lacryma-jobi</i> L. ^a	<i>Mentha piperita</i> L. ^a	<i>Thymus vulgaris</i> L. ^a
<i>Cordia ecalyculata</i> Vell. ^b	<i>Mentha pulegium</i> L. ^a	<i>Tilia cordata</i> Mill ^a
<i>Croton floribundus</i> Sprengel ^b	<i>Mentha spicata</i> L. ^a	<i>Tithonia diversifolia</i> (Hemsl.) Gray ^a
<i>Croton urucurana</i> Baillon ^{a,b}	<i>Mimosa pudica</i> L. ^{a,c}	<i>Trichilia clausenii</i> C.DC. ^b
<i>Cryptocarya aschersoniana</i> Mez ^{a,b}	<i>Momordica charantia</i> L. ^a	<i>Trichilia hirta</i> L. ^b
<i>Cupania vernalis</i> Cambess ^b	<i>Musa sapientum</i> L. ^a	<i>Tropaeolum majus</i> L. ^{a,c}
<i>Curcuma longa</i> L. ^a	<i>Nepeta catarica</i> (Catnip.) ^a	<i>Urtiga dioica</i> L. ^a
<i>Cynara scolymus</i> L. ^a	<i>Nicotiana tabacum</i> L. ^a	<i>Vochysia tucanorum</i> Mart. ^b
<i>Daphnopsis fasciculata</i> (Meisner) Nevling ^b	<i>Ocimum basiculum</i> L. ^a	<i>Zanthoxylum pohlianum</i> Engl. ^b
<i>Datura metel</i> L. ^a	<i>Ocimum gratissimum</i> L. ^a	<i>Zingiber officinale</i> Rosc. ^a

Part of plant: ^aleaves; ^bbarks; ^cflowers; ^dstalks.

and coated with a 20 nm gold layer by vacuum evaporation (Bal-tec SCD 050). All samples were observed in an Evo40 Leo scanning electron microscope.

Conidial germination assay

Based on the fungal growth assay results, a total of 20 plant extracts were selected for the conidial germination assay. Four milligrams of each plant extract were dissolved in 1 ml of 1% (v/v) Tween 80 and mixed with 100 µl of an *A. alternata* conidia suspension at $2.6\text{--}3.0 \times 10^5$ conidia/ml. An aliquot of 520 µl of each final suspension was added to 4.0 ml of solidified water-agar (WA; 80 g of agar and 555 mg of tetracycline) medium contained in a 6.0 cm Petri dish. After 12 h at 25 °C, under illumination, conidia were counted and those with the germinative tube length larger or equal to the smaller conidia diameter were considered germinated. This experiment was carried out with four replications (50 conidia each), using an aqueous 1% (v/v) Tween 80 solution as negative control and 3.5 mg/ml Dacobre WP and 0.16 mg/mL Amistar 500 WG solutions in 1% (v/v) Tween 80 as positive controls.

Mycelial growth assay

Plant extracts (20 samples) dissolved (7 mg/ml) in 1% (v/v) Tween 80 were poured into 9 cm Petri dishes (0.5 ml/dish) containing PDA (8 ml/dish) with tetracycline (Bunker, 0.55 mg/ml PDA). Subsequently, a 9 mm PDA disk with a seven-day old *A. alternata* colony (1.8 cm from the colony center) was placed upside down in the center of each Petri dish containing PDA impregnated with plant extract. After seven days at 25 °C, under a 12 h photoperiod, fungal colony diameters were measured and data were converted into percentage. This experiment was done with three replications,

employing 1% (v/v) Tween 80 as negative control and 3.5 mg/ml Dacobre WP and 0.16 mg/ml Amistar 500 WG solutions in 1% (v/v) Tween 80 as positive controls.

pH measurements

The pH of 20 plant extracts (2 mg) dissolved in 600 µl of 1% (v/v) Tween 80 1% were measured by dipping pH-indicator strips (Acilit® pH 0–6 and Neutralit® pH 5–10, Merck) into the solutions and comparing the resulting colors to the standard provided by the manufacturer.

Assay with Murcott tangor fruits

Ripe and healthy Murcott tangor fruits were washed with sterilized water and then with 70% ethanol solution. They were let to dry for 60 min in a biosafety hood. For injuring the fruits, four points were selected around the point of fruits insertion and, just after, four perforations (3 mm deep) were made with a needle in each selected point.⁵ Each plant extract (2.0 mg) was dissolved in 600 µl of a 10^6 conidia/ml in 1% (v/v) Tween 80 and 20 µl aliquots were deposited on each selected point. This experiment was carried out with four fruits per treatment. The conidia suspension in 1% (v/v) Tween 80 was used as negative control and mixtures of conidia suspension and the fungicides Dacobre WP (3.5 mg/ml) and Amistar 500 WG (0.16 mg/ml) as positive controls. Fruits with no conidia, but treated with 20 µl of 1% (v/v) Tween 80, plant extract, or fungicide, were also used in this experiment. All fruits were kept in a growth chamber at 25 °C under a 12/12 photoperiod. After 8 and 12 days, the lesion diameter around each selected location on fruits was measured with a ruler.

Prior to the statistical analysis, data were converted into spots development rate (ABS dr) by dividing the mean value for each fruit by the mean value of all fruits treated only with conidia in 1% (v/v) Tween 80 (negative control). All experiments were repeated twice.

Statistical analysis

Converted data (ABS dr) and data from the conidial germination and mycelial growth assays were submitted to analysis of variance, and average values were compared by Scott-Knott calculations ($P \leq 0.05$). Statistical analyses were done using SISVAR software.⁷

Results

Fungal growth, conidial germination and mycelial growth assays

Among the 126 plant extracts tested in the fungal growth assay, only those from leaves of *Artemisia annua*, barks of *Anadenanthera colubrina* and a mixture of flowers and leaves of *Ruta graveolens* were active against *A. alternata* (Table 2). Consequently, such extracts, as well as 17 plant extracts that presented no activity in the fungus growth assay, were submitted to the conidial germination and mycelial growth assays.

The results from the conidial germination assay (Table 2) showed that *A. colubrina* extract (3.0 germinated conidia) was not

statistically different from the commercial fungicides Dacobre WP and Amistar 500 WG, which afforded values of 3.5 and 7.2 germinated conidia, respectively. Other extracts such as those from *A. annua* and *F. carica* also inhibited *A. alternata* conidia germination, but in a lesser extent (20.7 and 23.2 germinated conidia, respectively).

The mycelial growth assay (Table 2) also showed that *A. colubrina* extract was not statistically different from the commercial fungicides. Similarly to the conidial germination assay, other plant extracts, such as *C. estrellensis* (barks), also presented some antifungal activity in the mycelial growth (34% of inhibition). The best results were obtained with *A. annua* and *A. colubrina* (43 and 46% of inhibition, respectively), that were as efficient as the commercial fungicides (45–48% of inhibition).

Scanning electron microscopy

Scanning electron microscopy (SEM) was used to further investigate the effects of extracts from *A. colubrina*, *A. annua* and *R. graveolens* on *A. alternata*. The samples exposed to the *A. colubrina* extract revealed no conidia or mycelia. Regarding conidia exposed to *A. annua* extracts, shorter germ tubes than those exposed to Dacobre WP (chlorotalonil and copper oxychloride) were observed, while *R. graveolens* extracts and Amistar 500 WG caused shrinking in conidia (Fig. 1). The negative control (1% Tween 80) did not cause any spore shrinking.

Assay with Murcott tangor fruits

Based on the in vitro assays (fungal growth, conidial germination and mycelial growth assays), five plant extracts (*A. colubrina*, *A. annua*, *C. estrellensis*, *F. carica* and *R. graveolens*) were selected for assays with Murcott tangor fruits. Only that from *A. colubrina* showed suppression of lesions caused by *A. alternata* (Table 3; Fig. 2). Although the efficiency of this extract reduced over time (ABS dr was 0.38A at 8 days and 0.49B at 12 days), the results obtained with such plant extract were statistically similar to those of the commercial fungicides (ABS dr were 0.63 and 0.61 at 8 days and 0.57 and 0.70 at 12 days for Dacobre WP and Amistar 500 WG, respectively). A 16-day evaluation was also tried in this experiment, but except for the fungicides and *A. colubrina* treatments, all inoculated surfaces on fruits were completely deteriorated by fusion of all lesions. No disease was

Table 2

Effect of plant extracts on *Alternaria alternata* in the fungal growth, conidia germination and mycelial growth assays.

Treatments ^a	Fungal growth assay ^b	Germinated conidia ^c	Mycelial growth (%) ^c
<i>Achillea millefolium</i> (leaves)	-	31.5 c	88 c
<i>Achillea millefolium</i> (flowers)	-	36.5 c	95 c
<i>Anadenanthera colubrina</i> (barks)	+	3.0 a	54 a
<i>Artemisia annua</i> (leaves)	+	20.7 b	57 a
<i>Cariniana estrellensis</i> (leaves)	-	45.0 e	91 c
<i>Cariniana estrellensis</i> (barks)	-	33.5 c	66 b
<i>Citrus aurantium</i> (leaves)	-	35.5 c	93 c
<i>Croton urucurana</i> (leaves)	-	34.2 c	74 b
<i>Datura metel</i> (leaves)	-	44.2 e	101 d
<i>Ficus carica</i> (leaves)	-	23.2 b	106 d
<i>Ficus trigona</i> (barks)	-	41.0 d	78 b
<i>Glechoma hederacea</i> (leaves)	-	46.7 e	93 c
<i>Guazuma ulmifolia</i> (barks)	-	42.2 d	87 c
<i>Jatropha curcas</i> (flowers)	-	46.0 e	125 e
<i>Ocimum basiculum</i> (leaves)	-	41.7 d	108 d
<i>Origanum vulgare</i> (leaves)	-	40.5 d	104 d
<i>Plantago lanceolata</i> (leaves)	-	34.5 c	105 d
<i>Ruta graveolens</i> (flowers and leaves)	+	34.5 c	72 b
<i>Trichilia clausenii</i> (barks)	-	46.2 e	87 c
<i>Trichilia hirta</i> (barks)	-	39.2 d	85 c
Chlorotalonil and copper oxychloride ^d	+	3.5 a	55 a
Azoxistrobin ^e	+	7.2 a	52 a
1% Tween 80	-	44.0 e	100 d
Coefficient of variation		11.2%	6.5%

^aExcept for *Ficus trigona* (pH 6.0), 3.3 mg/ml plant extracts solutions in aqueous 1% (g/ml) Tween 80 presented pH 5.0.

^b(+): Absence of mycelium growth, (-): Presence of mycelium growth.

^cMeans of four replicates with the same letter in a column are not statistically significant ($P \leq 0.05$) according to the Scott-Knott method.

^d0.87 and 1.76 g/l, respectively.

^e0.08 g/l.

Table 3

Plant extract and fungicide effects on Murcott tangor fruits inoculated with *Alternaria alternata*.

Treatments ^a	ABS dr ^b	
	8 days	12 days
<i>Anadenanthera colubrina</i> (barks)	0.38 aA	0.49 aB
<i>Artemisia annua</i> (leaves)	1.11 bA	1.10 bA
<i>Ruta graveolens</i> (flowers and leaves)	1.00 bB	0.86 bA
<i>Cariniana estrellensis</i> (barks)	0.89 bA	1.02 bB
<i>Ficus carica</i> (leaves)	0.94 bA	0.98 bA
Chlorotalonil and copper oxychloride ^c	0.63 aA	0.57 aA
Azoxistrobin ^d	0.61 aA	0.70 aB
1.0% Tween 80 ^e	1.00 b	1.00 b
Coefficient of variation	22.4%	21.8%

^a3.3 mg/ml plant extracts solutions in aqueous 1% (g/ml) Tween 80.

^bPreviously to the statistical analysis, data were converted into spots development rate (ABS dr) by dividing the mean value for each fruit by the mean value of all fruits treated only with conidia in 1% (v/v) Tween 80 (negative control). Means with the same lower case letter in a column and the same capital letter in a line are not statistically significant ($P \leq 0.05$) according to the Scott-Knott method.

^c0.87 and 1.76 g/l, respectively.

^d0.08 g/l.

^e10⁶ conidia/ml in a 1.0% (g/ml) Tween 80 solution.




Figure 1. Scanning electron micrograph details of *Alternaria alternata* conidia: A) Severe conidia shrinking caused by exposition to *Ruta graveolens* extract (3.3 mg/ml). An arrow shows the rings (septa) enhanced by whither of cells. B) Conidia shrinking after treatment with Azoxistrobin (0.08 g/l), but with less intensity than the observed for the *R. graveolens* extract. C and D) Conidia after exposition to 1% Tween 80. No deformity was observed.

observed on fruits treated with only 1% (v/v) Tween 80 solution.

Discussion

The screening for plants with antifungal activity in the present study started with the fungal growth assay, during which some forms of action of antifungal substances can apparently be detected (Fig. 1). Although only three extracts were active against the fungus, 20 extracts (3 active and 17 inactive) were submitted to the other two *in vitro* assays to facilitate the comparison of results. Most of the inactive extracts during the initial screening (fungal growth assay) inhibited the conidia germination. Analogously, some inactive extracts during the fungal growth assay inhibited the microorganism largely in the mycelial growth assay, showing that antifungal activity of the metabolites produced by the same plant can vary according to the phase of fungal development.

Considering all the *in vitro* assays, the most active extract was that from *A. colubrina*, which is a native plant from Caatinga (Brazilian savanna), popularly known as *angico*, that occurs in the northeast region of Brazil.¹ The ability of this extract to reduce mycelial growth of *F. oxysporum* f.sp. *tracheiphilum* when used alone or in combination with fungicides has recently been described by Silva et al.¹⁹ The authors suggested polyphenols as the active substances in the *A. colubrina* extract, since these compounds could form complexes with proteins and polysaccharides, inactivating enzymes essential for fungal growth.⁸ Furthermore, the barks of *A. colubrina* are rich in tannins, which may confer antimicrobial properties to the plant extract.^{6,13} The inexistence of conidia in the SEM image of the sample treated with the *A. colubrina* extract corroborates the strong inhibition of conidial germination by this plant, since without the germ tubes, conidia would be lost during the several washings or expositions to liquids used to prepare the sample for SEM analyses.

The inhibition of the fungus by the extracts of *A. annua* and *R. graveolens* in all *in vitro* assays is in accordance with the studies

carried out by Soylu et al.²⁰ and Meepagala et al.,¹² respectively. These authors have reported the activity of *A. annua* against *Sclerotinia sclerotiorum*, *Botrytis cinerea*, *Phytophthora infestans*, and *Verticillium dahliae*, and also antifungal activity of *R. graveolens* extracts against *Colletotrichum fragariae*, *C. gloeosporioides*, *C. acutatum*, and *B. cinerea*.

Furthermore, SEM analyses showed shrinkage of those conidia exposed to *R. graveolens*, which may be attributed to irreversible ultra structural changes.¹⁴ Regarding *A. annua*, the shorter germ tubes observed by SEM analyses suggest that the plant extract was able to delay fungal germination.

Although less efficiently than the above-mentioned extracts, those obtained from *C. estrellensis* and *F. carica* also inhibited the fungus. Although no report about antifungal activity for the former plant is known, the later was reported by Aqil and Ahmad as inhibitor of *Fusarium chlamydosporum*.²

The results obtained in the test with fruits inoculated with *A. alternata* suggest consistency between *in vitro* and *in vivo* assays, since the extract of *A. colubrina* was the most active in all experiments, reducing the effects of the fungus on fruits to levels comparable to those observed for the commercial fungicides. Regarding the other extracts employed in the experiments with fruits, their inactivity against the disease seems to correlate with their low to moderate antifungal activity observed during the *in vitro* assays. Thus, the fungal growth assay is a good protocol to start a screening program to select plant extracts active against *A. alternata*.

The results obtained in the present study suggests the fungal growth assay as a good method to be used in an initial step of a screening program aimed to select plant extracts for the control of ABS in Murcott tangor fruits. Among the plants studied, the most promising was *A. colubrina*, the extract of which reduced the development of *A. alternata* on such fruits to levels statistically equal to those observed for commercial fungicides. Consequently, *A. colubrina* seems to present a great potential for the development of new products for ABS control.




Figure 2. Effects of plant extracts and fungicides on the development of *Alternaria* brown spots in Murcott tanger fruits eight days after inoculation (10^6 *Alternaria alternata* conidia/ml): A) Chlorotalonyl and copper oxychloride at 0.87 and 1.76 g/l, respectively. B) Azoxistrobin at 0.08 g/l. C) *Ficus carica*. D) *Cariniana estrellensis*. E) *Ruta graveolens*. F) *Anadenanthera colubrina*. G) *Artemisia annua* extract at 3.3 mg/ml. H) 10^6 conidia/ml resuspended in Tween 80 solution 1.0% (v/v).

Financing

The authors gratefully acknowledge CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico), CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) and FAPEMIG (Fundação de Amparo à Pesquisa do Estado de Minas Gerais) for financial support and fellowships.

Conflict of interest

The authors report no conflict of interest.

Acknowledgements

The authors also express their sincere thanks to Dr. Paulo César Lima (Departamento de Ciências Exatas, Universidade Federal de Lavras) for helping with the statistical analysis.

References

- Albuquerque UP, Medeiros PM, Almeida ALS, Monteiro JM, Lins Neto EMF, Melo JG, et al. Medicinal plants of the *caatinga* (semi-arid) vegetation of NE Brazil: A quantitative approach. *J Ethnopharmacol.* 2007;114:325–54.
- Aqil F, Ahmad I. Broad-spectrum antibacterial and antifungal properties of certain traditionally used Indian medicinal plants. *World J Microb Biot.* 2003;19:653–7.
- Bowers JH, Locke JC. Effect of botanical extracts on the population density of *Fusarium oxysporum* in soil and control of Fusarium wilt in the greenhouse. *Plant Dis.* 2000;84:300–5.
- Bozzola JJ, Russel LD. *Electron Microscopy*. 2nd ed. Boston: Jones and Bartlett Publishers; 1998. p. 670.
- Carvalho DDC, Alves E, Batista TRS, Camargos RB, Lopes EAGL. Comparison of methodologies for conidia production by *Alternaria alternata* from citrus. *Braz J Microbiol.* 2008;39:792–8.
- Desmarchelier C, Romão RL, Coussio J, Ciccio G. Antioxidant and free radical scavenging activities in extracts from medicinal trees used in the 'Caatinga' region in northeastern Brazil. *J Ethnopharmacol.* 1999;67:69–77.
- Ferreira DF. Análises estatísticas por meio do Sisvar para Windows versão 4.0. *Reunião Anual da Região Brasileira da Sociedade Internacional de Biometria*, 45. São Carlos: Universidade Federal de São Carlos; 2000. p. 255–258.
- Haslam E. Natural polyphenols (vegetable tanins) as drugs: possible modes of action. *J Nat Prod.* 1996;59:205–15.
- Latha P, Anand T, Ragupathi N, Prakasam V, Samiyappan R. Antimicrobial activity of plant extracts and induction of systemic resistance in tomato plants by mixtures of PGPR strains and Zimmu leaf extract against *Alternaria solani*. *Biol Control.* 2009;50:85–93.
- Lindsey KL, Staden JV. Growth inhibition of plant pathogenic fungi by extracts of *Allium sativum* and *Tulbaghia violacea*. *South African J Bot.* 2004;70:671–3.
- McFayden REC. Biological control of weeds. *Annu Rev Entomol.* 1998;43:369–93.
- Meepagala KA, Schrader KK, Wedge DE, Duke SO. Algicidal and antifungal compounds from the roots of *Ruta graveolens* and synthesis of their analogs. *Phytochemistry.* 2005;66:2689–95.
- Monteiro JM, Albuquerque UP, Araújo EL. Taninos: uma abordagem da química a ecologia. *Quím Nova.* 2005;28:892–6.
- Pärvi M, Pärvi AE, Craciun C, Barbu-Tudoran L, Tamas M. Antifungal activities of *Chelidonium majus* extract on *Botrytis cinerea* in vitro and ultrastructural changes in its conidia. *J Phytopathol.* 2008;156:550–2.
- Peever TL, Su G, Carpenter-Boggs L, Timmer LW. Molecular systematics of citrus-associated *Alternaria* spp. *Mycologia.* 2004;96:119–34.
- Peres NA, Timmer LW. Evaluation of the Alter-Rater model for spray timing for control of *Alternaria* brown spot on Murcott tanger in Brazil. *Crop Prot.* 2006;25:454–60.
- Peres NAR, Agostini JP, Timmer LW. Outbreaks of *Alternaria* brown spot of citrus in Brazil and Argentina. *Plant Dis.* 2003;87:750.
- Prusky D, Kobiler I, Akerman M, Miyara I. Effect of acidic solutions and acidic prochloraz on the control of postharvest decay caused by *Alternaria alternata* in mango and persimmon fruit. *Postharvest Biol Technol.* 2006;42:134–41.
- Silva JA, Pegado CMA, Ribeiro VV, Brito NM, Nascimento LC. Efeito de extratos vegetais no controle de *Fusarium oxysporum* f.sp. *tracheiphilum* em sementes de caupi. *Ciênc Agrotec.* 2009;33:611–6.
- Soylu EM, Yigitbas H, Tok FM, Soylu S, Kurt S, Baysal O, et al. Chemical composition and antifungal activity of the essential oil of *Artemisia annua* L. against foliar and soil-borne fungal pathogens. *J Plant Dis Protect.* 2005;112:229–39.
- Teegne G, Pretorius JC, Swart WJ. Antifungal properties of *Agapanthus africanus* L. extracts against plant pathogens. *Crop Prot.* 2008;27:1052–60.
- Thangavelu R, Sundararaju P, Sathiamoorthy S. Management of anthracnose disease of banana caused by *Colletotrichum musae* using plant extracts. *J Hort Sci Biotechnol.* 2004;79:664–8.
- Timmer LW, Peever TL, Solel Z, Akimitsu K. *Alternaria* diseases of citrus—novel pathosystems. *Phytopathol Mediterr.* 2003;42:99–112.
- Vicent A, Armengol J, Sales R, Garcia-Jimenez J, Alfaro-Lassala F. First report of *alternaria* brown spot of citrus in Spain. *Plant Dis.* 2000;84:1044.
- Vicent A, Badal J, Asensi MJ, Sanz N, Armengol J, Garcia-Jimenez J. Laboratory evaluation of citrus cultivars susceptibility and influence of fruit size on Fortune mandarin to infection by *Alternaria alternata* pv. *citri*. *Eur J Plant Pathol.* 2004;110:245–51.