



## Original article

## *Flammulina velutipes*: An option for “alperujo” use



Maximiliano Rugolo, Laura Levin, Bernardo Ernesto Lechner\*

Laboratorio de Micología, PROPLAME-PRHIDEB-CONICET, Departamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, 1428, CABA, Buenos Aires, Argentina

## ARTICLE INFO

## Article history:

Received 17 April 2015

Accepted 23 December 2015

Available online 26 July 2016

## Keywords:

*Flammulina velutipes*

Alperujo

Mushroom production

Lignocellulolytic enzymes

## ABSTRACT

**Background:** Two-phase olive-mill wastes (or “alperujo”) exhibit highly phytotoxic properties, mainly due to phenols. A valuable option for alperujo is its agricultural use, provided that no phytotoxic effects occur.

**Aims:** The present investigation was aimed at evaluating the efficacy of two strains of the lignin-degrading fungus *Flammulina velutipes* to colonize alperujo in order to produce edible mushrooms and to achieve its detoxification.

**Methods:** Some important cultural characters related to mushroom production (earliness, biological efficiency and quality of basidiomes) were estimated. The production of lignocellulolytic enzymes, phenol removal and detoxification of the substrate was evaluated.

**Results:** High biological efficiencies (70.8%) were obtained at 12 °C with *F. velutipes* strain BAFC 670/06 in a substrate containing poplar wood shavings and 90% of alperujo. The nature of the substrate did not seem to exert an important influence on pileus and stem morphology; nevertheless shortest stems were observed at higher temperatures. Endo- $\beta$ -1,4-glucanase, endo- $\beta$ -1,4-xylanase, laccase and Mn-peroxidase activities were detected in the extracts recovered from the solid-state cultures. Both *F. velutipes* strains were effective in removing the phenolic compounds. The initial concentration in the substrate with 90% alperujo was reduced in the case of *F. velutipes* BAFC 1763 by 84.31%, and 40.15% by *F. velutipes* BAFC 670/06. Germinability experiments on *Raphanus sativus*, showed that alperujo phytotoxicity was significantly reduced by *F. velutipes* cultures.

**Conclusions:** The experimented changes by the spent mushroom substrate resulting from *F. velutipes* cultivation with high amount of alperujo would allow its reuse for agricultural purposes.

© 2016 Asociación Española de Micología. Published by Elsevier España, S.L.U. All rights reserved.

## *Flammulina velutipes*: una opción para el aprovechamiento del alperujo

## RESUMEN

## Palabras clave:

*Flammulina velutipes*

Alperujo

Producción de basidiomas

Enzimas lignocelulolíticas

**Antecedentes:** El alperujo, subproducto de las almazaras durante la extracción del aceite de oliva, tiene propiedades fitotóxicas importantes debido a su alto contenido fenólico. Su utilización en la agricultura es una opción válida, pero deben eliminarse sus efectos fitotóxicos.

**Objetivos:** El objetivo de este trabajo fue evaluar la eficacia de dos cepas del hongo ligninolítico *Flammulina velutipes* de crecer en sustratos con alto contenido de alperujo, detoxificarlo y producir basidiomas comestibles.

**Métodos:** Se estudiaron las principales características relacionadas con el cultivo para la producción de basidiomas: tiempo de aparición de primordios, eficiencia biológica y calidad. Se evaluó la producción de enzimas lignocelulolíticas, la remoción de compuestos fenólicos y la detoxificación del sustrato.

**Resultados:** Se obtuvieron altos valores de eficiencia biológica (70,8%) a 12 °C con la cepa BAFC 670/06 de *F. velutipes* en un sustrato compuesto de viruta de álamo y 90% de alperujo. La naturaleza del sustrato al parecer no ejerció influencias importantes en la morfología de los basidiomas, aunque a altas temperaturas los estípites presentaron una menor longitud. Se detectó actividad endo- $\beta$ -1,4-glucanasa, endo- $\beta$ -1,4-xilanas, lacasa y Mn-peroxidasa en extractos recuperados de cultivos en estado sólido.

\* Corresponding author.

E-mail address: [bernardoelechner@gmail.com](mailto:bernardoelechner@gmail.com) (B.E. Lechner).

Ambas cepas fueron efectivas en la reducción del contenido de fenoles del sustrato, reducción que alcanzó el 84,31% con *F. velutipes* BAFC 1763 y el 40,15% con *F. velutipes* BAFC 670/06. Los ensayos de germinación de semillas de *Raphanus sativus* mostraron una significativa reducción de la fitotoxicidad del alperujo.

**Conclusiones:** Los cambios experimentados por el sustrato remanente del cultivo de *F. velutipes* con altas concentraciones de alperujo podrían permitir su reutilización con fines agrícolas.

© 2016 Asociación Española de Micología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

The white-rot fungus *Flammulina velutipes* is a xylophagous fungus belonging to the Physalacriaceae family, with health and medicinal benefits, widely cultivated in East Asia.<sup>37</sup> *F. velutipes* is one of the six most cultivated mushroom species in the world; over 300,000 t of this mushroom are produced per year.<sup>28</sup> It is particularly known for its taste, and preventive as well as curative properties for liver diseases and gastro enteric ulcers. In addition, it has also been reported to contain immunomodulatory, antitumor and antibiotic substances.<sup>37</sup> Its distribution is limited to temperate zones of the world because a cold period is required for fruiting.<sup>37</sup> Its genome has been recently sequenced and reveals a high capacity for lignocellulose degradation.<sup>25</sup> Moreover *F. velutipes* strain Fv-1 demonstrated its potential for the conversion of lignocellulosic biomass to ethanol by consolidated bioprocessing, a methodology that combines enzyme production, enzymatic saccharification and ethanol fermentation in one step, thus reducing the total cost of bioethanol production.<sup>16</sup>

Since *F. velutipes* grows on wood in nature, mixtures of lignocellulosic materials have been utilized as substrates in the commercial production of the mushroom. A common substrate used for its production is sawdust with agricultural residues, such as corncobs, cottonseed husk, sugarcane bagasse, etc.<sup>37</sup>

The olive oil extraction industry produces high amounts of olive wastes. One emerging technology for olive-oil extraction consists of a continuous centrifugation two-phase process that generates a liquid phase (olive-oil) and a semi-solid organic waste (alperujo), which are then dried and extracted with solvents to obtain an extra yield of oil and a dry olive mill residue (DOR).<sup>34</sup> Alperujo is a fibrous lignocellulosic paste which has a water content of about 65%, a slightly acidic pH and a very high content of organic matter (90.66%), mainly composed of lignin, hemicellulose and cellulose (38.82%, 29.70% and 23.47%, w/w, of total organic matter, respectively). It has also a considerable proportion of N [TN (g·kg<sup>-1</sup>) 11.99] and phenolic compounds (1.36%, w/w, of total organic matter).<sup>21</sup>

To preserve the ecology of olive oil producer's countries, degradation of these wastes have been studied using thermal, physico-chemical processes and biological treatments.<sup>21</sup> Several studies evaluated its detoxification through solid state fermentation and the addition of appropriate co-composting materials to deal with the particular structure of this effluent, using aerobic or anaerobic microorganisms. Apart from the exploitation of mixed microbial communities present throughout the composting process, the inoculation with white-rot basidiomycetes has also been used to decrease the contents of phenolics and phytotoxicity of the alperujo.<sup>2,13,34</sup> White-rot fungi are able to secrete specific ligninolytic enzymes causing significant phenolic removal.<sup>31</sup> Moreover, if edible mushrooms were utilized, a double goal could be achieved, decreasing the toxicity of this waste by means of the ligninolytic enzymes secreted by the fungus, while producing appreciated basidiomes.<sup>42</sup>

Up to now, the use of by-products from the olive-oil industry for mushrooms production has been scarcely investigated.<sup>43</sup> Olive-mill wastewater and olive-press cake were used as substrates for *Pleurotus* cultivation,<sup>10,42</sup> olive-mill by-products were

used for *Agaricus bisporus* commercial exploitation,<sup>1</sup> and alperujo composted with wheat straw was evaluated as substrate for basidiome production of the edible mushrooms *Agrocybe cylindracea*, *Pleurotus cystidiosus*, *Pleurotus eryngii*, *Pleurotus ostreatus* and *Pleurotus pulmonarius*.<sup>30,43</sup>

The aims of this work were: (i) to study the capacity of two *F. velutipes* strains to grow with different alperujo concentrations and temperatures while producing lignocellulolytic enzymes, (ii) to evaluate if alperujo concentration influences yield values and morphological properties of the basidiomes obtained, and (iii) to assess the ability of *F. velutipes* for phenol removal and detoxification in this substrate.

## Materials and methods

### Fungal strains and culture conditions

*F. velutipes* strain BAFC 1763 (Fungal Collection of the Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires) and BAFC 670/06 (IIB-INTECH Collection of Fungal Cultures) were used. Both strains were maintained in Petri dishes at 4 °C on ME agar (malt extract 1.2%, glucose 1%, and agar 2%).

### Spawn production

Spawn production was prepared in 350 ml glass jars filled with boiled wheat grains (*Triticum durum*) and 1% (w/w) calcium carbonate (CaCO<sub>3</sub>). Jars were sterilized for 1.5 h at 121 °C, cooled and inoculated with a 1 cm diameter plug of mycelium grown on Nobles' medium,<sup>23</sup> and then they were incubated at 25 °C, in the dark, with periodical shaking. Time required for spawn production was recorded.

### Substrate preparation

Polypropylene bags of 20 cm × 40 cm were filled with 100 g (dry weight) of substrate [*Populus* wood shavings, alperujo (0%, 30%, 60% or 90%, w/w), and 3% CaCO<sub>3</sub>]. Humidity was adjusted (w/w) to 75%. Bags were stopped with cotton plugs held by PVC (polyvinyl chloride) cylinders and autoclaved at 121 °C for 2 h. After cooling, bags were inoculated with 5% (wet weight) of spawn, and were incubated at 25 °C in the dark until total substrate colonization (21–25 days).

### Basidiomes induction

After total colonization by the mycelium, polypropylene bags with the upper surface opened (with a neck of 10 cm approx. created from the same bag) were transferred to the basidiome production rooms. Cropping conditions to induce basidiome formation were 12 h light/12 h dark photoperiod (2000 lx of fluorescent light), 75–85% humidity levels and watering by automatic spray provided for 2 min, 3 times every day. Three different temperatures were assayed for basidiome production: 6 ± 1 °C, 12 ± 1 °C and 22 ± 1 °C.

### Cropping period, crop yield and morphological traits assessment

Up to three flushes were harvested during the cropping period (80 days between the induction day and the last harvest) in the different substrates. Mature basidiomes (80% of expanded pileus) were collected manually. The parameters to test the suitability of the substrates under study for the cultivation of *F. velutipes* were (a) earliness, defined as the time elapsed between the day of inoculation and the day of primordial formation, (b) biological efficiency (BE), calculated as the percentage ratio of fresh mushrooms weight over the dry weight of the substrate, and (c) morphological quality traits: (i) pileus width and length and (ii) stem length and diameter.

### Analytical determinations

#### Enzyme activities


Laccase activity (E.C.:1.10.3.2) was measured using 2,6-dimethoxyphenol (DMP) 5 mM in 0.1 M sodium acetate buffer (pH 3.6) at 50 °C. Oxidation of DMP was determined by the increase in  $A_{469}$  ( $\epsilon_{469} = 27 \text{ mM} \cdot \text{cm}^{-1}$ ).<sup>26</sup> Manganese peroxidase activity (MnP) (E.C.:1.11.1.13) was measured using phenol red as the substrate in 0.1 M sodium dimethyl succinate buffer (pH 4.5) ( $\epsilon_{610} = 22 \text{ mM} \cdot \text{cm}^{-1}$ ) at 50 °C.<sup>6</sup> Endo- $\beta$ -1,4-glucanase (E.C.:3.2.1.4) and endo- $\beta$ -1,4-xylanase (E.C.:3.2.1.8) activities were determined by measuring the reducing sugars released from carboxymethyl-cellulose or oat xylan, respectively, as substrates, in 50 mM sodium acetate buffer, pH 4.8. Liberated reducing sugars were quantified by the Somogyi–Nelson method,<sup>22</sup> using either glucose or xylose as standards.<sup>40</sup> Cellulolytic and xylanolytic activities were determined at 50 °C. Enzyme activity was expressed in International Units (U) as the amount of enzyme required to release 1  $\mu\text{mol}$  of product in 1 min. In terms of production, the activity was defined as  $\text{U} \cdot \text{g}^{-1}$  dry residue (substrate plus mycelium). Samples of substrates colonized by mycelium were collected at different stages of the solid-state fermentation: T1, after complete colonization of the substrate; T2, after the end of first flush; T3, after the end of the second flush (if present); T4, spent substrate after the last mushroom harvest. Crude extracts were obtained by adding distilled water to the samples from each freshly harvested culture (5:1, w/w), stirring for 20 min, followed by filtration and centrifugation. The supernatants were stored at 20 °C until needed.

#### Total phenolic compounds quantification

Dried solid samples of non-inoculated and spent substrates were extracted with methanol:water 80:20 (v/v) (5:1, w/w), stirring for 2 h at 120 rpm, followed by filtration and centrifugation. The quantification of the total phenolic compounds was based on the Folin–Ciocalteu reaction, according to the method of Singleton et al.,<sup>38</sup> measuring the absorbance at 765 nm. Gallic acid was used as standard for the quantification.

#### Phytotoxicity bioassay

The phytotoxicity of the extract obtained from *Flammulina*'s spent mushroom was determined according to the method described by Zucconi et al.<sup>44</sup> using *Raphanus sativus* seeds incubated for 48 h at 28 °C with either (a) control:water; (b) the extract obtained from the non-inoculated substrate and (c) the extract obtained from the spent substrate after the last mushroom harvest. The germination index (GI) was calculated according to the expression:  $\text{GI} = [(G/\text{Go})/(L/\text{Lo})] \times 100$ , where Go and Lo are, respectively, the germination percentage and radicle growth of the control. Crude extracts were obtained by adding distilled water to the



**Fig. 1.** Biological efficiencies attained by *F. velutipes* strain 670/06 with different alperujo concentrations and temperatures. Values with the same letter are not statistically different according to Tukey's test.

samples from each freshly harvested culture (5:1, w/w), stirring for 20 min, and followed by filtration and centrifugation.

#### Experimental design and statistical treatments

For basidiome production each treatment consisted of ten replicates. Data of analytical determinations are the average of the results of at least three replications with a standard error of less than 5%. Analysis of variance (ANOVA) and repeated measures analysis were tested by the software Statistica v.7.0. The assumptions of normality and homogeneity of variances were checked by means of Kolmogorov–Smirnov and Bartlett tests, respectively, for the validity of ANOVA method. The significant differences between treatments were compared by Tukey's test at 5% level of probability.

### Results

#### Effect of alperujo addition and temperature of incubation on *F. velutipes* basidiome production

Both strains of *F. velutipes* took 21–25 days to complete the colonization of all the substrates assayed. Growth was assessed visually, and while a dense mycelial mat could be observed in the substrates supplemented with alperujo, only scatter hyphae were seen on the control with solely poplar wood shavings. Lignocellulosic materials are generally low in protein content, insufficient for mushroom cultivation. Since the C:N ratio plays an important role in spawn running and the growth of basidiomes, nitrogen supplementation is an important factor for the growth and yield of mushrooms. The high content of organic matter provided by the alperujo might support better mycelial growth.

*F. velutipes* strain BAFC 670/06 demonstrated an exceptionally good adaptation to olive mill waste containing substrates since most of the alperujo media provided significantly higher BEs values in comparison to the control. Moreover, with alperujo addition mushroom primordia were formed within a shorter time period after substrate inoculation; the first flush was observed between 13 and 33 days after initial induction (Table 1). The highest BEs were obtained at 12 °C with *F. velutipes* strain BAFC 670/06, and BEs improved with increasing alperujo concentrations up to a maximum of 70.8 with 90% of this olive waste (Fig. 1). No significant differences were observed in BEs obtained with substrates containing 60 and 30% alperujo (46.60% and 38.68%, respectively) at 12 °C. More than 60, 75 and 90% of the total production was cropped during the first flush with 90%, 60% and 30% alperujo content,

**Table 1**Effect of alperujo addition and temperature of induction on *F. velutipes* strain 670/06 basidiome production.

| Temp. | Alperujo concentration | Earliness (days)          | Time between induction and first flush | First flush (g) | Second flush (g)                              | Pileus                     |                            | Stem                        |                             |
|-------|------------------------|---------------------------|----------------------------------------|-----------------|-----------------------------------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|
|       |                        |                           |                                        |                 |                                               | Width (cm)                 | Length (cm)                | Width (cm)                  | Length (cm)                 |
| 6 °C  | 0%                     | 50.50 ± 2.38 <sup>a</sup> | 30.25 ± 1.89 <sup>a</sup>              | 11.80 ± 7.54    | –                                             | 2.53 ± 0.73 <sup>a</sup>   | 2.38 ± 0.67 <sup>a</sup>   | 0.33 ± 0.09 <sup>bcd</sup>  | 10.67 ± 2.50 <sup>ab</sup>  |
| 6 °C  | 30%                    | 45.50 ± 1.00 <sup>a</sup> | 25.50 ± 1.00 <sup>a</sup>              | 16.80 ± 6.72    | 2.50 ± 1.73                                   | 1.43 ± 0.51 <sup>e</sup>   | 1.30 ± 0.46 <sup>f</sup>   | 0.25 ± 0.06 <sup>cdef</sup> | 9.46 ± 2.99 <sup>a</sup>    |
| 6 °C  | 60%                    | 41.80 ± 1.09 <sup>c</sup> | 21.80 ± 1.10 <sup>c</sup>              | 20.60 ± 5.94    | 5.50 ± 3.42                                   | 1.53 ± 0.56 <sup>e</sup>   | 1.41 ± 0.54 <sup>f</sup>   | 0.32 ± 0.17 <sup>fg</sup>   | 8.81 ± 2.84 <sup>bcd</sup>  |
| 6 °C  | 90%                    | 50.20 ± 1.09 <sup>b</sup> | 30.20 ± 1.10 <sup>b</sup>              | 34.17 ± 4.54    | –                                             | 2.00 ± 0.40 <sup>bcd</sup> | 1.81 ± 0.37 <sup>bcd</sup> | 0.36 ± 0.10 <sup>abc</sup>  | 11.42 ± 1.37 <sup>a</sup>   |
| 12 °C | 0%                     | 46.5 ± 1.00 <sup>b</sup>  | 26.50 ± 1.00 <sup>b</sup>              | 22.00 ± 1.41    | 1.40 ± 0.36                                   | 2.00 ± 0.37 <sup>cd</sup>  | 1.84 ± 0.38 <sup>cde</sup> | 0.26 ± 0.06 <sup>defg</sup> | 10.76 ± 2.10 <sup>ab</sup>  |
| 12 °C | 30%                    | 34.80 ± 1.09 <sup>c</sup> | 14.80 ± 1.10 <sup>c</sup>              | 36.50 ± 6.45    | 3.57 ± 1.69                                   | 2.59 ± 0.57 <sup>ab</sup>  | 2.42 ± 0.55 <sup>ab</sup>  | 0.38 ± 0.11 <sup>ab</sup>   | 12.27 ± 2.61 <sup>ab</sup>  |
| 12 °C | 60%                    | 42.40 ± 1.34 <sup>c</sup> | 22.40 ± 1.34 <sup>c</sup>              | 34.67 ± 3.72    | 11.60 ± 2.30<br>(third flush:<br>3.40 ± 0.34) | 2.26 ± 0.72 <sup>abc</sup> | 2.12 ± 0.66 <sup>abc</sup> | 0.34 ± 0.11 <sup>bcd</sup>  | 10.34 ± 2.46 <sup>abc</sup> |
| 12 °C | 90%                    | 40.33 ± 0.82 <sup>c</sup> | 20.33 ± 0.82 <sup>c</sup>              | 44.60 ± 2.70    | 26.20 ± 2.77                                  | 2.19 ± 0.54 <sup>cd</sup>  | 2.00 ± 0.50 <sup>cde</sup> | 0.39 ± 0.14 <sup>bcd</sup>  | 9.97 ± 2.41 <sup>cde</sup>  |
| 22 °C | 0%                     | 35.2 ± 1.09 <sup>c</sup>  | 15.20 ± 1.10 <sup>c</sup>              | 1.40 ± 0.55     | –                                             | 1.67 ± 0.39 <sup>de</sup>  | 1.42 ± 0.36 <sup>ef</sup>  | 0.23 ± 0.05 <sup>g</sup>    | 7.16 ± 1.73 <sup>fg</sup>   |
| 22 °C | 30%                    | 45.83 ± 0.75 <sup>d</sup> | 25.83 ± 0.75 <sup>d</sup>              | 9.60 ± 4.16     | –                                             | 2.08 ± 0.58 <sup>cde</sup> | 1.79 ± 0.51 <sup>def</sup> | 0.33 ± 0.11 <sup>efg</sup>  | 7.41 ± 1.55 <sup>g</sup>    |
| 22 °C | 60%                    | 40.86 ± 1.07 <sup>d</sup> | 20.86 ± 1.07 <sup>d</sup>              | 20.40 ± 3.85    | –                                             | 2.26 ± 0.61 <sup>bcd</sup> | 1.96 ± 0.52 <sup>cd</sup>  | 0.40 ± 0.12 <sup>abc</sup>  | 8.52 ± 1.63 <sup>efg</sup>  |
| 22 °C | 90%                    | 35.33 ± 1.03 <sup>d</sup> | 14.33 ± 1.03 <sup>d</sup>              | 21.33 ± 3.50    | –                                             | 1.90 ± 0.63 <sup>de</sup>  | 1.64 ± 0.46 <sup>def</sup> | 0.48 ± 0.16 <sup>a</sup>    | 9.73 ± 1.96 <sup>def</sup>  |

Data are the mean ± standard deviation of three determinations. Means followed by the same letter are not significantly different according to Tukey's test.


respectively, at 12 °C. Lower BEs were attained at 6 °C and 22 °C: at 6 °C BEs varied from 11.8 to 34.17%, while at 22 °C BEs ranged from 1.4 up to 21.33% with different amounts of alperujo; the highest values were obtained with 90% (Fig. 1).

The lowest BEs gotten at 22 °C might be related with various aborted primordia observed. Strain 1763 only produced basidiomes on 60% of alperujo at 12 °C and 22 °C, and the BEs were less than 10% (data not shown).

Pileus width and length, and stem length and diameter, were not significantly different among most of the treatments for the same strain, and hence the nature of the substrate did not seem to exert any important influence (exception i.e. in stem morphology with 90% of alperujo). On the contrary, the temperature of incubation held sway over morphology of *F. velutipes* basidiomes, resulting in mushrooms with shorter stems at 22 °C (Table 1).

### Enzyme activities

Cellulolytic, xylanolytic and ligninolytic activities were detected in the extracts recovered from the solid-state cultures. Table 2 summarizes the highest titers of the different enzyme activities attained by both strains (and the day and medium in which these values were achieved). Time course of enzyme production in the conditions that rendered the highest BE, *F. velutipes* strain BAFC 670/06 grown at 12 °C with 90% of alperujo, is depicted in Fig. 2. Hydrolytic activities showed their highest values at initial stages of cultivation: endo-β-1,4-glucanase utmost titers detected were  $5.46 \pm 0.03 \text{ U} \cdot \text{g}^{-1}$  (30%, 12 °C) and  $5.27 \pm 0.01 \text{ U} \cdot \text{g}^{-1}$  (60%, 12 °C) for 670/06 strain, and  $7.14 \pm 0.20$  (0%, 22 °C, and 30%, 22 °C) for 1763 strain, while endo-β-1,4-xylanase peaks were  $10.16 \pm 0.06 \text{ U} \cdot \text{g}^{-1}$  (60%, 12 °C) and  $6.09 \pm 0.01 \text{ U} \cdot \text{g}^{-1}$  (90%, 12 °C) for 670/06 strain. On the contrary, laccase activity showed its highest activity at final stages, at second flush or at the end of the culture period (80 days). *F. velutipes* BAFC 1763 displayed a higher activity of laccase [ $102.66 \pm 6.02 \text{ U} \cdot \text{g}^{-1}$  (60%, 22 °C)], and MnP activity was only detected in this strain ( $1.67 \pm 0.03 \text{ U} \cdot \text{g}^{-1}$  with 90% alperujo at 22 °C). Highest laccase activity detected in strain BAFC 670/06 was  $0.98 \pm 0.14 \text{ U} \cdot \text{g}^{-1}$  with 30% alperujo at 12 °C at the end of incubation. Laccase activity was influenced by the concentration of alperujo added to the substrate. Highest laccase activities recorded for strain BAFC 1763 at 12 °C with 0, 30, 60 and 90% alperujo were respectively:  $0.64 \pm 0.03$ ;  $5.46 \pm 0.23$ ;  $21.74 \pm 1.05$  and  $27.11 \pm 2.90 \text{ U} \cdot \text{g}^{-1}$ . Strain BAFC 670/06 produced the highest laccase activity with 30% alperujo ( $0.98 \pm 0.14 \text{ U} \cdot \text{g}^{-1}$  at 12 °C); laccase activity was  $0.68 \pm 0.11 \text{ U} \cdot \text{g}^{-1}$  with 90% alperujo at the same



**Fig. 2.** Kinetics of enzyme production in the conditions that rendered the highest BE, *F. velutipes* strain BAFC 670/06 grown at 12 °C with 90% of alperujo. Each point represents the mean of three replicate experiments. S.E.M. (errors not shown were within the heights of the symbols).

temperature while it not exceeded  $0.35 \pm 0.07 \text{ U} \cdot \text{g}^{-1}$  in the control without alperujo in the same conditions.

### Removal of phenolic compounds and detoxification of the substrate by *F. velutipes*

Both *F. velutipes* strains were effective in removing the phenolic compounds. Initial phenolic concentration in substrates with 30%, 60 and 90% of alperujo were respectively 2.61, 3.14 and  $3.76 \text{ mg} \cdot \text{g}^{-1}$  (expressed as gallic acid equivalents). Phenol concentration at the end of the incubation in the substrate with 90% alperujo was reduced by 84.31% at 12 °C by *F. velutipes* BAFC 1763, and 42.28% by *F. velutipes* BAFC 670/06. Phenol removal by *F. velutipes* BAFC 1763 reached values of 93.06% and 87.26% with 30% and 60% of alperujo content, while those attained by *F. velutipes* BAFC 670/06 with 30% and 60% of alperujo at 12 °C were respectively 33.71% and 46.49%. GI of the treated substrate (with 90% of alperujo) was similar, or even higher than the control value (100%): 127% and 97% for strains BAFC 670/06 and 1763, respectively. GI value in the non-treated substrate was 26.67%.

### Discussion

This work evaluated for the first time the potential use of olive mill wastes as substrate for *F. velutipes* cultivation. *F. velutipes* strain BAFC 670/06 formed primordia in a shorter time on substrates with alperujo. On the contrary, in previous works, always olive waste



**Table 2**Extracellular lignocellulolytic activities displayed by *F. velutipes* strains on poplar wood shavings with different alperujo concentrations.

| Enzyme activity      | Strain BAFC 670/06<br>(U·g <sup>-1</sup> substrate) <sup>a</sup>                                    | Strain BAFC 1763<br>(U·g <sup>-1</sup> substrate) |
|----------------------|-----------------------------------------------------------------------------------------------------|---------------------------------------------------|
| Endo-β-1,4-glucanase | 5.46 ± 0.03 (30%, 12 °C, after 1st flush) <sup>b</sup><br>5.27 ± 0.01 (60%, 12 °C, after 1st flush) | 7.14 ± 0.20 (30%, 22 °C, initial post-incubation) |
| Endo-β-1,4-xylanase  | 10.16 ± 0.06 (60%, 12 °C, after 1st flush)<br>6.09 ± 0.01 (90%, 12 °C, after 1st flush)             | 3.46 ± 0.57 (30%, 22 °C, initial post-incubation) |
| Laccase              | 0.98 ± 0.14 (30%, 12 °C, end of incubation)                                                         | 102.66 ± 6.02 (60%, 22 °C after 1st flush)        |
| Mn-peroxidase        | ND <sup>c</sup>                                                                                     | 1.67 ± 0.03 (90%, 22 °C, 90 days)                 |

<sup>a</sup> Data are the mean ± standard deviation of three determinations.<sup>b</sup> The values shown correspond to the peak of enzyme production. Data between parentheses indicate the conditions under which these values were achieved.<sup>c</sup> ND, not detected.

amendment had a distinct adverse effect on earliness.<sup>30,42,43</sup> Taking into account the nature of the residue used to cultivate the mushroom, BEs obtained in this work are acceptable and comparable to those obtained in other publications when growing *F. velutipes* using different formulation substrates: *i.e.* 50.9% using paddy straw with additives,<sup>14</sup> 90–106% on paddy straw,<sup>39</sup> 73% with maize straw,<sup>7</sup> 56% on coffee husks<sup>12</sup> and 78% while using coffee spent grounds.<sup>12</sup>

Pileus width and length, and stem length and diameter, were not significantly different among most of the treatments for the same strain when varying the amount of alperujo. Likewise, in the cases of *P. ostreatus* and *P. pulmonarius* no significant differences were observed in the size of the mushrooms obtained when assaying substrates with different alperujo concentrations.<sup>43</sup> Seven oyster mushroom strains were cultivated in wheat straw bags supplemented with up to 90% alperujo, and most of them showed no significant differences on cultivation parameters and basidiomes quality (except for color) between the control and the substrates supplemented with up to 50% alperujo, although high alperujo concentrations resulted in a significant yield, biological efficiency and productivity decrease, retarding of pinning and flushing.<sup>30</sup> Alperujo additions affected the yield, enzyme secretion, color and texture of the *Pleurotus* specimens obtained.<sup>27</sup> In both strains of *F. velutipes* assayed in this work, alperujo addition did not cause a decrease in mushroom size.

In order to obtain the nutrients required for growth and fruiting, *F. velutipes* is assumed to secrete the hydrolytic/oxidative enzymes that catalyze the degradation of the major macromolecular components (cellulose, hemicellulose and lignin) of its growth substrate. However, compared with other cultivated mushrooms, very little is known about the nature of the lignocellulolytic enzymes produced by *F. velutipes*, the parameters affecting their production, and enzyme activity profiles during different stages of the developmental cycle.<sup>17,36</sup> The production of these enzymes is important in substrate colonization, as well as decisive in basidiome production.<sup>24</sup> Moreover the production of ligninolytic enzymes by the fungus might be involved in its capacity to grow and detoxify substrates with high contents of phenolic compounds like alperujo.

Hydrolysis of cellulose and hemicellulose provides the nutrients required for vegetative growth as a prelude to basidiome production. Endoglucanase and endoxylanase activities increased with the incubation time and also during the formation of basidiomes for both strains. Matsumoto<sup>19</sup> also found that cellulase and xylanase activities increased during the development of *Lentinus edodes* basidiomes cultivated on eucalyptus sawdust, with highest levels during mushroom maturation. Luz et al.<sup>15</sup> registered similar results when incubating *P. ostreatus* in different agro industrial wastes. Kurt and Buyukalaca<sup>9</sup> detected the highest endoglucanase activities when incubating *P. ostreatus* and *Pleurotus sajor-caju* in different agro industrial wastes after first flush and on the 5th day of mycelial growth. The increase in the enzyme activities during

basidiome production may be due to the fungus' need to mobilize large amounts of carbon for mushroom formation.<sup>18</sup>

High ligninolytic activities during growth phases in lignocellulosic substrates, with drastic reductions during the period of basidiome formation, were observed in the white-rotting basidiomycetes *P. ostreatus*,<sup>9</sup> *P. sajor-caju*,<sup>9</sup> *Lentinus tigrinus*,<sup>11</sup> *L. edodes*,<sup>18,24</sup> *A. bisporus*<sup>41</sup> and *Grifola frondosa*.<sup>20</sup> In contrast, during *Volvariella volvacea* solid state fermentation, laccase activity increased greatly after vegetative growth at basidiome initiation and stayed at high level until the basidiomes matured.<sup>4</sup> Laccase activity was influenced by the concentration of alperujo added to the substrate. Ruiz Rodriguez et al.<sup>31</sup> also found an increase in laccase and peroxidase levels secreted by *P. ostreatus* and *P. pulmonarius* on substrates supplemented with olive mill wastes than on control.

DOR and its components have been described as a source of phenols and stimulators of fungal growth that induce several enzymes such as MnP and Lac.<sup>5,29,35</sup>

Ko et al.<sup>8</sup> evaluated endoglucanase, endoxylanase and laccase activities recovered from spent mushroom compost of *F. velutipes* grown in sawdust; specific activities obtained were respectively 151, 110 and 48.5 nkat·g<sup>-1</sup>. Compared with these data, our results showed higher recovery of all the enzyme activities evaluated. Moreover, as far as we know this is the first report of MnP activity in *F. velutipes*.

Due to its content in organic matter and mineral nutrients, alperujo might be employed for agronomic purposes.<sup>2</sup> However, the main technical constraint to its biological use is the presence of a relevant phenolic fraction, the concentration of which may easily range from 12 to 26 g·kg<sup>-1</sup>, exhibiting significant phytotoxicity.<sup>32</sup> Both spent mushroom substrates under trial showed low phytotoxicity (GI > 60%) within the range of safety for agronomic use.<sup>44</sup> Thus, the spent mushroom compost derived from *F. velutipes* cultivation might be used as organic fertilizer.<sup>1</sup> *F. velutipes* BAFC 1763 removed these toxic compounds more efficiently probably by synthesizing and excreting higher titers of ligninolytic enzymes (laccase and MnP) than the strain BAFC 670/06 into the alperujo. The effect of laccase and MnP on DOR, as well as their aqueous and organic extractives, has to some extent been described *in vivo* and *in vitro*.<sup>29</sup> *Coriopsis rigida*, another white-rot fungus, is capable of transforming certain phytotoxic monomeric phenols in DOR, into non-phytotoxic polymeric phenols.<sup>3</sup> The degree of polymerization of these phenols may limit the accessibility *via* plant cell membrane, decreasing phytotoxicity.<sup>33</sup> The incubation of DOR with *Bjerkandera adusta* also produced a sharp decrease in total phenolic content in 4 weeks, as well as a reduction in phytotoxicity.<sup>29</sup>

In conclusion, the results achieved in this investigation contribute to expand the knowledge on the lignocellulolytic enzyme system of *F. velutipes*, scarcely investigated up till now. Taking into account that *F. velutipes* successfully colonized this waste, this substrate formula might be considered for its growth. The inclusion of poplar wood shavings provides nutrients and enhances gas

exchange and water retention, thus facilitating the degradation of this toxic compound.<sup>13</sup> Both *F. velutipes* strains decreased the phenolic content whilst at the same time yielded a less phytotoxic substrate. *F. velutipes* BAFC 1763 cultures in alperujo/poplar wood shavings mixtures resulted in aqueous extracts containing about 80% less phenolic compounds, which inhibited seed germination to a much lesser extent. Experiments comparing abiotic control with the fungal treated substrate containing 90% of alperujo, showed that phytotoxicity was totally suppressed in the waste that underwent 100 days of fungal treatment. The experimented changes undergone by alperujo after degradation would allow its reuse (after the harvest of the mushrooms) for agricultural purposes as organic amendments or soil conditioner for growing plants.<sup>32</sup>

### Conflict of interest

The authors declare that there are no conflicts of interest.

### Acknowledgements

The authors thank University of Buenos Aires and CONICET (Argentina) for the financial assistance.

### References

- Altieri R, Esposito A, Parati F, Lobianco A, Pepi M. Performance of olive mill solid waste as a constituent of the substrate in commercial cultivation of *Agaricus bisporus*. *Int Biodeterior Biodegrad*. 2009;63:993–7.
- Bonanomi G, Giorgi V, Giovanni DS, Neri D, Scala F. Olive mill residues affect saprophytic growth and disease incidence of foliar and soilborne plant fungal pathogen. *Agric Ecosyst Environ*. 2006;115:194–200.
- Casa R, D'Annibale A, Pieruccetti F, Stazi SR, Giovannozzi S, Sermanni G, Locascio B. Reduction of the phenolic components in olive-mill wastewater by an enzymatic treatment and its impact on durum wheat (*Triticum durum* Desf.) germinability. *Chemosphere*. 2003;50:959–66.
- Chen SC, Ma D, Ge W, Buswell JA. Induction of laccase activity in the edible straw mushroom, *Volvariella volvacea*. *FEMS Microbiol Lett*. 2003;218:143–8.
- Díaz R, Saparrat MCN, Jurado M, García-Romera I, Ocampo JA, Martínez MJ. Biochemical and molecular characterization of *Coriopsis rigida* laccases involved in transformation of the solid waste from olive oil production. *Appl Microbiol Biotechnol*. 2010;88:133–42.
- Glenn JK, Gold MH. Purification and characterisation of an extracellular Mn(II)-dependent peroxidase from the lignin degrading basidiomycete *Phanerochaete chrysosporium*. *Arch Biochem Biophys*. 1985;242:329–41.
- Ji H, Wang Q, Wang H, Chen WJ, Zhu CH, Hou H, et al. A fundamental research of mushroom cultivation using maize straw. *Edible Fungi China*. 2001;20:10–2.
- Ko HG, Park SH, Kim SH, Park HG, Park WM. Detection and recovery of hydrolytic enzymes from spent compost of four mushroom species. *Folia Microbiol*. 2005;50:103–6.
- Kurt S, Buyukalaca S. Yield performances and changes in enzyme activities of *Pleurotus* spp. (*P. ostreatus* and *P. sajor-caju*) cultivated on different agricultural wastes. *Bioresour Technol*. 2010;101:3164–9.
- Lechner BE, Monaldi S. Utilization of garlic and maize wastes supplemented with olive mill waste water for *Pleurotus ostreatus* cultivation. *Rev Mex Micol*. 2011;34:17–22.
- Lechner BE, Papinutti L. Production of lignocellulosic enzymes during growth and fruiting of the edible fungus *Lentinus tigrinus* on wheat straw. *Process Biochem*. 2006;41:594–8.
- Leifa F, Pandey A, Soccol CR. Production of *Flammulina velutipes* on coffee husk and coffee spent ground. *Braz Arch Biol Technol*. 2001;44:205–12.
- Linares A, Caba JM, Ligerio F, de la Rubia T, Martínez J. Detoxification of semisolid olive-mill wastes and pine-chip mixtures using *Phanerochaete flavidio-alba*. *Chemosphere*. 2003;51:887–91.
- Lu QG, Gui YW, Tong XL. Mycelium growth and mushroom yield of *Flammulina velutipes* on different culture media. *Jiangsu Agric Sci*. 1989;3:26–7.
- Luz JMR, Nunes MD, Paes SA, Torres DP, Silva MCS, Kasuya MCM. Lignocellulolytic enzyme production of *Pleurotus ostreatus* growth in agroindustrial wastes. *Braz J Microbiol*. 2012;43:1508–15.
- Maehara T, Ichinose H, Furukawa T, Ogasawara W, Takabatake K, Kaneko S. Ethanol production from high cellulose concentration by the basidiomycete fungus *Flammulina velutipes*. *Fungal Biol*. 2013;117:220–6.
- Mallerman J, Papinutti L, Levin L. Characterization of  $\beta$ -glucosidase produced by the white rot fungus *Flammulina velutipes*. *JMB*. 2015;25:57–65.
- Mata G, Savoie JM. Extracellular enzyme activities in six *Lentinula edodes* strains during cultivation in wheat straw. *World J Microbiol Biotechnol*. 1998;14:513–9.
- Matsumoto T. Changes in activities of carbohydrases, phosphorylase, proteinases and phenol oxidases during fruiting of *Lentinula edodes* in sawdust cultures. *Tottori Mycol Inst*. 1998;26:46–54.
- Montoya S, Orrego CE, Levin L. Growth, fruiting and lignocellulolytic enzyme production by the edible mushroom *Grifola frondosa* (maitake). *World J Microbiol Biotechnol*. 2012;28:1533–41.
- Morillo JA, Antizar-Ladislao B, Monteoliva-Sánchez M, Ramos-Cormenzana A, Russell NJ. Bioremediation and biovalorisation of olive-mill wastes. *Appl Microbiol Biotechnol*. 2009;82:25–39.
- Nelson NJ. A photometric adaptation of the Somogyi method for the determination of glucose. *J Biochem*. 1944;153:375–80.
- Nobles MK. Identification of cultures of wood-inhabiting Hymenomycetes. *Can J Bot*. 1965;43:1097–139.
- Ohga S, Royse DJ. Transcriptional regulation of laccase and cellulase genes during growth and fruiting of *Lentinula edodes* on supplemented sawdust. *FEMS Microbiol Lett*. 2001;201:111–5.
- Park YJ, Baek JH, Lee S, Kim C, Rhee H. Whole genome and global gene expression analyses of the model mushroom *Flammulina velutipes* reveal a high capacity for lignocellulose degradation. *PLOS ONE*. 2014;9:546830:e93560. <http://dx.doi.org/10.1371/journal.pone.0093560>.
- Paszczynski A, Crawford RL. Degradation of azo compounds by ligninases from *Phanerochaete chrysosporium* involvement of veratryl alcohol. *Biochem Biophys Res Commun*. 1991;178:1056–63.
- Polónia I [Ph.D. thesis] Olive mill wastes as substrate to produce *Pleurotus* mushrooms. Lisboa, Portugal: Agro-Food Technology, Universidade Técnica de Lisboa, Instituto Superior de Agronomia; 2004.
- Psurtseva N. Modern taxonomy and medicinal value of the *Flammulina* mushrooms. *Int J Med Mushrooms*. 2005;7:449–51.
- Reina R, Liers C, Ocampo JA, García-Romera I, Aranda E. Solid state fermentation of olive mill residues by wood- and dung-dwelling Agaricomycetes: effects on peroxidase production, biomass development and phenol phytotoxicity. *Chemosphere*. 2013;93:1406–12.
- Ruiz-Rodríguez A, Soler-Rivas C, Polonia I, Wichers HJ. Effect of olive mill waste (OMW) supplementation to Oyster mushrooms substrates on the cultivation parameters and fruiting bodies quality. *Int Biodeterior Biodegrad*. 2010;64:638–45.
- Ruiz-Rodríguez A, Polonia I, Soler-Rivas C, Wichers HJ. Ligninolytic enzymes activities of Oyster mushrooms cultivated on OMW (olive mill waste) supplemented media, spawn and substrates. *Int Biodeterior Biodegrad*. 2011;65:285–93.
- Sampedro I, Romero C, Ocampo JA, Brenes M, García-Romera I. Removal of monomeric phenols in dry mill olive residue by saprobic fungi. *J Agric Food Chem*. 2004;52:4487–92.
- Sampedro I, D'Annibale A, Ocampo JA, Stazi SR, García-Romera I. Bioconversion of olive-mill dry residue by *Fusarium lateritium* and subsequent impact on its phytotoxicity. *Chemosphere*. 2005;60:1393–400.
- Sampedro I, Marinari S, D'Annibale A, Ocampo JA, Grego S, García-Romera I. Organic matter evolution and partial detoxification in two-phase olive mill waste colonized by white-rot fungi. *Int Biodeterior Biodegrad*. 2007;60:116–25.
- Sampedro I, D'Annibale A, Federici F, García-Romera I, Siles JA, Petruccioli M. Non-supplemented aqueous extract from dry olive mill residue: a possible medium for fungal manganese peroxidase production. *Biochem Eng J*. 2012;65:96–9.
- Sharma VP, Sharma SR, Kumar S. Effect of various supplements on lignocellulolytic enzyme production and yield of culinary-medicinal mushroom *Flammulina velutipes* (W. Curt.: Fr.) Singer (Agaricomycetidae). *Int J Med Mushrooms*. 2008;10:87–92.
- Sharma VP, Kumar S, Tewari RP. *Flammulina velutipes*, the culinary medicinal winter mushroom. Chambaghat: Directorate of Mushroom Research, Indian Council of Agricultural Research; 2009.
- Singleton VL, Orthofer R, Lamuela-Raventós RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods Enzymol*. 1999;299:152–78.
- Tang XN, Bian GQ, Zhang M, Yang HB, Yu JH. Studies on cultivating *Flammulina velutipes* with *Paspalum notatum*. *Edible Fungi China*. 2001;20:10–2.
- Wood TM, Bhat KM. Methods for measuring cellulase activities. *Methods Enzymol*. 1988;160:87–112.
- Wood DA, Goodenough PW. Fruiting of *Agaricus bisporus*: changes in extracellular enzyme activities during growth and fruiting. *Arch Microbiol*. 1977;114:161–5.
- Zervakis G, Yiatras P, Balis C. Edible mushrooms from olive oil mill wastes. *Int Biodeterior Biodegrad*. 1996;38:237–43.
- Zervakis GI, Koutrotsios G, Katsaris P. Composted versus raw olive mill waste as substrates for the production of medicinal mushrooms: an assessment of selected cultivation and quality parameters. *BioMed Res Int*. 2013. <http://dx.doi.org/10.1155/2013/546830>.
- Zucconi F, Pera A, Forte M, De Bertoldi M. Evaluating toxicity of immature compost. *Biocycle*. 1981;22:54–7.