



ELSEVIER

CIRUGÍA ESPAÑOLA

www.elsevier.es/cirugia



Image of the month

Jejuno-gastric Invagination After Billroth II Gastrectomy^{☆,☆☆}



Invaginación yeyuno-gástrica tras gastrectomía tipo Billroth II

Arturo Cruz Cidoncha,^{a,*} Laura García del Salto,^b Enrique González González,^a
Miguel Ángel García-Ureña^a

^a Servicio de Cirugía General y del Aparato Digestivo, Hospital Universitario del Henares, Coslada, Madrid, Spain

^b Servicio de Radiología, Hospital Universitario del Henares, Coslada, Madrid, Spain



Fig. 1



Fig. 2

A 69-year-old patient with a history of partial gastrectomy due to a gastroduodenal ulcer came to our emergency department with abdominal pain. Physical examination detected a distended abdomen with diffuse pain. Computed tomography identified the invagination of a long section of mesentery and jejunum in the gastric remnant (Figs. 1 and 2) with defined gastric dilatation that reached the pelvis (Fig. 1). During surgery, we found a jejuno-gastric invagination through a large gastrojejunral anastomosis with dilatation of the gastric remnant. We performed disinvagination and fixation of the efferent loop on the right antero-lateral wall. The patient presented and uneventful recovery with no complications.

[☆] Please cite this article as: Cruz Cidoncha A, García del Salto L, González González E, García-Ureña MÁ. Invaginación yeyuno-gástrica tras gastrectomía tipo Billroth II. Cir Esp. 2015;93:194.

^{☆☆} The case report presented in this article has been accepted for presentation in poster format at the 30th Spanish National Congress of Surgery.

* Corresponding author.

E-mail address: acidoncha@hotmail.com (A. Cruz Cidoncha).