



Allergologia et immunopathologia

www.elsevier.es/ai



ORIGINAL ARTICLE

Differential Th1/Th2 balance in peripheral blood lymphocytes from patients suffering from flea bite-induced papular urticaria

A. Cuéllar^a, A. Rodríguez^b, F. Rojas^a, E. Halpert^c, A. Gómez^d, E. García^{c,*}

^aDepartamento de Microbiología. Facultad de Ciencias. Pontificia Universidad Javeriana. Bogotá. Colombia

^bVicerrectoría de Investigaciones. Universidad Militar Nueva Granada. Bogotá. Colombia

^cDepartamentos de Alergia y Dermatología Pediátrica. Fundación Santa Fe de Bogotá. Bogotá. Colombia

^dInstituto de Genética Humana. Facultad de Medicina. Pontificia Universidad Javeriana. Bogotá. Colombia

KEYWORDS

Papular urticaria;
Interferon γ ;
Interleukin-4;
Intracellular cytokine;
Skin allergy;
Ectoparasite

Abstract

Background: The Th1/Th2 balance has not been characterized in patients suffering from flea bite-induced papular urticaria (FBPU). Our aim was to improve understanding of the immunopathogenesis of CD4+ and CD8+ T-cells in humans suffering from flea bite-induced papular urticaria. **Methods:** Peripheral blood mononuclear cells were obtained from 18 pediatric patients and 10 age-matched healthy controls. Cellular phenotypes, intracellular production of interferon gamma (IFN γ) and interleukin-4 (IL-4) in T-cells stimulated with polyclonal stimuli was determined by flow cytometry following short-term *in vitro* stimulation.

Results: The results revealed lower frequencies of IFN γ -secreting ($p = 0.02$) and higher frequencies of IL-4-secreting ($p = 0.03$) CD4+ T-cells in patient lymphocyte cultures compared to healthy control cultures in the presence of polyclonal stimuli. This is the first description of differential cytokine patterns in papular urticaria patients.

Conclusion: Patients suffering from papular urticaria have an atopic status compared to healthy children.

© 2008 SEICAP. Published by Elsevier España, S.L. All rights reserved.

Introduction

Allergic diseases are characterized by a complex immune response that is predominantly related to the secretion of Th2-associated cytokines. These cytokines are responsible for inducing and maintaining allergic inflammation^{1,2}.

Papular urticaria has been characterized as a chronic allergic disease, manifesting itself as hypersensitive reactions

on the skin caused by exposure to ectoparasites, such as fleas³. Previous studies concerning the immunological mechanisms involved in this disease have shown large eosinophil infiltrates in biopsies from patients' cutaneous lesions⁴. Moreover, it has been shown that simultaneous exposure of patients' dendritic cells to flea antigens and lipopolysaccharide induces increased expression of the CD86 co-stimulating molecule that is associated with the induction of a Th2 re-

*Corresponding author.

E-mail: eligarci@cable.net.co (E. García).

sponse⁵. Humoral immunity studies have revealed greater protein recognition in flea extract by IgG1 and IgG3 from healthy individuals and by IgE from patients with papular urticaria (unpublished data).

The presence of eosinophils in skin lesions and differential antigenic recognition by some antibody isotypes led to the hypothesis that patients suffering from flea bite-induced papular urticaria (FBPU) would have a Th2-dominant response, as has been described for other allergic diseases. The objective of the current work was to identify cytokines secreted by CD4⁺ and CD8⁺ lymphocytes (IFN γ and IL-4) as indicators of Th1 and Th2 responses, respectively, after stimulation with a polyclonal stimuli.

Materials and Methods

Study population

The sample included 18 patients, aged one to 13, who had been clinically diagnosed as suffering from FBPU. These patients were attended by the Pediatric Dermatology and Allergy Department of the Fundación Santa Fe de Bogotá, in Bogotá, Colombia. Patients excluded from consideration for participation in the study included those having secondary infected lesions or suffering from immuno-suppression caused by systemic disease, as well as those who had been treated with immuno-suppressive medication; those who had received antihistamines 5 days before the consultation, and/or who had been treated with flea extract. Ten healthy children admitted to the same institution for elective surgery and sharing the same age group and socioeconomic characteristics as the FBPU patients were included as controls. The investigation was approved by the ethics committees at both the Fundación Santa Fe de Bogotá and the Universidad Javeriana.

Disease diagnosis

FBPU was diagnosed according to clinical characteristics. Patients usually had lesions appearing as grouped papules, a symptom commonly associated with pruritus. Papules were often excoriated or crusted, appearing intermittently in a chronic course and leaving hypo- or hyper-pigmented macules behind. These were most often located in areas where clothing fits snugly such as the socks and the waist-band. In some patients, exposed areas of the extremities were also affected.

PBMC isolation and lymphocyte stimulation

Peripheral blood mononuclear cells (PBMC) were prepared from fresh heparinised blood using a Ficoll-hypaque density gradient (Sigma, St. Louis, MO). The cells were adjusted to 2×10^6 cells in a final 2 ml volume of RPMI 1640 medium containing antibiotics, non-essential amino acids, sodium pyruvate, and 10% FCS. PBMC were incubated with 1 μ g/ml of anti-CD28 and anti-CD49d. A positive control of 3.7 μ g/ml of staphylococcal enterotoxin B (SEB) was used⁶. A negative control (no stimulus) was included in every experiment. The cultures were incubated for 9 h at 37 °C in 5% CO₂, followed by an additional 3 h incubation in the presence of 10 μ g/ml

Brefeldin A (Sigma). Surface staining was performed with anti-CD3-FITC, anti-CD4-PerCP or anti-CD8-PerCP, anti-CD69-APC (BD Pharmingen); after fixation and permeabilisation, the cells were then stained with IFN γ -PE or IL-4-PE (BD Pharmingen). Data were acquired and analysed using a FACS-Calibur flow cytometer (BD Immunocytometry Systems) and Cell Quest software.

Quantifying cytokine levels in plasma

Cytokines were quantified in plasma using a Cytometric Bead Array kit (BD Biosciences), following the manufacturer's instructions. Beads displaying different PerCP fluorescence intensities were used for the quantification; they were coated with PE-conjugated capture antibodies that were specific for IFN γ and IL-4. Concentration was calculated using different cytokine patterns at known concentrations. Data were acquired using a FACSCalibur flow cytometer (BD Immunocytometry Systems), and Cell Quest software was used for analysis.

Statistical analysis

SPS statistical software was used for analysing the data. All categorical variables were described using frequencies, while continuous variables were reported as means or medians with their corresponding 95% confidence intervals and standard deviations. The U Mann-Whitney test was used for making comparisons; the significance level was set at < 0.05.

Results

PBMC were obtained and cultured in the presence of polyclonal stimuli to characterize the Th1/Th2 cytokine balance in lymphocytes from healthy controls and paediatric patients suffering from FBPU.

The patients were 50/50% with respect to sex, the age average was 3.5 years (SEM 0.7 years). Twenty-eight percent of the patients in this study reported a personal history of atopy (asthma, allergic rhinitis, or atopic dermatitis) while 78% of them reported a family history of atopy. Seventy percent of healthy children were males and the age average was 3.4 years (SEM 0.6 years); none reported a personal history of atopy, and 11% reported a family history of atopy.

Plasma levels of IFN γ and IL-4 were quantified by flow cytometry to establish differences in plasmatic cytokine levels between the patients and controls. The results (32.7 pg/ml, SEM 3.3 pg/ml in healthy children and 29.5 pg/ml, SEM 2.5 pg/ml in patients for IFN γ and 45.6 pg/ml, SEM 3.6 pg/ml in healthy children and 55.8 pg/ml, SEM 2.7 pg/ml for IL-4) revealed no quantitatively significant differences between patients and healthy controls for any of the cytokines evaluated.

Expression levels of cellular activation markers were examined in CD3⁺/CD4⁺ and CD3⁺/CD8⁺ T-cells to assess the activation status of the T-cells after various culture conditions. When cells were unstimulated or activated with SEB as a polyclonal stimulus, CD69 expression levels were similar in both patients and controls (24.2% in healthy children and 29.3% in patients).




Figure 1. CD3+/CD4+ and CD3+/CD8+ T-cell cytokine secretion after stimulation with SEB, from healthy controls (white) and patients (grey). The results are shown in terms of average values and standard errors.

Polyclonally stimulated CD3+/CD4+/CD69+ or CD3+/CD8+/CD69+ cells were analyzed to identify IFN γ - and IL-4-producing cells in both the patient and the control group. A greater percentage of IL-4-secreting CD4+ T-cells were found in patients (1.3 %) compared with healthy children (0.4 %) ($p = 0.03$), while a greater proportion of IFN γ -secreting CD4+ T-cells were found in healthy controls (3%) compared with patients (1.2 %) ($p = 0.02$). No differences were found in IL-4- or IFN γ -secreting CD8+ T-cells (fig. 1).

Discussion

FBPU in humans has classically been defined as a chronic allergic disease. However, no reports have been published to date indicating whether this pathology is related to a Th2-predominant response. IFN γ and IL-4 secreted by lymphoid subpopulations using polyclonal stimuli have been identified in this work, as well as plasmatic cytokine levels as indicators of Th1 and Th2.

Previous studies have reported greater circulating levels of IL-4 in allergic individuals compared to healthy controls⁷⁻⁹. Plasmatic levels of IL-4 and IFN γ were quantified in patients suffering from FBPU to establish whether the differences found at cellular level were reflected at the systemic level. According to our results, no significant difference was found between the patients and healthy controls with regard to plasmatic IL-4 and IFN γ levels. These results indicate that the disease's clinical manifestations were not associated with alterations at the systemic level.

CD69 expression was evaluated to establish whether allergic inflammation in papular urticaria was associated with functional alterations at the cellular level; CD69 has been used as both an *in vivo* and *in vitro* activation marker¹⁰. Lymphocytes from patients suffering from hypersensitive reactions to drugs have increased CD69 expression as compared to healthy individuals' lymphocytes^{11,12}, indicating that their functional activity is important. The results shown here indicate that both cells from patients suffering from FBPU and those from healthy controls were equally competent because no differences in CD69 activation marker expression were found after polyclonal stimulation.

With regard to the balance between Th2 and Th1 cytokine production in allergic diseases, it has been shown that children at risk for developing atopic dermatitis (followed-up

from birth to 3 years of age) are predominantly predisposed to respond with Th2 cytokine production after SEB stimulation¹³. Quantification of PBMC-secreted cytokines from children suffering from asthma, eczema, and/or atopic rhinitis has revealed increased IL-4 and IL-5 production and reduced IFN γ production after SEB stimulation as compared to cells from healthy children¹⁴.

The evaluation of Th2 and Th1 cytokines in this study through the quantification of IL-4 and IFN γ production (after polyclonal stimulation) demonstrates that cells from patients suffering from papular urticaria predominantly responded with Th2 cytokine production, indicating that patients underwent an atopic stage that may have predisposed them to develop the disease. A tendency towards greater IL-4 secretion was observed in the case of CD8+ T-cells, even though the difference was not significant. Differential cytokine secretion patterns by T-cells may be important contributing factors in the development of childhood papular urticaria.

These results are similar to those reported in the canine model of the disease¹⁵. Flea bite-induced allergic dermatitis is one of the most common allergic diseases in dogs, corresponding to around 50% of canine dermatological disorders. This allergic dermatitis generally coexists with other allergic conditions. Examination of peripheral blood cell-secreted cytokines in the canine model by mRNA quantification revealed greater mRNA expression for IL-4 in allergic dogs and increased mRNA expression for all cytokines quantified (including IFN γ) following exposure to fleabite¹⁵.

These findings support the hypothesis that susceptibility for the development of FBPU is related to a base atopic state and the predomination of Th2 cytokine production (such as IL-4).

This is the first description of differential cytokine patterns in papular urticaria patients.

Reported variations in the regulation of an allergic response to flea antigens during the development of papular urticaria could facilitate the implementation of antigen-specific therapies favoring allergy desensitisation.

Funding/Support

This study was financed by Fundación Santa Fe de Bogotá and the Vicerrectoría Académica, Pontificia Universidad Javeriana, Bogotá, Colombia.

Acknowledgments

The authors wish to thank the collaboration of doctors Manuel Forero, Mariela Tavera, Clara Ordoñez, Ana María Salazar, Armando Rojas, and Adriana Motta for help to get the patients.

References

1. Akdis AC. Allergy and hypersensitivity: mechanisms of allergic disease. *Curr Opin Immunol*. 2006;18:718-26.
2. Romagnani S. Immunologic influences on allergy and the TH1/TH2 balance. *J Allergy Clin Immunol*. 2004;113:395-400.
3. Feingold BF, Benjamini E. Allergy to flea bites. Clinical and experimental observations. *Ann Allergy* 1961;19:1275-89.
4. Garcia E, Halpert E, Rodriguez A, Andrade R, Fiorentino S, Garcia C. Immune and histopathologic examination of flea bite-induced papular urticaria. *Ann Allergy Asthma Immunol*. 2004;92:446-52.
5. Cuellar A, García E, Rodríguez A, Halpert E, Gómez A. Functional dysregulation of dendritic cells in patients with papular urticaria caused by fleabite. *Arch Dermatol*. 2007;143:1415-9.
6. Jaimes MC, Rojas OL, Gonzalez AM, Cajiao I, Charpilienne A, Pothier P, et al. Frequencies of virus-specific CD4(+) and CD8(+) T lymphocytes secreting gamma interferon after acute natural rotavirus infection in children and adults. *J Virol*. 2002;76:4741-9.
7. Qiao HL, Yang J, Zhang YW. Relationships between specific serum IgE, cytokines and polymorphisms in the IL-4, IL-4Ralpha in patients with penicillins allergy. *Allergy*. 2005;60:1053-9.
8. Borres MP, Bjorksten B. Peripheral blood eosinophils and IL-4 in infancy in relation to the appearance of allergic disease during the first 6 years of life. *Pediatr Allergy Immunol*. 2004;15:216-20.
9. Ohshima Y, Katamura K, Miura M, Mikawa H, Mayumi M. Serum levels of interleukin 4 and soluble CD23 in children with allergic disorders. *Eur J Pediatr*. 1995;154:723-8.
10. Maino VC, Suni MA, Ruitenberg JJ. Rapid flow cytometric method for measuring lymphocyte subset activation. *Cytometry*. 1995;20:127-33.
11. Beeler A, Zaccaria L, Kawabata T, Gerber BO, Pichler WJ. CD69 upregulation on T cells as an in vitro marker for delayed-type drug hypersensitivity. *Allergy*. 2008;63:181-8.
12. Golebiowska-Wawrzyniak M, Markiewicz K, Rytwinski K, Bulawa E, Czerwinska L, Wnuk A, Derentowicz P, Kozar A, Siwinska-Golebiowska H. [CD69 and PCNA expression of T-lymphocytes from children with cow's milk allergy (CMA)]. *Med Wieku Rozwoj*. 2003;7:565-75.
13. Sharp MJ, Rowe J, Kusel M, Sly PD, Holt PG. Specific patterns of responsiveness to microbial antigens staphylococcal enterotoxin B and purified protein derivative by cord blood mononuclear cells are predictive of risk for development of atopic dermatitis. *Clin Exp Allergy*. 2003;33:435-41.
14. Smart JM, Kemp A. Increased Th1 and Th2 allergen-induced cytokine responses in children with atopic disease. *Clin Exp Allergy*. 2002;32:796-802.
15. Wuersch K, Brachelente C, Doherr M, Reist M, Sattler M, Forster U, et al. Immune dysregulation in flea allergy dermatitis—a model for the immunopathogenesis of allergic dermatitis. *Vet Immunol Immunopathol*. 2006;110:311-23.