



Allergologia et immunopathologia

www.elsevier.es/ai



ORIGINAL ARTICLE

Prevalence of asthma and other allergic diseases in children born after in vitro fertilisation

F. Cetinkaya^{a,*}, S.A. Gelen^b, E. Kervancioglu^c, E. Oral^d

^aPediatrician and Pediatric Allergist in Sisli Etfal Education and Research Hospital, Istanbul, Turkey

^bPediatrician in Sisli Etfal Education and Research Hospital, Istanbul, Turkey

^cEmbryolog in Kibele ART Center in Istanbul, Turkey

^dGynecolog and Obstetrician in International Hospital, Istanbul, Turkey

KEYWORDS

Allergy;
Asthma;
Atopic dermatitis;
Allergic rhinitis

Abstract

Background: Children born after in vitro fertilisation (IVF) are under greater risk of development of some health problems than those children born after spontaneous conception. Yet it is not exactly known what the prevalence of asthma and other allergic diseases among these children is.

Aims of the study: To investigate the prevalence of asthma, allergic rhinitis and atopic dermatitis in children born after IVF, and controls born after spontaneous pregnancy using questionnaires of International Study of Asthma and Allergies in Children (ISAAC).

Methods: We recruited 158 children (mean age: 4.60 ± 2.14 years) born after IVF pregnancies and 102 children (mean age: 5.27 ± 2.8 years) as control group to investigate the prevalences of asthma, atopic dermatitis and allergic rhinitis. The questions in the questionnaire were asked to the parents by telephone or face to face.

Results: Wheezing ever has been reported in 19% of the children born after IVF and in 17.6% of the control group ($p > 0.05$). No significant differences were found between groups, in terms of use of any anti-asthma drugs, physician diagnosed asthma and admission to emergency room with wheezing ($p > 0.05$). In addition, prevalence of allergic rhinitis and atopic dermatitis were also comparable between two groups ($p > 0.05$).

Conclusions: Prevalences of asthma, allergic rhinitis and atopic dermatitis are similar in children born after IVF and children born after spontaneous conception.

© 2008 SEICAP. Published by Elsevier España, S.L. All rights reserved.

Introduction

The prevalence of childhood asthma and other allergic diseases have increased tremendously, especially in recent years, despite the absence of a satisfactory explanation for

this increase^{1,2}. Up to now, many factors in the antenatal and neonatal periods have been suggested as playing a role in the development of allergic diseases in children^{3,4}.

Since the first child conceived after in-vitro fertilisation (IVF) was born in the UK in 1978, more than one million ba-

*Corresponding author.

E-mail: feyzulkaya@myinet.com (F. Cetinkaya).

Table I. Prevalences of allergy and asthma symptoms in children born after IVF and controls

| Questions | IVF group | | Controls | | P |
|---|-----------|------|----------|------|-------|
| | n | % | n | % | |
| Wheeze ever | 30 | 19 | 18 | 17.6 | 0.786 |
| Wheeze in last 12 months | 21 | 13.3 | 9 | 8.8 | 0.271 |
| Exercise induced wheeze in last 12 months | 14 | 8.9 | 8 | 7.8 | 0.773 |
| Physician diagnosed asthma | 25 | 15.8 | 17 | 16.7 | 0.857 |
| Use of asthma drugs | 22 | 13.9 | 12 | 11.8 | 0.614 |
| Emergency room visit for acute wheezing | 14 | 8.9 | 8 | 7.8 | 0.773 |
| Runny nose or sneezing without a cold | 33 | 20.9 | 21 | 20.6 | 0.954 |
| Itchy eyes and nose in last 12 months | 26 | 16.5 | 13 | 12.7 | 0.413 |
| Physician diagnosed hay fever | 23 | 14.6 | 13 | 12.7 | 0.378 |
| Itchy rash in specific locations lasting at least 6 months in the previous year | 14 | 8.9 | 10 | 9.8 | 0.970 |

bies worldwide have been born with this method. Many papers have been published on the health problems of children born after IVF comparing them with children born after spontaneously conceived pregnancies^{5,6}.

The prevalence of allergic diseases and asthma have been studied extensively in normal populations and in some risk groups, yet it is not exactly known if there is an increased risk for allergic diseases and asthma among children born after IVF.

In this study we plan to document the prevalence of asthma, allergic rhinitis and atopic dermatitis among children born after IVF and compared with controls.

Materials and Methods

We recruited 158 children (mean age: 4.60 ± 2.14 years) born after IVF pregnancies in an international hospital between 2002-2004 and 102 children (mean age: 5.27 ± 2.8 years) born after spontaneous pregnancies as control group to investigate the prevalences of asthma, atopic dermatitis and allergic rhinitis. All of the parents of children born in the in-vitro fertilisation centre of international hospital were called and invited to join the study and the controls were randomly selected among the children of a day care centre in the hospital from a similar socioeconomic level. The questions in a Turkish version of the International Study of Asthma and Allergies in Children (ISAAC) questionnaire were asked to the parents by telephone or face to face⁷. The children from both groups with known causes of wheezing such as pneumonia, laryngomalacia/tracheomalacia, congenital heart diseases, cystic fibrosis and gastro-oesophageal reflux were excluded from the study. The study protocol was approved by the institutional ethics committee and verbal consent was obtained from the parents of the children after the aim of the study was described in detail.

Statistical analysis

Descriptive statistics were provided for the numeric and categorical variables using means, standard deviations and percent distributions where necessary. Student t test was used to compare quantitative data and Chi-square test and

Fisher's Chi-square test to compare qualitative data. A P value of less than 0.05 was used as the cut-off for statistical significance.

Results

Although more than 250 parents were invited, only parents of 158 children accepted to join the study. The total number of children in the day care centre was about 110 and we did not want to join more controls from other day care centres to maintain the socioeconomic similarity between the study and control groups. Therefore the study group consisted of 40 twins (80 children) and 78 single children; and the control group of 102 children. The female/male ratio was 0.82 among study group and 1.04 among controls ($p > 0.05$). The mean number of people in the home was also similar (4.3 people in study groups and 4.1 people in controls) ($p > 0.05$). All of the study and control groups were born in Istanbul. Wheezing ever has been reported in 19% of the children born after IVF and in 17.6% of the control group ($p > 0.05$) (table 1). No significant differences were found between groups, in terms of parental allergic diseases, use of any anti-asthma drugs, physician diagnosed asthma and admission to emergency room with wheezing ($p > 0.05$). In addition, prevalences of allergic rhinitis and atopic dermatitis were also comparable between two groups ($p > 0.05$) (table I).

Discussion

Although the reasons for the dramatic increase in allergic diseases have not been clarified yet, some important explanations and hypotheses have been proposed to date⁸. It is generally accepted that a genetic background and exposure to some environmental factors account for the rise in the prevalence of these diseases⁹.

Thousands of children are now born worldwide every year after IVF, but many studies have suggested that children born after IVF have an increased risk of developing certain health problems such as some malignancies¹⁰ and neurological problems¹¹. On the other hand the long-term effects of IVF on the development of allergies and asthma in children

born after IVF are not exactly known, although some increased risk has been reported for asthma¹². When planning this study we wondered if the hormones and drugs used in IVF caused epigenetic modification in DNA and effect gene expression and thus activated or inhibited the genes related to asthma and other allergic diseases. In our study prevalences of wheeze ever, wheezing in last 12 months, symptoms of allergic rhinitis and atopic dermatitis were comparable with former studies from our society^{13,14}. However, prevalences of physician diagnosed asthma and allergic rhinitis were found to be a little higher than in former studies. This may be due to the fact that both the study and control group come from the higher socioeconomic income groups of the population and they visit the hospitals and pediatricians more often than the middle or lower socioeconomic groups.

In conclusion: prevalences of asthma, allergic rhinitis and atopic dermatitis are similar in children born after IVF and children born after spontaneous pregnancies.

References

1. Aberg N, Hesselmar B, Aberg B, Eriksson B. Increase of asthma, allergic rhinitis and eczema in Swedish schoolchildren between 1979 and 1991. *Clin Exp All.* 1995;25:815-9.
2. Sly RM. Changing prevalence of allergic rhinitis and asthma. *Ann Allergy Asthma Immunol.* 1999;82:233-52.
3. Warner JA, Jones CA, Jones AC, Warner JO. Prenatal origins of allergic disease. *J Allergy Clin Immunol.* 2000;105:S493-8.
4. Klinnert MD, Nelson HS, Price MR, Adinoff AD, Leung DY, Mrazek DA. Onset and persistence of childhood asthma: predictors from infancy. *Pediatrics.* 2001;108:E69.
5. Klemetti R, Sevon T, Gissler M, Hemminki E. Health of children born as a result of in vitro fertilization. *Pediatrics.* 2006;118:1819-27.
6. Schieve LA, Rasmussen SA, Buck GM, Schendel DE, Reynolds MA, Wright VC. Are children born after assisted reproductive technology at increased risk for adverse health outcomes? *Obstet Gynecol.* 2004;103:1154-63.
7. Cetinkaya F, Uslu HS, Nuhoglu A. Effect of neonatal sepsis on the development of allergies and asthma in later childhood. *Int Arch Allergy Immunol.* 2007;142:145-50.
8. Strachan DP. Hay fever, hygiene, and household size. *BMJ.* 1989;299:1259-60.
9. Liu AH, Liu AH, Spahn JD, Leung DYM. Childhood asthma. In: Behrman RE, Kliegman RM, Jenson HB, editors. *Nelson Textbook of Pediatrics.* 17th ed. Philadelphia: Saunders; 2004. p. 760-74.
10. Kallen B, Finnstrom O, Nygren KG, Olausson PO. In vitro fertilization in Sweden: child morbidity including cancer risk. *Fertil Steril.* 2005;84:605-10.
11. Strömberg B, Dahlquist G, Ericson A, Finnström O, Köster M, Stjernqvist K. Neurological sequelae in children born after in-vitro fertilisation a population-based study. *Lancet.* 2002;359:461-5.
12. Ericson A, Nygren KG, Olausson PO, Källén B. Hospital care utilization of infants born after IVF. *Hum Reprod.* 2002;17:929-32.
13. Saraçlar Y, Sekerel BE, Kalayci O, Cetinkaya F, Adalıoğlu G, Tuncer A, Tezcan S. Prevalence of asthma symptoms in school children in Ankara, Turkey. *Respir Med.* 1998;92:203-7.
14. Kucukoduk S, Aydin M, Cetinkaya F, Dinc H, Gurses N, Saraçlar Y. The prevalence of asthma and other allergic diseases in a province of Turkey. *Turk J Pediatr.* 1996;38:149-53.