

# Primary hepatic perivascular epithelioid tumor (PEComa)


David F. Schaeffer, Michael P. Poulin

Division of Anatomic Pathology, Vancouver General Hospital, Vancouver, BC Canada.

A 49 year-old woman was found to have a 2.9 cm hepatic lesion in segment 8, initially identified on an abdominal ultrasound performed for chronic abdominal pain and new onset diabetes mellitus, type 2. On computed tomography study the lesion was arterial enhancing with washout on portal venous and delayed phase and was classified as a LI-RADS category 4 (LR-4), suggestive but not diagnostic of hepatocellular carcinoma (HCC); the remainder of the exam was normal with no evidence of metastatic deposits. Of note, the background liver did not show any features of advanced fibrosis, but the patient had a past medical history of non-alcoholic fatty liver disease (NAFLD). The patient subsequently underwent a partial hepatectomy and on gross examination a well circumscribed grey tan lesion was identified. Histologically, the lesion showed a nested architecture, composed of epithelioid cells with round to oval nuclei, variable nucleoli and prominent granular eosinophilic cytoplasm (Figures 1A and 1B; \* denotes background normal hepatic parenchyma). In areas the tumoral cells condensed around blood vessels (Figure 1B). The lesion also showed prominent fibrous trabeculae and pericellular fibrosis, raising the morphological differential diagnosis of a scirrhous or fibrolamellar subtype of hepatocellular carcinoma.

By immunohistochemistry the tumour cells were positive with antibodies to gp100 protein (HMB-45; Figure 1C), melan-A but negative for hepatocyte specific antigen (HSA) and S-100 protein.

The perivascular epithelioid group of tumors (PEComas) is characterized by their co-expression of melanocytic and muscle markers and includes angiomyolipoma, lymphangioliomyoma, and clear cell ('sugar') tumors of the lung, pancreas, and uterus.<sup>1-4</sup> Primary hepatic PEComas are rare, but usually present in non-cirrhotic younger patients and on imaging studies often are suggestive of hepatocellular carcinoma. Although renal angiomyolipoma and pulmonary lymphangioliomyomatosis are often



**Figure 1.** Representative microscopic photomicrographs of a primary hepatic perivascular epithelioid tumor (PEComa). **A.** The lesion is well circumscribed and separated by a fibrous capsule from the normal background hepatic parenchyma (\*). Also note the broad fibrous bands and prominent vasculature within the lesion itself. **B.** High-power image shows epithelioid tumor cells characterized by variable nucleoli and granular eosinophilic cytoplasm that condense around a vessel. **C.** The tumor cells strongly express the melanocytic marker HMB-45 (Magnifications: A 2x, B and C 200x).

seen in the setting of tuberous sclerosis complex, there has been no defined hereditary predisposition for hepatic PEComas to date. Fortunately, the majority of hepatic PEComas

reported in the literature behaved benign after complete resection with only one pulmonary metastasis reported.<sup>4</sup>

#### REFERENCES

1. Tan Y, Xiao E-H. Hepatic perivascular epithelioid cell tumor (PEComa): dynamic CT, MRI, ultrasonography, and pathologic features-analysis of 7 cases and review of the literature. *Abdominal Imaging* 2012; 37: 781-7.
2. Paiva CE, Neto FAM, Agaimy A, Domingues MAC, Rogatto SR. Perivascular epithelioid cell tumor of the liver coexisting with a gastrointestinal stromal tumor. *World J Gastroenterol* 2008; 14: 800-2.
3. Akitake R, Kimura H, Sekoguchi S, Nakamura H, Seno H, Chiba T, Fujimoto S. Perivascular epithelioid cell tumor (PEComa) of the liver diagnosed by contrast-enhanced ultrasonography. *Int Med* 2009; 48: 2083-6.
4. Folpe AL, Goodman ZD, Ishak KG, Paulino AF, Taboada EM, Meehan SA, Weiss SW, et al. Clear cell myomelanocytic tumor of the falciform ligament/ligamentum teres: a novel member of the perivascular epithelioid clear cell family of tumors with a predilection for children and young adults. *Am J Surg Pathol* 2000; 24: 1239-46.

#### Correspondence and reprint request:

David F. Schaeffer, MD, FRCPC  
Department of Pathology and Laboratory Medicine.  
The University of British Columbia.  
910 West 10th Avenue. Vancouver, BC V5Z 1M9.  
Tel.: 604-875-4480  
E-mail: david.schaeffer@vch.ca