

## Amoebic liver abscess production by *Entamoeba dispar*

Silvio S. Dolabella,\* Jesús Serrano-Luna,<sup>†</sup> Fernando Navarro-García,<sup>†</sup> René Cerritos,<sup>‡</sup> Cecilia Ximénez,<sup>‡</sup> José Manuel Galván-Moroyoqui,<sup>§,||</sup> Edward F. Silva,<sup>¶</sup> Víctor Tsutsumi,<sup>§</sup> Mineko Shibayama<sup>§</sup>

\* Department of Morphology, Laboratory of Parasitology, Federal University of Sergipe, Aracaju, Sergipe, 49100-000, Brazil.

<sup>†</sup> Department of Cell Biology, CINVESTAV-IPN, Mexico City, Mexico.

<sup>‡</sup> Department of Experimental Medicine, Faculty of Medicine, National Autonomous University of Mexico, Mexico City, Mexico.

<sup>§</sup> Department of Infectomics and Molecular Pathogenesis, CINVESTAV-IPN, Mexico City, Mexico.

<sup>||</sup> Department of Medicine and Health Sciences, University of Sonora, Hermosillo, Sonora, Mexico.

<sup>¶</sup> Department of Parasitology, Institute for Biological Sciences, Federal University of Minas Gerais, Minas Gerais, Belo Horizonte, 31270-901, Brazil.

### ABSTRACT

Although *Entamoeba dispar* displays a similar morphology to *Entamoeba histolytica*, cellular and molecular studies have revealed significant differences between these two amoebae, including the former being characterized as non-pathogenic and the later as pathogenic. However, recent *in vivo* and *in vitro* experiments have shown that *E. dispar* strains of different origin are capable of causing liver damage and destroying cell culture lines in the presence of common intestinal bacteria. These results suggested that *E. dispar* may present pathogenic behavior according to the specific *E. dispar* strain, culture and environmental conditions. To investigate this possibility, we carried out *in vivo* and *in vitro* studies using a xenic strain *E. dispar* (ICB-ADO) isolated from a symptomatic non-dysenteric Brazilian patient. This strain was able to induce liver necrosis in a hamster model that was more severe than that produced by *E. histolytica*. The ICB-ADO isolate also caused significantly more destruction of cultured MDCK cells and increased loss of transepithelial resistance than did the *E. histolytica*. Xenic *E. dispar* exhibited high proteolytic activity, which was partially inhibited by the addition of cysteine-protease inhibitors. Based on our biochemical and molecular characterization of *E. dispar* (ICB-ADO) xenic culture and its ability to produce liver abscesses, we conclude that this specific strain can indeed produce tissue damage, distinct from the frequently used non- pathogenic *E. dispar* SAW 760 strain.

**Key words.** Amoebiasis. *Entamoeba dispar*. *Entamoeba histolytica*. Xenic culture. Liver damage. Protease activity.

### INTRODUCTION

*Entamoeba histolytica*, the causative agent of human amoebiasis, remains as an important cause of morbidity and mortality in developing countries and is the responsible for up to 100,000 deaths worldwide each year. The *Entamoeba dispar* species is morphologically similar to *E. histolytica*, but has been considered a distinct strain based on differences in the isoenzymatic patterns observed between amoebae isolates from asymptomatic patients and

those with invasive disease.<sup>1</sup> This characterization has been reinforced by several studies that have demonstrated antigenic and genetic differences between the two strains.<sup>2-4</sup> The presence of *E. dispar* in the mammalian large intestine is widely believed to constitute a parasite colonization behaving as a commensal; however, this species has been isolated from patients with symptomatic non-dysenteric colitis.<sup>5</sup> More recently, *E. dispar* sequences of DNA have been detected and genotyped in samples obtained from patients with amoebic liver abscesses, suggesting that *E. dispar* can be also involved in the damage of human large intestine and liver.<sup>6</sup>

An important molecule that is known to be related to the pathogenicity of *E. histolytica*, the lectin Gal/GalNAc, has also been identified in *E. dispar*. However, some conformational differences exist on the *E. dispar* form of Gal/GalNAc lectin that may explain the reduced adherence and cytotoxicity of this strain that have been observed *in vitro*, and the

Correspondence and reprint request: Mineko Shibayama  
Department of Infectomics and Molecular Pathogenesis  
Center for Research and Advanced Studies  
National Polytechnic Institute. Av. IPN, No. 2508. Mexico City 07360, Mexico  
Tel.: (52) 55 5747-3348. Fax: (52) 55 5747-3398  
E-mail: mineko@cinvestav.mx

Manuscript received: June 02, 2011.  
Manuscript accepted: August 28, 2011.

significantly lower virulence of its trophozoites in animal models.<sup>7-9</sup> Interestingly, both *Entamoeba* species have almost the same set of genes that encode for the major virulence factors. Cysteine proteases (CP) are important virulence factors in the *E. histolytica* species; the production and release of these enzymes have been found to be about ten to 1,000 times higher in *E. histolytica* than in the *E. dispar* species.<sup>10,11</sup> The activity of small pore-forming amphipathic peptides, known as amoebapores, in *E. dispar* is about one-third of that observed in *E. histolytica*, despite the fact that about 95% amino acid homology exists between the two strain's amoebapores.<sup>12</sup>

Previous studies have suggested that molecular changes may be induced in *E. histolytica* trophozoites during the process of xenic culturing, resulting in an increased resistance to lysis by complement factors and/or modification of surface antigens of the amoeba. In this manner, the parasite may be able to effectively overcome the immune system's ability to recognize, attack, and ultimately eliminate the parasite.<sup>13-15</sup>

While it is generally accepted that the *E. histolytica* and *E. dispar* protozoans represent two distinct species, the pathogenic potential of the latter remains controversial. Recent studies have demonstrated that *E. dispar* trophozoites can cause intestinal focal lesions in experimental animal models and destruction of cultured monolayer epithelial cells; these findings suggest that *E. dispar* may induce at least some pathological changes in the human colon.<sup>16</sup> Other recent reports have indicated that *E. dispar* can also produce extensive hepatic lesions in hamsters when trophozoites are administered in conjunction with the host's original intestinal flora; the fact that these lesions are similar to those caused by *E. histolytica*, suggests that normal bacterial flora may interact with some *E. dispar* strains and influence their pathogenic behavior.<sup>5,17,18</sup> To date, very few experimental studies related to the use of xenic strains of *E. dispar* have been reported, and even fewer have focused on the role of bacteria play on influencing the pathogenic behavior of *E. dispar* strains.<sup>17,18</sup>

In the present study, we tested the *in vitro* and *in vivo* effect of a specific *E. dispar* strain (ICB-ADO) that in previous preliminary studies displayed a virulence phenotype of its trophozoites in xenic culture conditions similar to the *E. histolytica* strain HM1-IMSS.<sup>19</sup> Furthermore, this strain had been recently genotyped using non-coding intergenic DNA regions corresponding to tRNA genes. The four phylogenetic reconstructions performed had clearly

demonstrated that the *E. dispar* ICB-ADO strain is a genetic variant of *E. dispar* species.<sup>6</sup> We describe here our findings from the investigation into whether *E. dispar* presents pathogenic behavior according to the specific strains, culture and host conditions.

## MATERIAL AND METHODS

### Cell culture

*E. dispar* trophozoites from the ICB-ADO strain (zymodeme I-non-pathogenic)<sup>5</sup> were cultured under xenic conditions in cell culture bottles (30 mL; Corning, Lowell, MA, USA) using Pavlova's medium formula as modified by Costa, *et al.*<sup>17</sup> and containing bacterial flora. After a period of 72 h, the supernatant was removed in order to eliminate the bacterial flora and substituted with sterile PBS, pH 7.2; the mixture was incubated at 37 °C with gently shaking to wash the adhered trophozoites. This procedure was repeated once, after which the culture bottles were transferred to an ice bath for a total of 10 min. The prepared trophozoites were centrifuged at 200 x g for 5 min. The sediment amoebae were diluted in sterile PBS, pH 7.2, to obtain desired concentrations. The monoxenic strain of *E. dispar* trophozoites ICB-ADO (with *Crithidia fasciculata*) (non-pathogenic zymodeme I) and the axenic strain of *E. histolytica* trophozoites HM1-IMSS (pathogenic zymodeme XIX) were cultured in TYI-S-33 medium using borosilicate glass culture tubes.<sup>20</sup> Amoebae were harvested after 72 h of incubation by chilling the culture tubes in an ice bath for 10 min. After centrifugation at 200 x g for 5 min, the cell pellets were resuspended in fresh medium. Genetic identification of the strain was confirmed by repeating the analysis of the isoenzymatic pattern and by employing a PCR method for determining the conformational polymorphism of the fragment of 482 pb as described earlier.<sup>21</sup>

### Genotyping of ADO *Entamoeba* isolate

DNA was extracted from trophozoites (10<sup>6</sup>) harvested from a xenic culture using the TRIZOL reagent kit (Invitrogen Life Technologies, Carlsbad, CA, USA) and following the manufacturer's instructions.

In the present study the *E. dispar* ICB-ADO strain used had been previously re-characterized by PCR amplification of a DNA segment of the small rRNA gene subunit, consisting of 1,950 base pairs (bp) (the

RD amplicon) in both *E. histolytica* and *E. dispar*.<sup>22</sup> For amplification with RD primers, 2  $\mu$ L of DNA was used as the template and 1  $\mu$ L of obtained PCR product (diluted 1:10) was used as the template for amplifications with the primers Psp (specific for *E. histolytica*) and NPsp (specific for *E. dispar*).<sup>22</sup>

Both primer sets are expected to generate an 876-bp fragment that is part of the RD amplicon. The PCR reaction mix included 10 mM Tris-HCl, pH 8.3, 50 mM KCl, 0.001% gelatin, 2 mM MgCl<sub>2</sub>, 0.2 mM of each nucleotide, 0.025 U (Ampli Taq Gold, Applied Biosystems, Foster City, CA, USA) and 1  $\mu$ M of the respective primer. After an initial 10 min step at 95 °C, 35 cycles consisting of 94 °C for 1 min, 55 °C for 1.5 min and 72 °C for 2 min were carried out, followed by a final step of 72 °C for 8 min.

All PCR products were electrophoresed through 1.5% agarose gels (Gibco BRL Products); the gels were then stained with ethidium bromide 0.1% solution for band visualization and photographic images were recorded.

#### Intrahepatic inoculation of trophozoites in hamsters

Male adult golden hamsters (*Mesocricetus auratus*) two months-old and weighing approximately of 100 g were obtained from the Animal Center of the Center for Research and Advanced Studies of the National Polytechnic Institute, Mexico. The animals were anesthetized with sodium pentobarbital (40 mg/kg of body weight; Hypnol®, Lima, Peru) and inoculated intrahepatically with 5 x 10<sup>5</sup> trophozoites of *E. dispar* ADO strain (xenic culture) that had been obtained during log growth phase (72 h). All animal management protocols had been previously approved by the institutional committee. Our institution fulfills all the technical specifications for the production, care and use of laboratory animals and is certified by national law (NOM-062-Z00-1999).

Three animals were euthanized at 1, 3, 6, 12, 24, and 48 h and at the 7th day after inoculation. For controls, three individuals were inoculated with only bacteria that had been used in the xenic cultures; these animals were sacrificed at 6 and 24 h and 7 days post-inoculation. All hamsters were euthanized by an overdose of sodium pentobarbital and were handled according to the guidelines of the 2000 AVMA Panel of Euthanasia.

Liver tissue fragments were collected and fixed in 10% buffered formaldehyde, and processed for paraffin embedding. Sections were stained with hematoxylin-eosin (H&E).

#### Analysis of cytopathic effect

MDCK monolayer cells were cultured to 80% confluence in 8-well culture chambers (Nunc Inc., Rochester, NY, USA). A minimum essential medium (MEM; Gibco-Invitrogen) supplemented with 20% fetal bovine serum (FBS; Gibco-Invitrogen) was used and the cultures incubated at 37 °C in a 5% CO<sub>2</sub> atmosphere. Once the monolayer was formed, 3 x 10<sup>5</sup> trophozoites of *E. dispar* ICB-ADO strain were added and incubated for 5, 15, 30, and 60 min at 37 °C. After each time-point, cultures were washed three times with MEM without FBS, fixed with 10% methanol, stained with 5% Giemsa dye for 40 min and mounted with resin. For controls, the bacteria associated with the xenic culture of the ICB-ADO strain were used.

Microscopic images were digitalized by using a JVC TK-1270/JGB camera attached to a Nikon H500L light microscope. Morphometric analysis was performed using Kontron KS300 v.2.0 software (Kontron Electronics, Poway, CA, USA).

#### Transepithelial electric resistance (TER) measurement

Measurements of TER were evaluated using an EVOM epithelial voltmeter (World Precision Instruments Inc., Sarasota, FL, USA).

MDCK cells were grown in Transwell® chambers (Corning) using MEM medium supplemented with 20% FBS at 37 °C in a 5% CO<sub>2</sub> atmosphere. The chambers were placed in a multiwell plate to form two independent compartments. After reaching confluence (approximately 100,000 cells per well), cells were washed twice with MEM without FBS and trophozoites of *E. histolytica* or *E. dispar* (at a 1:2 trophozoite:MDCK ratio) were added to the Transwell®. TER was measured at 5, 15, 30, and 60 min, and calculations were carried out based on Ohm's law. The normalized values reported here represent the mean  $\pm$  SEM at each time point (n = 6).

#### Protease activities in Azocoll assays

Total crude extracts were obtained by lysing trophozoites with 5 cycles of freeze-thawing. Total protein concentration was quantified by Bradford's method.<sup>23</sup>

Two milligrams of Azocoll (Sigma-Aldrich, St. Louis, MO, USA) were aliquoted into Eppendorf tubes (1.5 mL) containing 100  $\mu$ g of crude extract; the

final volume was adjusted to 500  $\mu$ L with activation buffer (100 mM Tris-OH and 10 mM of  $\text{CaCl}_2$ , pH 7.0). The samples were then incubated at 36 °C for 16 h. The reaction was terminated by addition of 500  $\mu$ L of 10% trichloroacetic acid. Tubes were centrifuged at 4,500 x g for 5 min and supernatants were collected for spectrophotometric absorbance reading at 540 nm.

To determine the effect of inhibitors of cysteine proteases, serine proteases and metalloproteases, the following chemicals were added directly to the extracts and the mixtures allowed to incubate at room temperature for 1 h prior to addition of the substrates: p-hydroxy-mercuribenzoic acid (PHMB; 10 mM), phenylmethylsulphonyl fluoride (PMSF; 5 mM) or ethylenediaminetetracetic acid (EDTA; 2 mM), respectively. The experiments were performed in triplicate for each sample and statistical analysis was carried out using the Bonferroni test to identify significant differences among protease activities.

## RESULTS

### Molecular characterization of the ADO isolate

This isolate had been obtained from the feces of a Brazilian symptomatic patient clinically diagnosed with non-dysenteric colitis. Cysts from the same patient were cultured in Pavlova's medium modified by Costa, *et al.*, as previously described,<sup>17</sup> and cultures were maintained in xenic conditions. The major bacterial components of this culture were determined to be *Escherichia coli*, *Acinetobacter lwoffii* and a rough *E. coli* colony that were presumed as components of the patient's original intestinal flora. Previously, this isolate was genetically identified by Tannich, *et*

*al.*<sup>3</sup> as an *E. dispar* species, based on the conformational polymorphism of the 482 bp DNA fragment of the m17 gene that was amplified with the primers P1-S17 (forward) and P1AS20 (reverse). In our study we sought to re-test the *Entamoeba* species of the ICB-ADO isolate by targeting the gene for the small subunit of rRNA, as described in table 1. Figure 1A shows the PCR products that were obtained with the RD primer which was expected to amplify a 1950 bp segment of DNA in both *E. histolytica* and *E. dispar* species.<sup>22</sup> Our amplification products agreed with the previous genotyping which identified the ICB-ADO strain as an *E. dispar* genetic variant.<sup>6</sup>

### Hepatic lesions analysis


*E. dispar* xenic strain ICB-ADO was found to cause extensive lesions similar to those reported by others for the strain *E. histolytica* HM1-IMSS at seven days post-inoculation.<sup>17,19</sup> However, during the initial periods of infection the speed of abscess development observed was slower. Histological analysis indicated that liver lesions produced by ICB-ADO trophozoites at 3 h showed an important number of parasites in close contact with hepatocytes. Moreover, many of these parenchymal cells presented visual signs of cell damage, as shown in figure 2A. At 6 h post-inoculation, the amoebae induced a discrete inflammatory reaction, and the liver parenchyma proximal to the Glisson capsule appeared damaged (Figure 2B). At 12 h post-infection, the necrotic areas increased in size and numbers of inflammatory foci surrounding the amoebae were also more severe (Figure 2C). At 48 h, extensive areas of liver necrosis with scarce number of trophozoites were seen (Figure 2D).

Table 1. Primers used for genotyping.

Primer name	Gene	Sequence (5' to 3')	Size range (bp)	Conditions
RD5 RD3	SS rRNA <i>E.h./E.d.</i>	GGAAGCTTATCTGGTTGATCCTGCCAGTA GGGATCCTGATCCTTCCGCAGGTTACCTAC	1950	94 °C 1.0' 55 °C 1.5' 72 °C 2.0'
Psp5 Psp3	SS rRNA <i>E.h.</i>	GGCCAATTCAATGAATTGAG GGGATCCTGATCCTTCCGCAGGTTACCTAC	876	94 °C 1.0' 55 °C 1.5' 72 °C 2.0'
NPsp5 NPsp3	SS rRNA <i>E.d.</i>	GGCCAATTTATGTAAGTAAATTGAG CTTGGATTTAGAAACAATGTTCTTC	876	94 °C 1.0' 55 °C 1.5' 72 °C 2.0'


When trophozoites of the *E. histolytica* HM1:IMSS strain was intrahepatically inoculated, several trophozoites were found to be surrounded by inflammation at 12 h; although, the tissue damage was

less extensive than with the ICB-ADO strain (Figure 2E). A granulomatous reaction was more clearly observed with the *E. histolytica* liver lesions than with the *E. dispar* ICB-ADO strain at 48 h post-in-



**Figure 1.** PCR amplification of the xenic strain ICB-ADO and reference strains of species *E. dispar* SAW760 and *E. histolytica* HM1:IMSS. Primer

RD was used to amplify DNA of species from the genus *Entamoeba* (A). Primer NPsp was specific for *E. dispar* species (B) and Psp primer was specific for *E. histolytica* species (C). Line 1: DNA from *E. histolytica* HM1:IMSS strain; line 2: DNA from *E. dispar* SAW760 strain; line 3: DNA from xenic ICB-ADO strain; line 4: negative controls without DNA; line L: DNA size markers.



**Figure 2.** Amoebic liver abscesses produced by *E. dispar* ICB-ADO (A-D) and *E. histolytica* HM1-IMSS (E-F). A. Three hours post-inoculation of *E. dispar* trophozoites. Numerous amoebae were (arrows) in close contact with hepatocytes and scarce inflammatory reaction (ir) was observed. Magnification: 60 x. B. Six hours post-*E. dispar* inoculation. Trophozoites (arrow) are surrounded by a discrete inflammatory infiltrate; associated hepatocytes show signs of damage. Magnification: 60 x. C. At 12 h post-inoculation of *E. dispar* trophozoites, liver parenchyma showed extensive necrotic areas and important number of inflammatory cells (PMN) associated with amoebae (arrows). Magnification: 20 x. D. Forty-eight hours post-*E. dispar* inoculation. The areas of necrosis (n) increased significantly with chronic inflammatory infiltrates (arrows) bordering the damaged parenchyma. Magnification: 10 x. E. Hepatic lesions were produced by *E. histolytica* trophozoites at 12 h post-inoculation. Several trophozoites (arrows) were surrounded by polymorphonuclear leukocytes (PMN) with some hepatocytes showing signs of damage. Magnification: 40 x. F. At 48 h post-inoculation with *E. histolytica* typical granulomatous reaction (g) was observed and associated with significant necrotic areas (n). Magnification: 10 x. All tissues were stained with H&E.

fections (Figure 2F). The strain ICB-ADO that had been monoxenically cultured was unable to cause lesions in the hamster liver (data not shown).

### Cytopathic effect of *E. histolytica* and *E. dispar* strains on MDCK cells

*E. dispar* monoxenic strain ICB-ADO produced a smaller extent of damage to the MDCK monolayer cells (< 50%) as compared with that of *E. histolytica* strain HM1: IMSS (Table 2). *E. dispar* xenic strain ICB-ADO produced more cell damage than its corresponding monoxenic strain. However, this damage was less extensive than that produced by *E. histolytica* strain HM1-IMSS.

### Electrophysiological measurements

The decrease in TER observed when using xenic *E. dispar* ICB-ADO strain trophozoites was slower than with the *E. histolytica* strain. At 30 min, measurements of TER on *E. dispar* monoxenic and xenic

ICB-ADO strains were larger than that observed for *E. histolytica* strain HM1:IMSS. However, after 60 min of interaction all the tested strains presented a total loss of TER (Figure 3).


### Proteolytic activity using Azocoll substrate

The crude extract of monoxenic *E. dispar* ICB-ADO strain showed lower proteolytic activity than the *E. histolytica* strain HM1: IMSS. However, the crude extract from the xenic ICB-ADO culture exhibited stronger proteolytic activity than the *E. histolytica* strain (Figure 4).


The nature of the proteolytic activity observed in the crude extracts was determined by using specific inhibitors for cysteine, serine and metalloproteases. The substrate used for all the inhibitors was the nonspecific protease azo-dye-impregnated collagen (Azocoll). The inhibitor for cysteine protease PHMB had a significant inhibition capacity over the proteolytic activity of the xenic strain (Figure 5). The

**Table 2.** Cytopathic effect of *E. histolytica* (*E.h*) and *E. dispar* (*E.d*) strains on MDCK cell monolayers.


Strain	Percentage of monolayer destruction		
	15 min	30 min	60 min
HM1-IMSS ( <i>E.h.</i> )	27	55	89
ICB-ADO monoxenic ( <i>E.d.</i> )	13	18	37
ICB-ADO xenic ( <i>E.d.</i> )	17	31	60



**Figure 3.** Effect of *E. dispar* (monoxenic and xenic ICB-ADO strains) and *E. histolytica* (axenic HM1: IMSS strain) trophozoites on the transepithelial electrical resistance of MDCK cell monolayers. Trophozoites of each species were added at a 1:2 trophozoite:cells ratio. A faster decrease of TER was observed in *E. histolytica* when compared with *E. dispar*. At 60 min, all strains showed a loss in TER.



**Figure 4.** Proteolytic activity of crude extracts of *E. histolytica* (HM1: IMSS strain) and *E. dispar* ICB-ADO monoxenic and polixenic strains using Azocoll substrate. The data represent the mean  $\pm$  SD of at least three independent experiments (\* $p < 0.05$ ).



**Figure 5.** Proteolytic activity of crude extracts of *E. dispar* ICB-ADO monoxenic and polyxenic strains using Azocoll substrate and the effect of protease inhibitors (PHMB, PMSF and EDTA). Data represent the mean  $\pm$  SD of at least three independent experiments (\* $p < 0.05$ ).

same occurred in the case of the monoxenic strain, although with a less intensity. The PMSF and EDTA inhibitors of serine and metalloproteases did not reduce the proteolytic activities.

## DISCUSSION

Most of the current knowledge regarding the biology and physiopathology of *E. dispar* is based on a unique species isolated from the feces of a single English male who was an asymptomatic cyst passer; this strain is classified as SAW760 and cultured axenically.<sup>24</sup> However, little is known about other isolates of the genetically similar *E. dispar* species that has been cultured monoxenically or xenically by other groups.<sup>5,16,17,25</sup> Previous work on the pathogenicity and virulence of *E. dispar* support the idea that this strain, in certain conditions, such as alterations of the immune system or interactions with bacteria of host intestinal flora, can produce a pathogenic behavior in humans, contrarily as was originally proposed by Brumpt<sup>26</sup> that *E. dispar* was always a non-pathogenic strain. Moreover, there are some reports in the literature using other different isolates of *E. dispar*, such as those described by Costa, *et al.*,<sup>17</sup> Furst, *et al.*<sup>18</sup> and Costa, *et al.*,<sup>25</sup> which have shown the production of liver damage in experimental animals. The scarce information available for the biological behavior of other strains, different from the original *E. dispar* SAW760, has been a direct result of the technical difficulties associated with axenic cul-

ture conditions, and has impeded our understanding of the pathogenic capabilities of this species. Thus, it is necessary to perform more experimental studies using several *E. dispar* isolates to attain a better understanding of their pathogenic potential and to determine whether this species is capable of invading the mammalian intestine and liver.

To this end, we investigated the *E. dispar* ICB-ADO strain that was originally isolated from a stool sample of a symptomatic Brazilian patient with non-dysenteric colitis. This particular isolate has been successfully cultured by others in monoxenic and xenic conditions<sup>17</sup> and has been genotyped using different targets of polymorphic intergenic DNA region corresponding to tRNA genes (SQD3-5, Dsp5-6, StgD35, and NKD3-5). The phylogenetic reconstruction performed by the Neighbor-Joining method for *E. dispar*, *E. histolytica* and ICB-ADO strains using the sequences of PCR products obtained with the appropriated primers clearly showed that the ICB-ADO strain is included in the *E. dispar* group in all phylogenies.<sup>6</sup> In addition, this work demonstrates the presence of genetic markers of *E. dispar* in aspirates from liver abscess of different patients, indicating the possible invasive role of *E. dispar* species. The authors discuss the importance of a new data observed in *E. dispar* emphasizing the importance of this species in the damage of intestine and liver. Therefore, our results introduce again the important question: whether *E. dispar* is a simple commensal parasite (Brumpt hypothesis) or is capable to produce, in certain conditions, important lesions as described by several authors? We have raised four important hypotheses regarding the physiopathology of the amoebic liver abscess:

- First, *E. histolytica* invades the intestinal mucosa and then, both species (*E. histolytica* and *E. dispar*) reach the liver through the portal circulation. *E. dispar* does not produce tissue damage but may take advantage of the pathogenic capacity of the *E. histolytica* species.
- Second possible explanation proposes that the two species are similarly responsible of both the intestinal and liver damage.
- Third hypothesis deals with bacteria-mediated pathogenicity. This suggests that the pathogenesis of at least some *E. dispar* strains may be mediated by a type of bacterial flora in a particular host flora.
- The last hypothesis suggests that a process of recombination event between *E. histolytica* and *E. dispar* species may be occurring.

Therefore, these previously mentioned hypotheses are introducing again to the highly important discussion, whether *E. dispar* is just a commensal strain in the human gut or it has the ability to invade tissues and organs.

The induction of amoebic liver abscess (ALA) in the hamster model (susceptible to liver damage) was performed by injecting the xenic trophozoites of this ICB-ADO strain and resulted in serious hepatic damage, surpassing 40% of the total organ weight by 7 days post-infection. This unexpected result led us to question the variable pathogenic potential of all available strains of *E. dispar*. The previously reported histological events preceding the formation of the liver abscess in hamsters<sup>27-29</sup> served as a basis by which we compared the morphological and other biological findings after inoculation with the *E. dispar* ICB-ADO strain. In the early stages of infection (1-12 h) we observed inflammatory foci that gradually increased in size and number; a small number of amoebae were associated with the inflammation and some showed an atypical morphology. One of the major findings at 6 h after intrahepatic inoculation was the presence of trophozoites in close contact with hepatocytes where the liver parenchyma displayed signs of damage (unpublished results). These findings differed from those published by Tsutsumi, *et al.*<sup>26</sup> and Chadee & Meerovitch,<sup>30</sup> which indicated that amoebae surrounded by inflammatory infiltrates were lysed and in turn damaged the hepatic parenchyma. Another important difference with the previous reports of ALA produced by *E. histolytica* in hamster and gerbil models<sup>28,30</sup> is that in our *E. dispar* infected hamsters the granulomatous reaction was practically absent. The liver lesions produced in our study by the *E. dispar* ICB-ADO strain (although with slow evolution at the beginning) progressed in a rapid manner after 24 h post-intrahepatic inoculation and resulted in extensive necrotic areas, as compared to the damage induced by *E. histolytica* HM1: IMSS.

The possible role of host associated bacteria present in ICB-ADO strain cultured xenically was discarded after inoculating only the bacterial flora in hamster livers. This procedure resulted only in a small periportal inflammatory infiltrate in animals euthanized at 7 days post-inoculation. In addition, inoculation of *E. dispar* monoxenic strain ICB-ADO did not cause any damage to the liver of the animals, only a scarce inflammatory reaction was observed in the portal areas (data not shown).

Studies reported by Espinosa-Cantellano, *et al.*<sup>16</sup> using an axenic strain of *E. dispar* (strain SAW) de-

monstrated that trophozoites of this specific strain presented a discreet cytopathic effect on MDCK cells, requiring double the time to reach equal *E. histolytica* capacity to completely break down the transepithelial electrical resistance. To confirm the pathogenic behavior of our *E. dispar* species (ICB-ADO), we used MDCK cells and estimated the cell damage by measuring the TER. This technique is a highly sensitive procedure to detect the early damages produced by trophozoites in a well-established and controlled cell monolayer.<sup>31</sup> The extent of cell damage depends on the strain and the trophozoite/epithelial cells ratio. We observed that *E. dispar* ADO xenic strain presented less damage to MDCK cells than did the *E. histolytica* strain HM1-IMSS.

Some data have suggested that adhesion constitutes an important and crucial phenomenon during the early cell damage by *E. histolytica* trophozoites.<sup>32,33</sup> It is possible that in the case of *E. dispar* in xenic culture, at the beginning of an *in vitro* interaction with MDKC cells, the amoeba-bacteria association may act to physically interfere with the adhesion capability to cell monolayers. A similar work carried out by Costa, *et al.*,<sup>17</sup> using VERO cell line cultures, indicated a small degree of cell destruction occurred. However, at later stages the presence of bacteria associated with amoeba may instead stimulate some virulence factors produced and secreted by the parasite, such as the proteases; this phenomenon may characterize the pathophysiology of intestinal amoebiasis.<sup>34</sup>

To examine the proteolytic activity of this amoeba as mechanisms involved in tissue destruction, we used Azocoll as a chromogen substrate. The *E. dispar* ICB-ADO xenic strain displayed a strong proteolytic activity, a particularly surprising result since *E. dispar* is widely considered to be a poor protease producer.<sup>35</sup> Since the bacteria alone were unable to produce significant protease activity, the protease activity observed in the ADO-ICB xenic strain was presumed to be due to the trophozoites of *E. dispar*.

Several studies have suggested that bacteria/*E. histolytica* association has a significant effect on the trophozoites behavior, thus modulating the phenotypical changes and virulence properties of amoebae.<sup>13,33,36-38</sup> Studies reported by Bruchhaus, *et al.*<sup>39</sup> demonstrated that only three (*ehcp1*, *ehcp2*, *ehcp5*) of 50 genes of cysteine proteases identified in the *E. histolytica* genome<sup>40</sup> were strongly expressed under culture conditions. However, Singh, *et al.*<sup>41</sup> demonstrated that the association of *E. histolytica*/*E. coli*, resulted in a strong activity of cysteine pro-

teases, a high expression of the *ehcp2* and *ehcp5* genes, and significant cellular cytotoxicity. The regulation of these genes in the proteolytic pathway may also occur in some strains of *E. dispar* when associated with bacteria.

Early studies conducted through the interaction of *E. histolytica* trophozoites with bacteria have shown that after this interaction the virulence of the amoebae was amplified, increasing their cytopathic effect to epithelial cells and amoebic liver abscesses in hamster model.<sup>42-44</sup> Recent studies by Galván-Moroyoqui, *et al.*<sup>34</sup> have demonstrated that following association of *E. histolytica* with enteropathogenic bacteria there is an increase of Gal/GalNAc lectin expression on the surface of trophozoites, which results in an increased adhesion capacity and higher cytopathic effect. Moreover, these authors demonstrated that the presence of pathogenic enterobacteria could induce changes in epithelial cells (production of pro-inflammatory cytokines) that facilitate adhesion capacity and tissue damage induced by the invasive trophozoites.

It has been proposed that bacteria in the gastrointestinal tract may be one of the factors that modify or modulate the virulence of the trophozoites of *E. histolytica*, experiments *in vitro* have shown that pathogenic bacteria can increase the virulence of *E. histolytica* causing further damage and adhesion to target cells.<sup>34</sup> In this context it is possible that *E. dispar* naturally interact with pathogenic bacteria in the intestinal tract, however, previous reports have shown that the interaction of *E. dispar* SAW760 with pathogenic bacteria does not change the virulence of this strain of amoeba.<sup>34</sup> In contrast, here we show that xenical culture with bacteria increases the virulence of *E. dispar* ADO causing damage to cultured epithelial cells (MDCK), enhancing the protease activity and producing larger amoebic liver abscess in hamsters.

Studies using the xenic strains of *E. dispar* are scarce in the literature<sup>17,18,25</sup> and, of those, very few have sought to correlate or explain the influence of bacteria on these *E. dispar* strains. The emerging recognition of the functional relationships that exist between commensal communities, the host immune system, and the susceptibility to inflammatory diseases emphasizes the need for a better understanding of the pathways that initiate and maintain amoebae-bacterial symbiosis.

The paradigm that only *E. histolytica* should be considered as a potential pathogen that can cause serious damage to the large intestine and extraintestinal organs must be reconsidered, since recent stu-

dies have shown that *E. dispar* can act as a potential agent capable to produce tissue damage in humans. In the work presented herein we have demonstrated that *E. dispar* may assume a pathogenic capacity when cultured in xenic conditions. This can take place during the colonization of the gut by this amoeba, which interacts not only with commensal bacteria but also with pathogenic bacteria, which could have an important role in the invasive form of the disease. Our results reinforce the existence of potentially pathogenic strains of *E. dispar*, and this in turn may change the epidemiological situation (based on molecular biology findings) of the complex *E. histolytica*/*E. dispar* with an important impact in the treatment, similarly as occurs with patients with *E. histolytica* infection. Currently, we are carrying out additional studies focused on the specific role of bacteria related to the pathogenesis of *E. dispar*. In summary, our results strongly suggest that the presence of bacteria could contribute to the regulation of virulence factors, including protease activities and, consequently, may have an important impact on the manifestation of virulence by *E. dispar* ADO strain in xenic culture.

## ABBREVIATIONS

- **Gal/GalNAc:** Galactose/N-acetyl Galactosamine.
- **PBS:** phosphate buffer saline.
- **MDCK:** Madin-Darby canine kidney.
- **MEM:** minimum essential medium.
- **FBS:** fetal bovine serum.
- **TER:** transepithelial resistant.
- **PHMB:** P-Hydroxy-MercuriBenzoic acid.
- **PMSF:** PhenylMethylSuphonyl fluoride.
- **EDTA:** EthyleneDiamineTetracetic acid.

## ACKNOWLEDGEMENTS

We wish to thank to João da Costa Viana, Edna Maria Pires, Silvia Galindo-Gómez, Liliana Rojas and Patricia Morán for their valuable technical assistance.

## FINANCIAL SUPPORT

This work was partially support by Grant No 79220 from the National Council for Sciences and Technology (CONACyT) Mexico and Foundation for Research Development of Minas Gerais State (FAPEMIG), Brazilian Coordination for the Improvement of Higher Education Personnel (CAPES).

## REFERENCES

1. Sargeant PG, Williams JE, Grene JD. The differentiation of invasive and non-invasive *Entamoeba histolytica* by isoenzyme electrophoresis. *Trans R Soc Trop Med Hyg* 1978; 72: 519-21.
2. Strachan WD, Chiodini PL, Spice WM, Moody AH, Ackers JP. Immunological differentiation of pathogenic and non-pathogenic isolates of *Entamoeba histolytica*. *Lancet* 1988; 1: 561-8.
3. Tannich E, Horstmann RD, Knobloch J, Arnold HH. Genomic DNA differences between pathogenic and nonpathogenic *Entamoeba histolytica*. *Proc Natl Acad Sci USA* 1989; 86: 5118-22.
4. WHO, Organization WH, PAHO-UNESCO W. Report of a consultation of experts on amoebiasis. Mexico; 1997.
5. Costa AO, Viana JC, Assis D, Rocha OA, Silva EF. Comparison of xenic and monoxenic *Entamoeba dispar* cultures using hepatic inoculation in hamster. *Arch Med Res* 2000; 31(4 Suppl.): S247-S248.
6. Ximenez C, Cerritos R, Rojas L, Dolabella S, Moran P, Shibayama M, González E, et al. Human amebiasis: breaking the paradigm? *Int J Environ Res Public Health* 2010; 7: 1105-20.
7. Mann BJ, Dodson JM, Schroeder JW, Lockhart LA, Leo B, Clark CG. Structure and function of the galactose/N-acetyl D-galactosamine inhibitable adhesin of *Entamoeba dispar*. *Arch Med Res* 1997; 28(Spec No.): 168-9.
8. Petri WA Jr., Smith RD, Schlesinger PH, Murphy CF, Ravdin JI. Isolation of the galactose-binding lectin that mediates the in vitro adherence of *Entamoeba histolytica*. *J Clin Invest* 1987; 80: 1238-44.
9. Pillai DR, Kobayashi S, Kain KC. *Entamoeba dispar*: molecular characterization of the galactose/N-acetyl-d-galactosamine lectin. *Exp Parasitol* 2001; 99: 226-34.
10. North MJ, Mottram JC, Coombs GH. Cysteine proteinases of parasitic protozoa. *Parasitol Today* 1990; 6: 270-5.
11. Reed SL, Keene WE, McKerrow JH, Gigli I. Cleavage of C3 by a neutral cysteine proteinase of *Entamoeba histolytica*. *J Immunol* 1989; 143: 189-95.
12. Leippe M. Amoebores. *Parasitol Today* 1997; 13: 178-83.
13. Bhattacharya A, Ghildyal R, Prasad J, Bhattacharya S, Diamond LS. Modulation of a surface antigen of *Entamoeba histolytica* in response to bacteria. *Infect Immun* 1992; 60: 1711-3.
14. Campos-Rodríguez R, Jarillo-Luna A. The pathogenicity of *Entamoeba histolytica* is related to the capacity of evading innate immunity. *Parasite Immunol* 2005; 27: 1-8.
15. Meerovitch E, Chadee K. In vivo models for pathogenicity in amebiasis. In: Ravdin J (ed.). *Amebiasis Human infection by Entamoeba histolytica*. 1988, p. 177-90.
16. Espinosa-Cantellano M, González-Robles A, Chávez B, Castañón G, Argüello C, Lazaro-Haller A, Martínez-Palomo A. *Entamoeba dispar*: ultrastructure, surface properties and cytopathic effect. *J Eukaryot Microbiol* 1998; 45: 265-72.
17. Costa AO, Gomes MA, Rocha OA, Silva EF. Pathogenicity of *Entamoeba dispar* under xenic and monoxenic cultivation compared to a virulent *E. histolytica*. *Rev Inst Med Trop Sao Paulo* 2006; 48: 245-50.
18. Furst C, Gomes MA, Tafuri WL, Silva EF. Biological aspects of a Brazilian strain of *Entamoeba dispar*. *Pathologica* 2002; 94: 22-7.
19. Shibayama M, Dolabella SS, Silva EF, Tsutsumi V. A Brazilian species of *Entamoeba dispar* (ADO) produces amoebic liver abscess in hamsters. *Ann Hepatol* 2007; 6: 117-8.
20. Diamond LS, Harlow DR, Cunick CC. A new medium for the axenic cultivation of *Entamoeba histolytica* and other *Entamoeba*. *Trans R Soc Trop Med Hyg* 1978; 72: 431-2.
21. Gomes MA, Pesquero JB, Furst C, Valle PR, Pesquero JL, Silva EF. An improved method to distinguish *Entamoeba histolytica* and *Entamoeba dispar*. *Parasitology* 1999; 119: 359-62.
22. Clark CG, Diamond LS. Differentiation of pathogenic *Entamoeba histolytica* from other intestinal protozoa by riboprinting. *Arch Med Res* 1992; 23: 15-6.
23. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976; 72: 248-54.
24. Clark CG. Axenic cultivation of *Entamoeba dispar* Brumpt 1925, *Entamoeba insolita* Geiman and Wichterman 1937 and *Entamoeba ranarum* Grassi 1879. *J Eukaryot Microbiol* 1995; 42: 590-3.
25. Costa CA, De Brito KN, Gomes MA, Caliri MV. Histopathological and immunohistochemical study of the hepatic lesions experimentally induced by *Entamoeba dispar*. *Eur J Histochem* 2010; 54: e39.
26. Brumpt E. Étude sommaire de l'*Entamoeba dispar* n. sp. Amibe à kystes quadrinucléés, parasite de l'homme. *Bull Acad Med* 1925; 94: 943-52.
27. Tsutsumi V, Martínez-Palomo A. Inflammatory reaction in experimental hepatic amebiasis. An ultrastructural study. *Am J Pathol* 1988; 130: 112-9.
28. Tsutsumi V, Mena-Lopez R, Anaya-Velazquez F, Martínez-Palomo A. Cellular bases of experimental amebic liver abscess formation. *Am J Pathol* 1984; 117: 81-91.
29. Tsutsumi V, Shibayama M. Experimental amebiasis: a selected review of some in vivo models. *Arch Med Res* 2006; 37: 210-20.
30. Chadee K, Meerovitch E. The pathogenesis of experimentally induced amebic liver abscess in the gerbil (*Meriones unguiculatus*). *Am J Pathol* 1984; 117(1): 71-80.
31. Orozco ME, Fernandez Castelo S, Martínez Palomo A. Detection of early cellular damage produced by *Entamoeba histolytica* in epithelia. *Arch Invest Med* 1982; 13(Suppl. 3): 169-75.
32. Ravdin JI, Guerrant RL. A review of the parasite cellular mechanisms involved in the pathogenesis of amebiasis. *Rev Infect Dis* 1982; 4: 1185-207.
33. Silva EF, Gomes MA, Barral de Martínez AM. *Entamoeba histolytica*: axenization and characterization of isolated samples from symptomatic and asymptomatic patients from different regions of Brazil. *Arch Med Res* 1997; 28(Spec No.): 288-9.
34. Galvan-Moroyoqui JM, Del Carmen Dominguez-Robles M, Franco E, Meza I. The interplay between *Entamoeba* and enteropathogenic bacteria modulates epithelial cell damage. *PLoS Negl Trop Dis* 2008; 2: e266.
35. Talamas-Rohana P, Aguirre-García MM, Anaya-Ruiz M, Rosales-Encina JL. *Entamoeba dispar* contains but does not secrete acid phosphatase as does *Entamoeba histolytica*. *Exp Parasitol* 1999; 92: 219-22.
36. Padilla-Vaca F, Ankri S, Bracha R, Koole LA, Mirelman D. Down regulation of *Entamoeba histolytica* virulence by monoxenic cultivation with *Escherichia coli* O55 is related to a decrease in expression of the light (35-kilodalton) subunit of the Gal/GalNAc lectin. *Infect Immun* 1999; 67: 2096-102.
37. Bhattacharya A, Anand MT, Paul J, Yadav N, Bhattacharya S. Molecular changes in *Entamoeba histolytica* in response to bacteria. *J Eukaryot Microbiol* 1998; 45: 285-335.
38. Bhattacharya A, Ghildyal R, Prasad J, Bhattacharya S, Diamond LS. Modulation of cell surface antigens of *Entamoeba*

- histolytica in response to bacteria. *Infect Immun* 1992; 60: 711-3.
39. Bruchhaus I, Loftus BJ, Hall N, Tannich E. The intestinal protozoan parasite *Entamoeba histolytica* contains 20 cysteine protease genes, of which only a small subset is expressed during in vitro cultivation. *Eukaryot Cell* 2003; 2: 501-9.
40. Loftus B, Anderson I, Davies R, Alsmark UC, Samuelson J, Amedeo P, Roncaglia P, et al. The genome of the protist parasite *Entamoeba histolytica*. *Nature* 2005; 433: 865-8.
41. Singh D, Naik SR, Naik S. Role of cysteine proteinase of *Entamoeba histolytica* in target cell death. *Parasitol* 2004; 129: 127-35.
42. Mirelman D, Bracha R, Chayen A, Aust-Kettis A, Diamond LS. *Entamoeba histolytica*: effect of growth conditions and bacterial associates on isoenzyme patterns and virulence. *Exp Parasitol* 1986; 62: 142-8.
43. Mirelman D. Ameba-bacterium relationship in amebiasis. *Microbiol Rev* 1987; 51: 272-84.
44. Sepúlveda B, Tanimoto-Weki M, Guerrero A, Solís G. Immunity in hamsters after vaccination with monoxenic acid axenic cultures of *E. histolytica*. *Arch Invest Med* 1973; Suppl 1: 159-64.