


phosphatase (ALP) $\geq 1.67 \times \text{ULN}$, and total bilirubin $\leq 2 \times \text{ULN}$; they received seladelpar 10 mg or placebo (2:1 randomisation) for 12 months. Cirrhosis was defined by medical history, liver biopsy, transient elastography, laboratory findings, and radiological features. Safety and changes in laboratory parameters were assessed.

Results: Of 193 patients, 27 (14%) had Child-Pugh A compensated cirrhosis at baseline (18 seladelpar, 9 placebo). Mean ALP change was -121.4 U/L for seladelpar vs 23.2 U/L for placebo patients with cirrhosis, and -134.8 U/L for seladelpar vs -18.0 U/L for placebo patients without cirrhosis. Greater decreases in other laboratory parameters were observed with seladelpar vs placebo regardless of cirrhosis status. Adverse events with seladelpar vs placebo were similar in patients with and without cirrhosis. No patients with cirrhosis discontinued seladelpar due to adverse events. Elevations in alanine aminotransferase or aspartate aminotransferase of $>3 \times \text{ULN}$ occurred in 3 patients with cirrhosis.

Conclusions: Seladelpar reduced biomarkers of cholestasis and was overall safe and well tolerated in patients with PBC with or without cirrhosis.

Conflict of interest: Yes, Gilead Sciences, Inc.

Changes in ALP in Patients With PBC With or Without Cirrhosis at Baseline Treated With Seladelpar or Placebo in the RESPONSE Trial



*P<.0001. **P<.05

ALP, alkaline phosphatase; BL, baseline; LS, least squares; PBC, primary biliary cholangitis; SE, standard error.

<https://doi.org/10.1016/j.aohep.2025.102052>

#131

CLINICAL AND EPIDEMIOLOGICAL CHARACTERISTICS OF HEPATOCELLULAR CARCINOMA IN A REFERRAL HOSPITAL IN LIMA, PERU

Fortunato Sandro Príncipe Meneses¹, Zuly Placido Damian², Estefania Liza Baca², Javier Díaz Ferrer²

¹ Unidad de Investigación. Servicio de Hígado. Hospital Nacional Edgardo Rebagliati Martins. Essalud. Lima. Perú.

² Servicio de Hígado. Hospital Nacional Edgardo Rebagliati Martins. Essalud. Lima. Perú.

Introduction and Objectives: Hepatocellular carcinoma (HCC) is the most common primary liver tumor (85–90%) and one of the leading causes of death among cirrhotic patients.

To determine the clinical and epidemiological characteristics of patients with HCC at a referral hospital in Lima, Peru.

Materials and Methods: A cross-sectional observational study including 282 patients diagnosed with HCC at the Liver Unit of HNERM-EsSalud between 2016 and 2023.

Results: 62% of patients were male, with a mean age of 65.4 years (range: 16–92). Cirrhosis was present in 78.4% of cases. The most common etiology was MASLD (52%), followed by hepatitis B (16.3%) and hepatitis C (13%). Liver function was classified as Child-Pugh A in 53%, B in 30%, and C in 17%. Only 40% were enrolled in a screening

program. Tumor stage according to BCLC was: 0–A in 40%, B in 18.6%, C in 6%, and D in 33%. Serum AFP>200 ng/mL was observed in 45% of cases. Treatments included transarterial chemoembolization (21%), radiofrequency ablation (5.3%), surgery (15.6%), liver transplantation (6.4%), systemic therapy (6%), and palliative care (39%).

When comparing cirrhotic vs non-cirrhotic patients, hepatitis B was more frequent in the non-cirrhotic group (P<0.001), with larger tumors (11.1 cm vs. 5.3 cm, P<0.001), higher AFP levels, and lower screening rates.

Conclusions: MASLD was the leading cause of HCC overall, while hepatitis B predominated in non-cirrhotic patients. Only 40% underwent screening. Patients in early stages had access to better treatment options.

Conflict of interest: None

Variable	Non-cirrhosis (N=61)	Cirrhosis (N=221)	p-value
Age (mean, SD)	56.2 (18.1)	67.9 (9.8)	<0.001
Sex			0.733
Female	22 (36.1%)	85 (38.5%)	
Male	36 (63.9%)	136 (61.5%)	
Etiology			0.001
MASLD	22 (36.1%)	125 (56.6%)	
Others	14 (23%)	39 (17.6%)	
HVB	19 (31.1%)	27 (12.2%)	
HVC	6 (9.8%)	30 (13.6%)	
Diabetes			<0.001
Yes	11 (18%)	93 (42.1%)	
No	50 (82%)	128 (57.9%)	
Hypertension			0.057
Yes	18 (29.5%)	95 (43%)	
No	43 (70.5%)	126 (57%)	
Coronary artery disease			0.193
Yes	0 (0%)	6 (2.7%)	
No	61 (100%)	215 (97.3%)	
Renal disease			0.361
Yes	2 (3.3%)	14 (6.3%)	
No	59 (96.7%)	215 (97.3%)	
Diagnosis exam for HCC			<0.001
Surveillance	4 (6.6%)	108 (48.9%)	
Symptoms	57 (93.4%)	113 (51.1%)	
Diagnosis methods			<0.001
Image	28 (45.9%)	204 (92.3%)	
Biopsy	35 (54.1%)	17 (7.7%)	
Tumor diameter (mean, SD)	11.1 (6)	5.8 (3.7)	<0.001
Nodules number			0.074
<=3	47 (77%)	191 (86.4%)	
>3	14 (23%)	30 (13.6%)	
AFP			0.015
<20	12 (19.7%)	66 (29.9%)	
20-200	13 (21.3%)	65 (29.4%)	
200-400	3 (4.9%)	20 (9%)	
>400	33 (54.1%)	70 (31.7%)	
Treatment			<0.001
Palliative	24 (39.3%)	86 (38.9%)	
TACE	0 (0%)	59 (26.7%)	
RFA	1 (1.6%)	14 (6.3%)	
Systemic	7 (11.5%)	10 (4.5%)	
Surgery	23 (37.7%)	23 (10.4%)	
Liver transplant	0 (0%)	16 (7.2%)	
Others	6 (9.8%)	13 (5.9%)	

<https://doi.org/10.1016/j.aohep.2025.102053>

#133

IMPORTANCE OF P450 POLYMORPHISM IN THE DEVELOPMENT OF DILI/HILI: HISTOLOGICAL AND BIOCHEMICAL FINDINGS

Tayna da Silva Domingos¹, Vinicius Santos Nunes¹, Luiz Antônio Rodrigues de Freitas¹, Genario Oliveira Santos Júnior¹, Raymundo Paraná Ferreira Filho¹, Maria Schinoni Protti¹

¹ Hospital Universitario Prof Edgard Santos, Brasil.

Introduction and Objectives: Drug-induced liver injury (DILI) and herb-induced liver injury (HILI) represent diagnostic and prognostic challenges in hepatology. Objective: To evaluate histological, epidemiological, biochemical variables and the prevalence of cytochrome P450 (CYP450) genotypes in individuals with DILI/HILI at a hepatotoxicity clinic.

Materials and Methods: Cross-sectional study with individuals who developed DILI/HILI and underwent liver biopsy.