

centers in Argentina. Each participant underwent a rapid HCV antibody test, followed by point-of-care confirmatory testing using HCV RNA PCR when reactive. Demographic data, testing history, and general knowledge about HCV were collected.

Results: Of the 160 participants, 2 tested reactive on the rapid test and are currently undergoing confirmation by PCR. A total of 84.4% were male and 15.6% female, with a median age of 36 years. Among participants, 95% were unaware of having been previously tested for HCV, and 76.9% had no knowledge about the disease. Additionally, 74.4% lacked medical insurance coverage.

Conclusions: This preliminary study highlights the feasibility of implementing HCV micro-elimination strategies in addiction centers. The findings emphasize the limited awareness of HCV and the absence of prior testing in most participants, underlining the importance of targeted screening and educational interventions in this high-risk population.

Conflict of interest: None

<https://doi.org/10.1016/j.aohep.2025.102082>

#179

ASSESSMENT OF QUALITY OF LIFE IN PATIENTS WITH PRIMARY BILIARY CHOLANGITIS USING THE PBC-27 QUESTIONNAIRE AT A TERTIARY CARE CENTER

Deborah Espinoza Lopez¹, Rodrigo Toledo Galvan¹, Maria Beatriz Jourdan Rodriguez¹, Viridiana Lopez Ladron de Guevara¹, Maria de Fatima Higuera de la Tijera¹

¹ Hospital General de México Dr. Eduardo Liceaga.

Introduction and Objectives: Primary biliary cholangitis (PBC) is a chronic liver disease that significantly impacts patients' quality of life due to persistent symptoms like fatigue, pruritus, and emotional or cognitive disturbances. Understanding these effects is vital for guiding patient-centered care.

This study aimed to assess quality of life in patients with PBC using the validated PBC-27 questionnaire, identifying the most affected domains.


Materials and Methods: We conducted a cross-sectional, observational, and descriptive study in 50 adult patients with confirmed PBC, evaluated at a tertiary care center between March and May 2025. Variables included age, sex, comorbidities, disease duration, and PBC-27 domain-specific scores. Descriptive statistics were used, with medians and interquartile ranges for continuous variables, and absolute and relative frequencies for categorical data.

Results: All 50 patients completed the PBC-27 questionnaire. The mean overall score was 77.1 ± 4.2 , ranging from 68.5 to 86.2. Fatigue was the most affected domain, with a mean score of 22.6 ± 3.2 , followed by general symptoms (16.8 ± 2.9), pruritus (10.3 ± 2.1), and cognitive symptoms (8.9 ± 1.8). Emotional symptoms were also prevalent (8.7 ± 2.0). Domains related to dry mouth and dry eyes had the lowest impact, with mean scores of 5.3 ± 1.7 and 5.1 ± 1.5 , respectively.

Conclusions: These findings underscore the multidimensional burden of PBC on quality of life, particularly in physical and neuro-cognitive domains. Regular assessment using structured tools like the PBC-27 is crucial for tailoring follow-up and improving patient outcomes.

Conflict of interest: None

Chart 1. PBC-27 Questionnaire Results



<https://doi.org/10.1016/j.aohep.2025.102083>

#180

CORRELATION BETWEEN LI-RADS CLASSIFICATION AND HISTOPATHOLOGICAL FINDINGS IN PATIENTS WITH SUSPECTED HEPATOCELLULAR CARCINOMA

Alonso Vera Torres¹, Valentina Mejia Valdes¹, Diana Fernanda Bejarano Ramirez¹

¹ Hospital Universitario Fundación Santa Fe de Bogotá, Colombia.

Introduction and Objectives: The Liver Imaging Reporting and Data System (LI-RADS) standardizes the interpretation of liver imaging in patients at risk of hepatocellular carcinoma (HCC). However, its correlation with histopathological parameters has not been sufficiently validated in Latin American settings.

To evaluate the correlation between LI-RADS classification assigned through contrast-enhanced computed tomography (CT) or magnetic resonance imaging (MRI) and histopathological findings in patients with suspected hepatocellular carcinoma treated at Fundación Santa Fe de Bogotá.

Materials and Methods: A retrospective observational study was conducted in patients with histologically confirmed HCC and at least one contrast-enhanced imaging study available for blinded review. Clinical, radiological, and histopathological variables were collected. The correlation between LI-RADS category and tumor differentiation grade (Edmondson-Steiner scale) was evaluated using the chi-square test. Statistical analysis was performed with SPSS v25, and a p-value < 0.05 was considered statistically significant.

Results: A total of 154 patients were included, with a mean age of 64.4 years (95% CI: 63.3–65.4) and a male predominance (65%). Most patients were classified as Child-Pugh A or B and had solitary tumors ≤ 3 cm. No statistically significant association was found between the LI-RADS category and Edmondson-Steiner grade ($\chi^2 = 18.77$, $p = 0.537$). Histological grading was heterogeneously distributed within LI-RADS categories 4 and 5, with no clear predominance of poorly differentiated tumors.

Conclusions: In this cohort, LI-RADS classification did not show a significant correlation with histological differentiation of HCC. Although LI-RADS remains a useful non-invasive diagnostic tool, it should not be used in isolation to infer tumor aggressiveness. Further studies with greater statistical power are required to validate its use as a predictor of tumor histology in Latin American clinical contexts.

Conflict of interest: None