



# BRAZILIAN JOURNAL OF MICROBIOLOGY

<http://www.bjmicrobiol.com.br/>



## Environmental Microbiology

# Phylogenetic MLSA and phenotypic analysis identification of three probable novel *Pseudomonas* species isolated on King George Island, South Shetland, Antarctica



Felipe Vásquez-Ponce, Sebastián Higuera-Llantén, María S. Pavlov, Sergio H. Marshall, Jorge Olivares-Pacheco\*

Pontificia Universidad Católica de Valparaíso, Facultad de Ciencias, Instituto de Biología, Laboratorio de Genética e Inmunología Molecular, Valparaíso, Chile

### ARTICLE INFO

#### Article history:

Received 13 August 2017

Accepted 14 February 2018

Available online 15 March 2018

Associate Editor: Vânia Melo

#### Keywords:

*Pseudomonas*

Antarctica

*P. fluorescens* complex

Multilocus Sequence Analysis

### ABSTRACT

Antarctica harbors a great diversity of microorganisms, including bacteria, archaea, microalgae and yeasts. The *Pseudomonas* genus is one of the most diverse and successful bacterial groups described to date, but only eight species isolated from Antarctica have been characterized. Here, we present three potentially novel species isolated on King George Island. The most abundant isolates from four different environments, were genotypically and phenotypically characterized. Multilocus sequence analysis and 16S rRNA gene analysis of a sequence concatenate for six genes (16S, *aroE*, *glnS*, *gyrB*, *ileS* and *rpoD*), determined one of the isolates to be a new *Pseudomonas mandelii* strain, while the other three are good candidates for new *Pseudomonas* species. Additionally, genotype analyses showed the three candidates to be part of a new subgroup within the *Pseudomonas fluorescens* complex, together with the Antarctic species *Pseudomonas antarctica* and *Pseudomonas extremaustralis*. We propose terming this new subgroup *P. antarctica*. Likewise, phenotypic analyses using API 20 NE and BIOLOG® corroborated the genotyping results, confirming that all presented isolates form part of the *P. fluorescens* complex. *Pseudomonas* genus research on the Antarctic continent is in its infancy. To understand these microorganisms' role in this extreme environment, the characterization and description of new species is vital.

© 2018 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Antarctica is arguably one of the planet's harshest environments.<sup>1</sup> Intense cold, low humidity, a limited availability of liquid water, as well as periods of high solar radiation and extended total darkness, mean that the continent

\* Corresponding author.

E-mail: [jorge.olivares@pucv.cl](mailto:jorge.olivares@pucv.cl) (J. Olivares-Pacheco).

<https://doi.org/10.1016/j.bjm.2018.02.005>

1517-8382/© 2018 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

represents a relatively inhospitable environment for the development of the life.<sup>1</sup> Nevertheless, Antarctica is now considered to host a very high diversity of microorganisms, including bacteria, archaea, yeasts and microalgae.<sup>2,3</sup> As a result of intense selective pressures throughout their evolutionary history, all of these microorganisms are fully adapted to the Antarctic conditions.<sup>4,5</sup> These adaptations are of great interest to the biotechnology industry, since they harbor the possibility of obtaining bio-products that could be used in various processes of conservation at low temperatures.<sup>6</sup> There is therefore an increasing need to investigate the diversity and to exploit the potential of the microorganisms of the cold continent.

A bacterial genus that has appeared frequently in Antarctic research is *Pseudomonas*. One of the most well-studied bacterial genera, it is present in most environments and can be considered one of the most successful bacterial groups on Earth.<sup>7</sup> *Pseudomonas*' success is mainly based on its low nutritional requirements and great metabolic diversity, which allows it to use numerous organic compounds as a source of both carbon and energy.<sup>8,9</sup> In addition, these bacteria produce a large amount of secondary metabolites, which are essential for their survival and even enable some species to degrade aromatic and halogenated compounds.<sup>10,11</sup> Based on these characteristics, the bacteria of the genus *Pseudomonas* represent an interesting alternative for production of different and biotechnologically useful molecules.

Despite being extremely well-studied on a global level, little is known about the species diversity of the genus *Pseudomonas* in Antarctica. To date, only eight species have been described on this continent: *Pseudomonas antarctica*, *Pseudomonas meridiana*, *Pseudomonas proteolytica*,<sup>12</sup> *Pseudomonas guineae*,<sup>13</sup> *Pseudomonas extremaustralis*,<sup>14</sup> *Pseudomonas deceptionensis*,<sup>15</sup> *Pseudomonas prosekii*<sup>16</sup> and, most recently, *Pseudomonas gregormendelii*.<sup>17</sup> In order to further our understanding of this genus' biogeographic distribution, as well as its function as part of the ecosystem of the extreme Antarctic environment, it is of utmost importance to identify and describe new Antarctic species of *Pseudomonas*. With this in mind, the present study aims to carry out a phylogenetic and phenotypic characterization of four bacterial isolates originating from different environments on King George Island, Antarctica. Specifically, we set the following specific objectives: (1) an evaluation of the growth rate of different strains at different temperatures; (2) the amplification and sequencing of the 16S rRNA gene; (3) the amplification and sequencing of the genes *aroE*, *glnS*, *gyrB*, *ileS* and *rpoD*, all of which were analyzed both individually and, after sequence concatenation, by Multilocus Sequence Analysis (MLSA), a robust, high-resolution technique for the identification of novel species and the confirmation of species identity of new isolates<sup>18–20</sup>; and (4) a phenotypic characterization using API 20 NE and BIOLOG GN2. Based on the both the phylogenetic analysis of 16S-rRNA gene and the MLSA results, as well as on the phenotypic characterization, we show that the four isolates belong to the genus *Pseudomonas* and specifically to the *P. fluorescens* complex, and that three of them might represent novel species, while one strain could be classified within the species *P. mandelii*.

## Materials and methods

### Strains and culture conditions

Different Antarctic environments on King George island, South Shetland archipelago, were sampled in order to isolate specifically bacteria members of the *Pseudomonas* genus. These samples were collected on Austral summer season on January 2011. Samples of seawater from the Fildes bay at 5 m depth (62°11'50.3" S; 58°54'50.3" W); freshwater from a summer lake on Fildes peninsula (62°10'9.2" S; 58°55'29" W); marine sediments from Fildes bay (62°13'28.8" S; 58°58'42.7" W) and soil from the Fildes Peninsula (62°10'9.2" S; 58°55'29" W); were taken in order to isolate bacteria from all environmental diversity. 10 L of water samples were first filtered by a 0.1 mm filters and then the samples were concentrated in a 0.2 µm filter. This filter was then resuspended in 3 mL of saline buffer (0.85% NaCl). 5 g of sediments and soil samples were resuspended in 5 mL of saline buffer solution (0.85% NaCl). 100 µL of the solution were plated onto Nutrient Agar (BD™ DIFCO™, New Jersey, NJ, USA) and incubated at 4, 10, 15 and 25 °C for 72 h. In order to isolate bacteria of the genus *Pseudomonas*, all colonies were replated on *Pseudomonas* Isolation Agar (PIA) (SIGMA–Aldrich, Los Angeles, CA, USA) and incubated at 4, 10, 15 and 25 °C for 72 h. All obtained colonies were 16S rRNA gene sequenced. The most abundant isolate of each environment was selected and designed as a particular strain. Growth rates of all selected strains were estimated in Lysogeny Broth (LB) (BD™ DIFCO™, New Jersey, NJ, USA) at 4, 10, 15, 25, 30 and 37 °C. All tests were performed in 100 mL flasks in 20 mL medium and at constant agitation rate of 240 rpm, in duplicates and on three different days (making for a total of six samples). The strain *P. fluorescens* A506 (ATCC® 31948™) served as a control for the phenotypic characterization.

### Amplification of the 16S rRNA gene and phylogenetic analysis

Genomic DNA was extracted from all strains using the GeneJET Genomic DNA Purification Kit (Thermo Fisher Scientific, Waltham, MA, USA) following the manufacturer's instructions. The 16S rRNA gene was amplified by PCR using universal primers 27F (5'-AGAGTTTGATCMTGGCTCAG-3') and 1492R (5'-GGTTACCTTGTACGACTT-3'). All PCR products were sequenced by Sanger method for five times (Forward and reverse). The obtained sequences were deposited in GenBank (NCBI) (Table S1). The sequences were then BLASTed<sup>21</sup> against the 16S rRNA GenBank database (NCBI). Sixty-one 16S rRNA gene sequences of the genus *Pseudomonas* were selected for the phylogenetic analysis. The 16S rRNA gene of *Escherichia coli* strain MG1655 K12 was used as an outgroup. Sequences of all members of the genus *Pseudomonas* were obtained from the database on the website [www.pseudomonas.com](http://www.pseudomonas.com)<sup>22</sup> and aligned using ClustalW (using standard parameters) as implemented in the software MEGA 7.0.14.<sup>23</sup> Gaps were treated with the partial deletion method. A maximum likelihood phylogenetic tree was constructed using the Tamura-Nei model. Bootstrap values were calculated based on 1000 replicas.

## MLSA

MLSA was performed on six housekeeping genes: 16S-rRNA, *aroE* (Shikimate 5-dehydrogenase), *glnS* (glutaminyl-tRNA transferase), *gyrB* (DNA gyrase subunit B), *ileS* (isoleucine-tRNA transferase), and *rpoD* (RNA polymerase sigma factor). PCR primers were taken from Andreani et al.<sup>24</sup> For all genes, PCR consisted of an initial denaturalization step of 10 min at 95 °C, followed by 30 cycles of 45 s at 95 °C; 45 s at 55 °C and 1 min at 72 °C, and a final extension step of 10 min at 72 °C. All PCR products were sequenced by Sanger method for five times (Forward and reverse). The partial sequences obtained for the genes *aroE*, *glnS*, *gyrB*, *ileS* and *rpoD* of strains IB20, 12B3 and 6A1 were deposited in GenBank (NCBI) (Table S1). The corresponding sequences for strain KG01 were taken from its whole genome sequence, which we have published previously.<sup>25</sup> The obtained sequences were individually BLASTed<sup>21</sup> against GenBank in order to identify the homologues with greatest sequence identity. For the phylogenetic analysis, we concatenated the partial gene sequences after the multisequence alignment in the following order: 16S-*aroE*-*glnS*-*gyrB*-*ileS*-*rpoD*, resulting in a single sequence. For the same genes, we took the orthologues from 51 of the species that are most representative of the genus *Pseudomonas*, whose sequences were obtained from the database on the website [www.pseudomonas.com](http://www.pseudomonas.com).<sup>22</sup> As an outgroup, we used the concatenate of the orthologues from the strain *E. coli* K12 MG1655. The phylogenetic analysis was carried out in the same way as for the 16S rRNA gene.

## Phenotypic characterization

The phenotypic characterization of the Antarctic isolates IB20, 12B3, 6A1 and KG01, as well as of the control strain *P. fluorescens* A506 (ATCC® 31948™), was carried out using API 20 NE (Biomérieux, L'Etoile, France) and BIOLOG GN2 (BIOLOG®, Hayward, CA, USA). For both API 20 NE and BIOLOG assays, we used previously isolated colonies that had been grown for 24 h at 25 °C on LB agar plates (BD™ DIFCO™, New Jersey, NJ, USA). These colonies were re-suspended in saline solution (0.85% NaCl) and inoculated according to the manufacturer's instructions. Oxidase production was determined using Oxidase Reagent Droppers (Becton, Dickinson and Company, New Jersey, NJ, USA) following the manufacturer's specifications. All phenotypic experiments were performed in triplicate using six colonies in all rounds.

## Results

### Selection of most abundant *Pseudomonas* isolates based on 16S rRNA sequence

At the first-round selection in nutrient agar, more than 6000 colonies were obtained in all temperatures used. All colonies were replated in PIA medium in order to select only members of the *Pseudomonas* genus. Only 268 colonies grow in PIA in all temperatures for all environments selected. These colonies were 16S rRNA gene sequenced and the most abundant isolates for environment were used in this work. Isolate from

**Table 1 – The optimum growth temperature of the Antarctic bacterial isolates is 25 °C. The table shows the doubling times of the four Antarctic bacterial strains at different temperatures (N/G = no growth).**

	Doubling time (min) + SD					
	4 °C	10 °C	15 °C	25 °C	30 °C	37 °C
IB20	341 ± 38	118 ± 21	65 ± 6	43 ± 8	75 ± 11	N/G
12B3	412 ± 46	132 ± 16	77 ± 4	56 ± 6	89 ± 11	N/G
6A1	381 ± 26	158 ± 18	69 ± 8	49 ± 9	78 ± 8	N/G
KG01	376 ± 31	141 ± 22	72 ± 10	48 ± 5	92 ± 6	N/G

seawater was named IB20; from freshwater 12B3; from marine sediments 6A1 and from soil KG01.


### The Antarctic bacterial isolates are psychrotolerant, with optimum growth at 25 °C

In order to determine the optimum growth conditions for the Antarctic isolates, we performed growth experiments at different temperatures. Growth curve analysis and the calculation of doubling times clearly showed that the optimum growth temperature is 25 °C for all of the Antarctic isolates (Table 1). While no growth could be detected at 37 °C, all isolates grew well at 4 °C, as is typical of psychrotolerant bacteria.<sup>26</sup> The optimum growth temperature of 25 °C is shared by *Pseudomonas* strains across latitudes. On the Antarctic South Shetland Islands, however, this temperature has not been reached for the past 25,000 years,<sup>27</sup> suggesting that members of the genus *Pseudomonas* might have colonized Antarctica before it became a polar continent.

### Phylogenetic analysis based on 16S rRNA gene and MLSA

In order to determine the genus of the Antarctic bacterial isolates, we sequenced the 16S rRNA gene. The obtained sequences were deposited in GenBank (NCBI; Table S1). In a first step, the sequences were BLASTed against NCBI's 16S rRNA GenBank.<sup>21</sup> For isolates IB20, 12B3 and KG01, sequence identity with any species contained in the database was below 97%, suggesting that all three isolates could represent new bacterial species.<sup>28</sup> However, querying the sequence of strain 6A1 revealed a 99.9% sequence identity with the strain *P. mandelii* JR-1, meaning that the isolate might in fact belong to that species. On the other hand, the phylogenetic tree (Fig. 1) shows all of these strains as part of the *P. fluorescens* lineage and within that, the *P. fluorescens* complex, one of the most diverse and ubiquitous lineages of the genus.<sup>29</sup> With regards to the other bacteria of the genus *Pseudomonas* that have been isolated in Antarctica, it can be seen that both strains of *P. guineae* form a lineage, which is separate from the ones that have been described for *P. aeruginosa* and *P. fluorescens*. However, these results are insufficient to draw conclusions.

In order to expand on the results of the 16S rRNA gene analysis, we partially sequenced the five housekeeping genes *aroE*, *glnS*, *gyrB*, *ileS* and *rpoD*. Each one of these genes was then BLASTed against the GenBank nucleotide database (NCBI).<sup>21</sup> The results are listed in Table 2. With the exception of the strain 6A1, where four of five genes show extremely high similarities (≤98%) demonstrating that this is the first strain of



**Fig. 1** – Maximum-likelihood phylogenetic tree based on the partial 16S rRNA gene sequences of members of the *Pseudomonas* genus. The 16S sequence of *E. coli* K12 was used as an outgroup. The tree shows all bacteria of the genus *Pseudomonas* isolated in Antarctica to date (**bold**). Isolates presented in this study are shown in **bold and underlined**. It can be seen that the majority of the Antarctic bacteria are part of the *P. fluorescens* lineage, with the exception of the *P. guineae* isolates, which seem to be part of a separate lineage.

*P. mandelii* isolated in Antarctica; all three strains showed low similarities in at least four genes ( $\geq 96\%$ ). These results confirm that IB20, 12B3 and KG01 are serious candidates to be classified as a novel species of the *Pseudomonas* genus.

**Concatenated sequence analysis indicates that the Antarctic strains could form a novel sub-group of the *fluorescens* complex**

The sequences of the genes 16S, *aroE*, *glnS*, *gyrB*, *ileS* and *rpoD* were concatenated to form a single sequence of 4150 bp

**Table 2** – Percentage identity values obtained by BLASTing the five housekeeping genes of the four Antarctic isolates. The letters represent the different species and strains of *Pseudomonas*, to which the percentage identity refers.

	<i>aroE</i>	<i>glnS</i>	<i>gyrB</i>	<i>ileS</i>	<i>rpoD</i>
IB20	$\leq 90\%$ <sup>g,h,i</sup>	$\leq 95\%$ <sup>h,i,j</sup>	$\leq 94\%$ <sup>i,k,l</sup>	$\leq 96\%$ <sup>h,m,n</sup>	$\leq 99\%$ <sup>o,p,q</sup>
12B3	$\leq 91\%$ <sup>g,h,i</sup>	$\leq 96\%$ <sup>h,i,j</sup>	$\leq 95\%$ <sup>i,s,t</sup>	$\leq 96\%$ <sup>h,m,n</sup>	$\leq 99\%$ <sup>o,p,q</sup>
6A1	$\leq 94\%$ <sup>a,b,c</sup>	$\leq 98\%$ <sup>a,b,y</sup>	$\leq 98\%$ <sup>a,b,d</sup>	$\leq 98\%$ <sup>a,b,e</sup>	$\leq 99\%$ <sup>a,b,f</sup>
KG01	$\leq 88\%$ <sup>m,u,v</sup>	$\leq 94\%$ <sup>w,p,q</sup>	$\leq 94\%$ <sup>i,x,w</sup>	$\leq 96\%$ <sup>i,h,n</sup>	$\leq 99\%$ <sup>p,q,w</sup>


<sup>a</sup> = *P. mandelii* JR-1, <sup>b</sup> = *P. fluorescens* NCIMB11764, <sup>c</sup> = *P. chlororapis* subsp. *aurantiaca* JD37, <sup>d</sup> = *P. fluorescens* PS1, <sup>e</sup> = *P. mandelii* DSM17967T, <sup>f</sup> = *P. fluorescens* FW300-N2E3, <sup>g</sup> = *P. fluorescens* SBW25, <sup>h</sup> = *P. antarctica* PAMC 27494, <sup>i</sup> = *P. fluorescens* LBUM 636, <sup>j</sup> = *P. fluorescens* UK4, <sup>k</sup> = *P. reactans* USB131, <sup>l</sup> = *P. reactans* USB20, <sup>m</sup> = *P. trivialis* IHBB745, <sup>n</sup> = *P. fluorescens* KENGFT3, <sup>o</sup> = *P. fluorescens* A506, <sup>p</sup> = *P. fluorescens* PICF7, <sup>q</sup> = *P. fluorescens* PCL1751, <sup>r</sup> = *P. fluorescens* NCIMB11764, <sup>s</sup> = *P. yamanorum* 8H1, <sup>t</sup> = *P. putida* MG2010, <sup>u</sup> = *P. fluorescens* L111, <sup>v</sup> = *P. fluorescens* L321, <sup>w</sup> = *P. azotoformans* S4, <sup>x</sup> = *P. reactans* USB94, <sup>y</sup> = *P. koreensis* D26.

and compared against a concatenate of orthologous genes of the genus *Pseudomonas*. A concatenate of the same genes of *E. coli* strain K12 MG1655 was used as an outgroup. Sequences were aligned, and a phylogenetic tree was constructed using the maximum likelihood model with 1000 bootstraps replicates (Fig. 2). The two major lineages of the genus *Pseudomonas*, that of *P. fluorescens* and that of *P. aeruginosa*, can be clearly distinguished from one another. Likewise, within the *P. fluorescens* lineage, three subgroups can be clearly distinguished; these represent the *P. fluorescens* complex, the *P. syringae* group and the *P. putida* group. It can be seen that all Antarctic bacteria of the genus *Pseudomonas* described to date (in bold) are part of the *P. fluorescens* complex. Together with *P. antarctica* PAMC27494 and *P. extremaustralis* 14-3, the strains described here appear to form a new subgroup within this complex, which we have called the *P. antarctica* subgroup. The strain 6A1, on the other hand, forms a cluster with *P. mandelii* JR-1, confirming that it belongs to that particular species. Likewise, *P. deceptionensis*, though part of the *P. fluorescens* complex, belongs to another, as yet undescribed, subgroup.

### Phenotypic analysis

The results of the MLSA analysis had already classified all strains as part of the *P. fluorescens* complex. To confirm this, a comprehensive phenotypic analysis of the Antarctic strains was carried out using BIOLOG GN2 and API 20 NE. The phenotypic analyses are summarized in Table 3. It is noteworthy that API 20 NE analyses classified all isolates within the species *P. fluorescens*. It is, however, known that the resolution of this technique is not particularly high, since it classifies most members of the *P. fluorescens* complex within this species.<sup>29</sup> BIOLOG analyses yielded significant differences between the Antarctic isolates and the control strain *P. fluorescens* A506 (ATCC<sup>®</sup> 31948<sup>TM</sup>). Based on these differences, three of the four new isolates can be considered good candidates for new species. The following specific findings are worth mentioning. The strain KG01 is the only one unable to assimilate D-mannose, L-proline, putrescine, pyruvic acid





**Fig. 2 – Maximum-likelihood phylogenetic tree based on the concatenated sequences of the genes 16S-aroE-glnS-gyrB-ileS-rpoD, showing the evolutionary relationship among selected members of the *Pseudomonas* genus. The bacteria isolated in Antarctica are shown in bold, and isolates presented in this study are shown in bold and underlined. The tree clearly shows the lineages of *P. aeruginosa* and *P. fluorescens*. Likewise, the *P. fluorescens* complex and its subgroups can be clearly distinguished. All Antarctic *Pseudomonas* can be classified within the *P. fluorescens* complex. Among them, *P. deceptionensis* seems to form a separate subgroup, and the isolate *Pseudomonas* sp. 6A1 appears in a branch of the *P. mandelii* subgroup. Together with the species *P. antarctica* and *P. extremaustralis*, the isolates *Pseudomonas* sp. IB20, *Pseudomonas* sp. 12B3 and *Pseudomonas* sp. KG01 form the new subgroup *P. antarctica*.**

methyl ester, L-serine, and D,L- $\alpha$ -glycerol phosphate, while being the only strain able to assimilate adipic acid and p-hydroxyphenylacetic acid. IB20 is the only strain unable to assimilate D-arabitol and propionic acid; however, it is the only one able to assimilate L-phenylalanine (weakly), formic acid (weakly) and L-alaninamide. 12B3 is the only strain unable to

assimilate malonic acid, urocanic acid and inosine. Finally, the strain 6A1 is the only one capable of assimilating N-acetyl-D-glucosamine, D-serine (weakly) and dextrin (weakly), while being unable to assimilate adonitol, D-galacturonic acid, D-glucuronic acid and L-ornithine. Additionally, this strain is unable to hydrolyze L-arginine and gelatin, thereby differing significantly from the other strains studied. Finally, all four Antarctic strains are able to assimilate D-trehalose, D-galactonic acid lactone, glycyl-L glutamic acid and D,L-carnitine, while the control, *P. fluorescens* A506 (ATCC 31948<sup>®</sup> TM), one of the most representative strains of the *P. fluorescens* complex, is not. Nevertheless, in the great majority of phenotypic tests, the results for the Antarctic strains are similar to those for the control, including the inability to assimilate nitrate, thereby corroborating that the Antarctic isolates are part of the *P. fluorescens* complex.

## Discussion

In the present study, we describe four different Antarctic isolates, which we classified as part of the genus *Pseudomonas* using both molecular and phenotypic strategies. 16S rRNA gene analysis, MLSA and phenotypic tests using BIOLOG and API 20 NE all suggest that the Antarctic isolates *Pseudomonas* sp. KG01, *Pseudomonas* sp. IB20 and *Pseudomonas* sp. 12B3 are serious candidates to classify as new species of the genus. Likewise, we determined that the isolate *Pseudomonas* sp. 6A1 is the first strain of the species *P. mandelii* to be isolated in Antarctica.

There has been little research on the genus *Pseudomonas* on the Antarctic continent, and only eight species have been described in the past 20 years. Recent research has thoroughly revised the taxonomy and classification of the genus *Pseudomonas*,<sup>29</sup> dividing it into two major lineages: that of *P. aeruginosa*, which includes almost exclusively bacteria isolated from clinical settings, and that of *P. fluorescens*, which contains most of the genus' environmental bacteria, including the species *P. putida* and *P. syringae*, both of which have been widely studied. One of the most diverse groups within this lineage is called the "*P. fluorescens* complex". It has been further subdivided into nine subgroups, *P. protegens*, *P. chlororaphis*, *P. corrugata*, *P. koreensis*, *P. jessenii*, *P. mandelii*, *P. fragi*, *P. gessardii* and *P. fluorescens*.<sup>29</sup> All Antarctic strains studied to date have been classified within this complex. This is in line with our findings for the three candidates for new species, *Pseudomonas* sp. KG01, *Pseudomonas* sp. IB20 and *Pseudomonas* sp. 12B3. Likewise, the isolate *Pseudomonas* sp. 6A1, classified in this study as a strain of the species *P. mandelii*, also forms part of this complex. An original contribution of our work is that with these three potentially new species, we have identified a novel subgroup, which we have called *P. antarctica*, since it also incorporates the strains *P. antarctica* PAMC27494 and *P. extremaustralis* 14-3. These were previously considered to be part of the subgroup *P. fluorescens*.<sup>12</sup> The hugely diverse *P. fluorescens* complex has a wide environmental distribution, and due to their versatile metabolism, most of its members play key roles at the ecological level.<sup>30,31</sup> It is worth mentioning that some of the species of the *P. fluorescens* complex play an important role in biocontrol by

**Table 3 – Phenotypic characterization of the Antarctic bacterial isolates. The phenotypic characterization was performed using BIOLOG GN2 and API 20 NE. + = positive test, – = negative test, W = weak reaction, a = API20NE, b = BIOLOG, \* = *P. fluorescens* A506 (ATCC® 31948™).**

Test (Active ingredients)	Detail	KG01	IB20	6A1	12B3	<i>P. fluorescens</i> *
D-mannose <sup>a,b</sup>	Assimilation	–	+	+	+	+
L-Proline <sup>b</sup>	Assimilation	–	+	+	+	+
L-Serine <sup>b</sup>	Assimilation	–	+	+	+	+
Putrescine <sup>b</sup>	Assimilation	–	+	+	+	W
Pyruvic Acid Methyl Ester <sup>b</sup>	Assimilation	–	+	+	+	W
D,L- $\alpha$ -Glycerol Phosphate <sup>b</sup>	Assimilation	–	+	W	+	–
D-Arabitol <sup>b</sup>	Assimilation	+	–	+	+	+
Propionic Acid <sup>b</sup>	Assimilation	+	–	W	+	+
Adonitol <sup>b</sup>	Assimilation	+	+	–	+	–
D-Galacturonic Acid <sup>b</sup>	Assimilation	+	+	–	+	–
D-Glucuronic Acid <sup>b</sup>	Assimilation	+	+	–	+	–
L-Ornithine <sup>b</sup>	Assimilation	W	+	–	+	–
Gelatine, bovine origin <sup>a</sup>	Hydrolysis	+	+	–	+	–
Malonic Acid <sup>b</sup>	Assimilation	+	W	+	–	+
Urocanic Acid <sup>b</sup>	Assimilation	+	+	+	–	+
Inosine <sup>b</sup>	Assimilation	+	+	+	–	+
Adipic acid <sup>a</sup>	Assimilation	+	–	–	–	–
p-Hydroxy Phenylacetic Acid <sup>b</sup>	Assimilation	+	–	–	–	+
L-Alaninamide <sup>b</sup>	Assimilation	–	+	–	–	–
L-Phenylalanine <sup>b</sup>	Assimilation	–	W	–	–	–
Formic Acid <sup>b</sup>	Assimilation	–	W	–	–	–
N-Acetyl-Dglucosamine <sup>b</sup>	Assimilation	–	–	+	–	+
D-Serine <sup>b</sup>	Assimilation	–	–	W	–	–
Dextrin	Assimilation	–	–	W	–	–
D-Trehalose <sup>b</sup>	Assimilation	+	+	+	+	–
D-Galactonic Acid Lactone <sup>b</sup>	Assimilation	+	+	+	+	–
Glycyl-Lglutamic Acid <sup>b</sup>	Assimilation	W	+	+	+	–
D,L-Carnitine <sup>b</sup>	Assimilation	+	W	+	+	–
Succinic Acid Mono-Methyl-Ester <sup>b</sup>	Assimilation	–	+	–	+	–
Succinamic Acid <sup>b</sup>	Assimilation	–	+	–	+	–
Glucuronamide <sup>b</sup>	Assimilation	–	+	–	+	–
Glycyl-Laspartic Acid <sup>b</sup>	Assimilation	–	+	–	W	–
i-Erythritol <sup>b</sup>	Assimilation	–	+	–	W	–
Xylitol <sup>b</sup>	Assimilation	–	W	–	+	–
Thymidine <sup>b</sup>	Assimilation	–	+	–	+	–
2-Aminoethanol <sup>b</sup>	Assimilation	–	+	–	+	W
Tween 40 <sup>b</sup>	Assimilation	–	+	+	+	+
L-Leucine <sup>b</sup>	Assimilation	–	+	W	+	W
D-Fructose <sup>b</sup>	Assimilation	–	W	+	+	+
L-Threonine <sup>b</sup>	Assimilation	–	+	W	–	–
Hydroxy-L Proline <sup>b</sup>	Assimilation	–	–	W	+	+
Uridine <sup>b</sup>	Assimilation	–	W	–	+	+
D-Sorbitol <sup>b</sup>	Assimilation	+	–	–	+	–

producing secondary metabolites, including antibiotics and fungicides, which help protect plants from infections by fungi and pathogenic bacteria.<sup>32,33</sup> In addition, it has been reported that certain subgroups of this complex can act in the bioremediation of heavy metals and pesticides.<sup>34</sup> Furthermore, several members of the complex play a vital role in the carbon cycle.<sup>30</sup> It has been suggested that the use of members of the subgroup *P. fluorescens* as plant growth-promoting rhizobacteria (PGPR) in plants of agronomical interest could reduce CO<sub>2</sub> emissions, thereby mitigating global warming and the resulting climate change.<sup>35</sup> On this background, the biotechnological potential of the three candidates for new species is obvious. Future studies are needed to analyze their possible use as plant growth-promoting rhizobacteria (PGPR), as well as their use in biocontrol and bioremediation.

It is worth noting that the four isolates were collected from different environments, including seawater, marine sediment and freshwater, suggesting that the *P. fluorescens* complex might have a notable capacity of colonization of diverse environments in the Antarctic continent. This does not seem to be uncommon in extreme environments, and a study from the Himalayas, another environment of extreme cold, also reported the *P. fluorescens* complex to be highly represented within the genus *Pseudomonas*.<sup>36</sup> Of 336 specimens of the genus *Pseudomonas*, 308 isolates belonged to this complex. Of note, the temperatures of the sampling sites ranged from 2 to 11 °C in the Himalayas, thus confirming that members of this complex are tolerant of the cold. Future studies examining a large number of Antarctic isolates are needed to determine the abundance of this complex on the Antarctic continent and to elucidate the ecological role played by these bacteria.

Due to the high volume of whole genome sequencing data and metagenomic analyses, the description of new bacterial species is turning increasingly more complex. MLSA is a robust technique to determine whether a particular isolate belongs to a previously described species, or whether it represents a new species. Several studies have shown that the study of concatenated sequences of carefully selected housekeeping genes is a powerful and reliable tool to identify new species within a genus.<sup>37,38</sup> Here, we suggest that, in addition to 16S-rRNA, the genes *aroE*, *gyrB*, *ileS*, *glnS* and *rpoD* are ideal for use in MLSA in the genus *Pseudomonas*, since they combine all necessary features: they are “housekeeping” genes, they are protein-coding, and they are not transmitted horizontally.<sup>37,39</sup> Although previous studies have suggested that the gene *ileS* is not a good candidate to be used in MLSA,<sup>40</sup> in this study, it proved to be highly efficient. On the other hand, the gene *rpoB* is commonly used in MLSA,<sup>29,41–44</sup> however, despite trying several sets of primers, we failed to obtain a unique product suitable for sequencing from our Antarctic bacterial isolates. The gene was therefore excluded from the analyses. One of the strengths of our work is that we performed MLSA on a concatenate of six genes (16S, *aroE*, *gyrS*, *glnS*, *ileS* and *rpoD*), as opposed to the three or four genes used in most studies characterizing members of the genus *Pseudomonas*. This resulted in a significantly more robust analysis. Without doubt, one of the main problems of this study lay in the shortage of available Antarctic *Pseudomonas* sequences. Sequences for all genes (*aroE*, *glnS*, *gyrB*, *ileS* and *rpoD*) were only available for the strains *P. antarctica* PAMC27494, *P. extremaustralis* 14-3 and *P. deceptionensis* DSM26521. The phylogenetic tree obtained through MLSA could therefore be further complemented when the sequences of other species become available in the database. For the 16S rRNA gene, however, sequences from all Antarctic species were available.

Molecular identification by MLSA has proven to be an effective tool to determine whether a bacterial isolate belongs to a particular species. However, in addition to phylogenetic, phenotypic or serological analyses, the most recommended method to discriminate between bacterial species, and to characterize new bacterial species, is DNA-DNA hybridization.<sup>45</sup> This method does, however, have limitations: it takes long to develop and relies on specialized expertise, it does not define the phylogenetic distance between species, and it is not cumulative.<sup>42</sup> A fast and robust method of classification for the genotypic characterization based on the sequences of protein-coding genes, MLSA has allowed researchers to classify a diverse and previously undescribed group of prokaryotes obtained from metagenomics analyses, many of which cannot be cultured.<sup>28,42</sup> It has therefore even been argued that DNA-DNA hybridization analysis may no longer be necessary for the description of a new species.<sup>42,43,46</sup>

In conclusion, our results strongly suggest that the Antarctic isolates *Pseudomonas* sp. KG01, *Pseudomonas* sp. IB20 and *Pseudomonas* sp. 12B3 are candidates for new species of the genus *Pseudomonas*, that all of them belong to the *P. fluorescens* complex, and that, together with the Antarctic strains *P. antarctica* PAMC27494 and *P. extremaustralis* 14-3, they form part of the new subgroup “*P. antarctica*”. Future studies including metagenomics studies, are needed to determine the abundance of

members of the genus *Pseudomonas* in Antarctica and to elucidate their possible ecological role on the cold continent.

## Funding

This work was supported by the Dirección de Investigación of the Pontificia Universidad Católica de Valparaíso [Project DI 037.431/2015], by an INACH PhD scholarship [DT.02-13], and by the Programa de Investigación Asociativa [PIUAs-PUCV 037.293-2015].

## Conflicts of interest

The authors declare that they have no conflict of interest.

## Acknowledgments

We would like to thank Dr. Marcelo González of the Chilean Antarctic Institute (INACH) for providing the Antarctic bacterial isolates.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.bjm.2018.02.005.

## REFERENCES

1. Cowan DA, Tow LA. Endangered Antarctic environments. *Annu Rev Microbiol.* 2004;58:649–690.
2. Smith J, Tow L, Stafford W, Cary C, Cowan D. Bacterial diversity in three different Antarctic cold desert mineral soils. *Microb Ecol.* 2006;51(4):413–421.
3. Tytgat B, Verleyen E, Obbels D, et al. Bacterial diversity assessment in Antarctic terrestrial and aquatic microbial mats: a comparison between bidirectional pyrosequencing and cultivation. *PLOS ONE.* 2014;9(6):e97564.
4. Loperena L, Soria V, Varela H, et al. Extracellular enzymes produced by microorganisms isolated from maritime Antarctica. *World J Microbiol Biotechnol.* 2012;28(5):2249–2256.
5. Cavicchioli R. Microbial ecology of Antarctic aquatic systems. *Nat Rev Microbiol.* 2015;13(11):691–706.
6. Martínez-Rosales C, Fullana N, Musto H, Castro-Sowinski S. Antarctic DNA moving forward: genomic plasticity and biotechnological potential. *FEMS Microbiol Lett.* 2012;331(1):1–9.
7. Anzai Y, Kim H, Park JY, Wakabayashi H, Oyaizu H. Phylogenetic affiliation of the pseudomonads based on 16S rRNA sequence. *Int J Syst Evol Microbiol.* 2000;50 Pt 4(4):1563–1589.
8. Rojo F. Carbon catabolite repression in *Pseudomonas*: optimizing metabolic versatility and interactions with the environment. *FEMS Microbiol Rev.* 2010;34(5):658–684.
9. Palleroni NJ. The *Pseudomonas* story. *Environ Microbiol.* 2010;12(6), 1377–1383.10.
10. Puchałka J, Oberhardt MA, Godinho M, et al. Genome-scale reconstruction and analysis of the *Pseudomonas putida* KT2440 metabolic network facilitates applications in biotechnology. *PLoS Comput Biol.* 2008;4(10):e1000210.

11. Spiers AJ, Buckling A, Rainey PB. The causes of *Pseudomonas* diversity. *Microbiology*. 2000;2345–2350.
12. Reddy GSN, Matsumoto GI, Schumann P, Stackerbrandt E, Shivaji S. Psychrophilic pseudomonads from Antarctica: *Pseudomonas antarctica* sp. nov., *Pseudomonas meridiana* sp. nov. and *Pseudomonas proteolytica* sp. nov. *Int J Syst Evol Microbiol*. 2004;54(3):713–719.
13. Bozal N, Montes MJ, Mercadé E. *Pseudomonas guineae* sp. nov., a novel psychrotolerant bacterium from an Antarctic environment. *Int J Syst Evol Microbiol*. 2007;57(11):2609–2612.
14. López N, Pettinari MJ, Stackerbrandt E, et al. *Pseudomonas extremaustralis* sp. nov., a poly(3-hydroxybutyrate) producer isolated from an Antarctic environment. *Curr Microbiol*. 2009;59(5):514–519.
15. Carrión O, Miñana-Galbis D, Montes MJ, Mercadé E. *Pseudomonas deceptionensis* sp. nov., a psychrotolerant bacterium from the Antarctic. *Int J Syst Evol Microbiol*. 2011;61(Pt 10):2401–2405.
16. Kosina M, Barták M, Mašláňová I, et al. *Pseudomonas prosekii* sp. nov., a novel psychrotrophic bacterium from Antarctica. *Curr Microbiol*. 2013;67(6):637–646.
17. Kosina M, Švec P, Černohlávková J, et al. Description of *Pseudomonas gregormendelii* sp. nov., a novel psychrotrophic bacterium from James Ross Island, Antarctica. *Curr Microbiol*. 2016;73(1):84–90.
18. Schleifer KH. Classification of bacteria and archaea: past, present and future. *Syst Appl Microbiol*. 2009;32(8):533–542.
19. Schleifer K-H, Amann R, Rosselló-Móra R. Taxonomy in the age of genomics. Introduction. *Syst Appl Microbiol*. 2015;38(4):207–208.
20. Tindall BJ, Rosselló-Móra R, Busse H-J, Ludwig W, Kämpfer P. Notes on the characterization of prokaryote strains for taxonomic purposes. *Int J Syst Evol Microbiol*. 2010;60(Pt 1):249–266.
21. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol*. 1990;215(3):403–410.
22. Winsor GL, Lam DKW, Fleming L, et al. *Pseudomonas* Genome Database: improved comparative analysis and population genomics capability for *Pseudomonas* genomes. *Nucleic Acids Res*. 2011;39(suppl 1):D596–D600.
23. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets.
24. Andreani NA, Martino ME, Fasolato L, et al. Tracking the blue: a MLST approach to characterise the *Pseudomonas fluorescens* group. *Food Microbiol*. 2014;39:116–126.
25. Pavlov MS, Lira F, Martínez JL, Olivares J, Marshall SH. Draft genome sequence of Antarctic *Pseudomonas* sp Strain KG01 with full potential for biotechnological applications. *Genome Announc*. 2015;3(4).
26. Bölter M. Ecophysiology of psychrophilic and psychrotolerant microorganisms. *Cell Mol Biol (Noisy-le-grand)*. 2004;50(5):563–573.
27. Kejna M. Air temperature on King George Island, South Shetland Islands, Antarctica. *Pol Polar Res*. 1999;20(3):183–201.
28. Gevers D, Cohan FM, Lawrence JG, et al. Opinion: re-evaluating prokaryotic species. *Nat Rev Microbiol*. 2005;3(9):733–739.
29. Garrido-Sanz D, Meier-Kolthoff JP, Göker M, et al. Genomic and genetic diversity within the *Pseudomonas fluorescens* Complex Vinatzer BA, ed. *PLOS ONE*. 2016;11(2):e0150183.
30. Remold SK, Purdy-Gibson ME, France MT, Hundley TC. *Pseudomonas putida* and *Pseudomonas fluorescens* species group recovery from human homes varies seasonally and by environment. *PLOS ONE*. 2015;10(5):e0127704.
31. Liu X, Ecartot M, Kontro MH. The physicochemical conditions of isolation source determine the occurrence of *Pseudomonas fluorescens* group species. *Ann Microbiol*. 2015;65(4):2363–2377.
32. Gross H, Loper JE. Genomics of secondary metabolite production by *Pseudomonas* spp. *Nat Prod Rep*. 2009;26(11):1408–1446.
33. Weller DM, Landa BB, Mavrodi OV, et al. Role of 2,4-diacetylphloroglucinol- producing fluorescent *Pseudomonas* spp. in the defense of plant roots. *Plant Biol*. 2007;9(1):4–20.
34. Wasi S, Tabrez S, Ahmad M. Use of *Pseudomonas* spp. for the bioremediation of environmental pollutants: a review. *Environ Monit Assess*. 2013;185(10):8147–8155.
35. Nie M, Bell C, Wallenstein MD, Pendall E. Increased plant productivity and decreased microbial respiratory C loss by plant growth-promoting rhizobacteria under elevated CO<sub>2</sub>. *Sci Rep*. 2015;5:9212.
36. Negi YK, Prabha D, Garg SK, Kumar J. Genetic diversity among cold-tolerant fluorescent *Pseudomonas* isolates from Indian Himalayas and their characterization for biocontrol and plant growth-promoting activities. *J Plant Growth Regul*. 2010;30(2):128–143.
37. Konstantinidis KT, Tiedje JM. Towards a genome-based taxonomy for prokaryotes. *J Bacteriol*. 2005;187(18):6258–6260.
38. Martens M, Dawyndt P, Coopman R, Gillis M, De Vos P, Willems A. Advantages of multilocus sequence analysis for taxonomic studies: a case study using 10 housekeeping genes in the genus *Ensifer* (including former *Sinorhizobium*). *Int J Syst Evol Microbiol*. 2008;58(Pt 1):200–214.
39. Mulet M, Lalucat J, García-Valdés E. DNA sequence-based analysis of the *Pseudomonas* species. *Environ Microbiol*. 2010;12(6):1513–1530.
40. Andreani NA, Martino ME, Fasolato L, et al. Tracking the blue: a {MLST} approach to characterise the *Pseudomonas fluorescens* group. *Food Microbiol*. 2014;39:116–126.
41. Gomila M, Peña A, Mulet M, Lalucat J, García-Valdés E. Phylogenomics and systematics in *Pseudomonas*. *Front Microbiol*. 2015;6:214.
42. Bennasar A, Mulet M, Lalucat J, García-Valdés E. Pseudo-MLSA: a database for multigenic sequence analysis of *Pseudomonas* species. *BMC Microbiol*. 2010;10:118.
43. Glaeser SP, Kämpfer P. Multilocus sequence analysis (MLSA) in prokaryotic taxonomy. *Syst Appl Microbiol*. 2015;38(4):237–245.
44. Alvarez-Pérez S, de Vega C, Herrera CM. Multilocus sequence analysis of nectar pseudomonads reveals high genetic diversity and contrasting recombination patterns. *PLOS ONE*. 2013;8(10):e75797.
45. Stackerbrandt E, Frederiksen W, Garrity GM, et al. Report of the ad-hoc committee for the re-evaluation of the species definition in bacteriology. *Int J Syst Evol Microbiol*. 2002;52(Pt 3):1043–1050.
46. Gevers D, Coenye T. Phylogenetic and genomic analysis. *Man Environ Microbiol*. 2007:157–168.