



Image of the month

Mesenteric Fibromatosis With Complete Aortic Thrombosis as an Indication for Intestinal Transplant[☆]



Fibromatosis mesentérica con trombosis completa de la aorta como indicación de trasplante intestinal

Oscar Caso-Maestro,^{*} Iago Justo-Alonso, Félix Cambra-Molero, Carlos Jiménez-Romero

Cirugía General y Trasplante de Órganos Abdominales, Hospital Universitario 12 de Octubre, Madrid, Spain




Fig. 1

We present the case of a 35-year-old male who was referred to our department due to mesenteric fibromatosis with severe infiltration at the root of the mesentery. After completing the study with an abdominal MRI and aortography, the presence of a mass was confirmed, which measured 92 mm×55 mm×73 mm and completely affected the root of the mesentery. This caused complete thrombosis of the infrarenal abdominal aorta, with hypertrophy of Riolan's arch from the middle colic artery to the inferior mesenteric artery. With these findings, we decided to study the patient as a candidate for bowel transplantation (Fig. 1).

[☆] Please cite this article as: Caso-Maestro O, Justo-Alonso I, Cambra-Molero F, Jiménez-Romero C. Fibromatosis mesentérica con trombosis completa de la aorta como indicación de trasplante intestinal. Cir Esp. 2015;93:599.

^{*} Corresponding author.

E-mail address: oscarcasomaestro@hotmail.com (O. Caso-Maestro).