



CIRUGÍA ESPAÑOLA

www.elsevier.es/cirugia



Video of the month

Laparoscopic Transduodenal Ampullectomy for Ampullary Tumor[☆]



Ampulectomía transduodenal laparoscópica por tumor ampular

Esteban Cugat Andorrà, Eric Herrero Fonollosa,^{*} Maria Isabel García Domingo, Judith Camps Lasa

Unidad de Cirugía de Hepatobiliopancreática, Hospital Universitari Mutua Terrassa, Terrassa, Barcelona, Spain

Tumors of the ampulla of Vater are rare, and their treatment usually involves pancreaticoduodenectomy (PD).

In selected cases of benign lesions, such as adenomas or carcinoma in situ, resection of the ampullary tumor could be an alternative, and PD should be carried out depending on the pathology results.

We present the case of a 78-year-old patient diagnosed with jaundice from a well-defined solid nodular tumor in the ampulla of Vater. A laparoscopic transduodenal

ampullectomy with choledochoscopy was performed. The postoperative course was correct, and the definitive pathology result was compatible with adenoma.

Appendix A. Supplementary Data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.ciresp.2018.09.012](https://doi.org/10.1016/j.ciresp.2018.09.012).

[☆] Please cite this article as: Cugat Andorrà E, Herrero Fonollosa E, García Domingo MI, Camps Lasa J. Ampulectomía transduodenal laparoscópica por tumor ampular. Cir Esp. 2019;97:50.

^{*} Corresponding author.

E-mail address: eherrero@mutuaterrassa.cat (E. Herrero Fonollosa).