



CIRUGÍA ESPAÑOLA

www.elsevier.es/cirugia


Original article

Starting a robotic thoracic surgery program: From wedge resection to sleeve lobectomy in six months. Initial conclusions[☆]



Pablo Luis Paglialunga,^{a,*} Laureano Molins,^{a,b} Rudith Guzmán,^a Angela Guirao,^{a,b} Leandro Grando,^a David Sanchez-Lorente,^a Carlos Guerrero,^a Irene Bello,^{a,b} Nestor Quiroga,^a Marc Boada^a

^a Department of Thoracic Surgery, Institut Clínic Respiratori (ICR), Hospital Clínic of Barcelona, University of Barcelona, Barcelona, Catalonia, Spain

^b Institut d'Investigacions Biomèdiques August Pi I Sunyer (IDIBAPS), Barcelona, Catalonia, Spain

ARTICLE INFO

Article history:

Received 20 October 2022

Accepted 8 April 2023

Available online 6 August 2023

Keywords:

Robot-assisted thoracic surgery (RATS)

Minimally invasive thoracic surgery (MITS)

Learning curve

A B S T R A C T

Introduction: Robot-assisted thoracic surgery (RATS) is a rapidly expanding technique. In our study, we aimed to analyze the results of the process to adopt robotic surgery in our Department of Thoracic Surgery.

Methods: This is an intention-to-treat analysis of a series of consecutive patients operated on using the RATS approach in our hospital from January 2021 to March 2022. Data were registered for patient characteristics, type of surgery, operative times, conversion rate, chest tube duration, length of hospital stay and complications.

The IBM SPSS[®] statistical software was used for the statistical analysis. A cumulative sum analysis of the operating time was performed to define the learning curve.

Results: During the study period, 51 patients underwent robotic surgery, including pulmonary and non-pulmonary interventions. In addition, 15 patients (29.4%) underwent non-pulmonary interventions: one pleural (2%), 2 diaphragmatic (3.9%), and 12 mediastinal (23.5%). Among the mediastinal surgeries, one conversion was necessary (8.3%) for a complex vascular malformation, and 11 were completed by RATS, including 7 (58.3%) thymomas, 3 (25%) pleuro-pericardial cysts, and one (8.3%) neurogenic tumor. Mean operative time was 141 min (104–178), mean chest tube duration was 0.9 days (0–2), and mean length of stay was 1.45 days (1–2).

Thirty-six patients underwent lung surgery (70.6%). The complete RATS resections (34; 94.4%) included: 3 wedge resections (11.1%), 2 segmentectomies (3.7%), 28 lobectomies (81.5%), and one sleeve lobectomy (3.7%). Mean surgery time was 194.56 min (141–247), chest tube duration was 3.92 days (1–8), and length of stay was 4.6 days (1–8). Complications occurred in 4 patients (11.1%). No 90-day mortalities were registered.

[☆] Please cite this article as: Paglialunga P, Boada M, Guzmán R, Sanchez-Lorente D, Guirao A, Bello I, et al. Inicio de un programa de cirugía torácica robótica: de resección en cuña a lobectomía con broncoplastia en seis meses. Conclusiones iniciales. Cir Esp. 2023.

* Corresponding author.

E-mail address: pablopaglialunga@gmail.com (P.L. Paglialunga).

<http://dx.doi.org/10.1016/j.cireng.2023.04.020>

2173-5077/© 2023 AEC. Published by Elsevier España, S.L.U. All rights reserved.

Conclusions: The implementation of RATS was achieved with good clinical results and operative times for all indications. A rapid learning curve was accomplished in short time. Previous VATS experience, patient selection, team training and program continuity are fundamental to successfully develop a RATS program.

© 2023 AEC. Published by Elsevier España, S.L.U. All rights reserved.

Inicio de un programa de cirugía torácica robótica: de resección en cuña a lobectomía con broncoplastia en seis meses. Conclusiones iniciales

R E S U M E N

Palabras clave:

Cirugía torácica asistida por robot
Cirugía torácica mínimamente
invasiva
Curva de aprendizaje

Introducción: La cirugía torácica asistida por robot (RATS) es una técnica en rápida expansión. Nuestro objetivo fue analizar el resultado del proceso de adopción de la cirugía robótica en nuestro Departamento de Cirugía Torácica.

Métodos: Este es un análisis por intención de tratamiento de una serie de pacientes consecutivos operados mediante el método RATS en nuestro centro desde enero de 2021 a marzo de 2022. Se registraron las características de los pacientes, tipo de cirugía, tiempos operatorios, tasa de conversión, duración del drenaje torácico, estancia hospitalaria y complicaciones. Para el análisis estadístico se utilizó el software IBM SPSS®. Se realizó un análisis de suma acumulada del tiempo de operación para definir la curva de aprendizaje.

Resultados: Durante el periodo de estudio, 51 pacientes fueron sometidos a cirugía robótica. 15 pacientes (29,4%) fueron sometidos a intervenciones no pulmonares: 1 pleural (2%), 2 diafragmáticas (3,9%) y 12 mediastínicas (23,5%). Entre las cirugías mediastínicas, fue necesaria una conversión (8,3%) por malformación vascular compleja y 11 fueron completadas por RATS, incluidos 7 (58,3%) timomas, 3 (25%) quistes pleuro-pericárdicos y 1 (8,3%) tumores neurogénicos. El tiempo operatorio medio fue de 141 minutos [104–178], la duración media del tubo torácico fue de 0,9 días [0–2] y la estancia media fue de 1,45 días [1–2].

36 pacientes tuvieron cirugías pulmonares (70,6%). Las resecciones RATS completas (34; 94,4%) incluyeron: 3 resecciones en cuña (11,1%), 2 segmentectomías (3,7%), 28 lobectomías (81,5%) y 1 lobectomía con broncoplastia (3,7%). El tiempo medio de cirugía fue de 194,56 minutos [141–247], la duración del tubo torácico fue de 3,92 días [1–8] y la estancia hospitalaria fue de 4,6 días [1–8]. No se registró mortalidad a los 90 días.

Conclusiones: La implementación de RATS se logró con buenos resultados clínicos y tiempos operatorios en todas las indicaciones. Se completó una rápida curva de aprendizaje en poco tiempo.

La experiencia previa en VATS, la selección de pacientes, la capacitación del equipo y la continuidad del programa son fundamentales para desarrollar con éxito un programa RATS.

© 2023 AEC. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

While video-assisted thoracoscopy (VATS) has several advantages compared with open surgery, as smaller incisions, less pain, shorter hospital stay, quicker recoveries and a faster return to routine daily activity, it has some limitations, including lack of articulation of the instrument, two-dimensional visualization, and counterintuitive movement of the instrument. Robot-assisted thoracic surgery (RATS) is an evolving technique which has helped overcome some of these limitations. High-definition three-dimensional stereo video, improved ergonomics and tremor suppression are some of RATS benefits.^{1–3} In addition, the articulating robotic instrument is also better at performing procedures within deep and narrow spaces. These attributes may facilitate the

resection of different elements, like hilar and mediastinal lymph nodes in robotic surgery compared with VATS.^{4–7}

Such are some of the reasons why robotic surgery is growing exponentially worldwide. Since Melfi et al. firstly applied it to thoracic surgery in 2002,⁸ several authors have demonstrated the feasibility and safety of RATS for complex thoracic procedures.^{9–13} In recent years, multiple studies have compared other minimally invasive approaches, such as VATS, with RATS. In some of them, RATS was associated with less blood loss, lower conversion rates, a higher number of harvested lymph nodes, shorter postoperative chest tube drainage and less prolonged hospital stays.^{14–19}

Despite reported benefits a slow implementation rate of RATS programs is being observed across Europe. In Spain, according to the work published by Varela et al., robot-assisted thoracic surgery is not implemented in most Thoracic Surgery

Departments. This occurs in some reason because there is a reticence to go through a learning curve while the clinical and economic benefits are still under evaluation.²⁰ Not only that, the increasing demand for RATS, requires structured and standardized training modules that will translate into efficient programs with good short-term results. Accordingly, meticulous and continuous analysis of the programs is necessary to demonstrate their effectiveness; some use learning curves to evaluate that.^{21,22}

In the present study we want to analyze and evaluate initial clinical results, surgical times and learning curves after initiating a Robotic-Assisted Thoracic Surgery Program at our center.

Methods

This is an intention to treat analysis of a series of consecutive patients operated on using Robotic assisted thoracic surgery approach in our center from January 2021 to March 2022. Da Vinci™ Xi surgical system (Intuitive, California, USA/ Abex Excellence Robotics, Spain) was used in all cases.

Demographics and patients' characteristics, including age, gender, comorbidities, and functional status in case of lung resection, were duly registered before surgery. Patients were classified into four groups according to the type of surgery. *Mediastinal surgery*, including anterior tumors (thymoma and pleuro-pericardiac cyst) and posterior tumors (neurogenic and bronchogenic cyst); *pleural surgery*, including parietal and visceral pleura; *lung surgery*: including all lung resections, anatomic and non-anatomic; Fourth group was reserved for other interventions as diaphragmatic.

In general terms, the selection criteria used do not differ from those used for VATS in our hospital. In the case of

pulmonary interventions, tumors up to clinical stage IIB were selected according to the 8th edition of the TNM. In the case of mediastinal lesions, full capsulated nodules with no image of invasion of mediastinal structures in computed tomography (CT) scan and/or magnetic resonance imaging (MRI) were selected for RATS approach. Likewise, in the other two groups, patients with lesions without involvement of central structures that could be resected by a minimally invasive approach were included. However, in order to achieve a safe program, start first cases selected for robotic surgery were small posterior mediastinal tumors, thymomas and non-anatomic lung resections which required lymphadenectomy. After that, progression to major lung resections was done starting with lower lobes previously diagnosed for non-small cell lung cancers.

Intraoperatively, surgical time, blood loss, rate of conversion, reason for conversion, final used approach (VATS or open surgery), and air leak were registered. Postoperatively, need and length of stay at intensive care unit (ICU) was registered. Complications and outcome at discharge were also registered. Day of chest tube removal and day of discharge were registered to calculate days of chest tube and total length of stay (LOS).

Lung surgery

Lung surgeries were performed in the lateral decubitus position, with lateral flexion of the thorax. All robotic ports were placed according to Fig. 1 (right) and CO2 insufflation at low pressure and flow was started (Pressure 5, Flow 5).

Specific analysis of lung interventions was done. Diagnostic for lung surgery was registered before intervention whether it was for lung metastasis, solitary lung nodule or diagnosed primary lung cancer. Thoracoscopic tunnel technique for




Fig. 1 – Left: Port placement for RATS thymectomy. 4 ports were used. One port was inserted in the third intercostal space and two in the fifth intercostal space. The assistance was placed at the level of the sixth intercostal space. Right: Port placement for RATS lung resection. All ports including assistance (A) were placed in the eighth intercostal space, except for the anterior port, which was placed in sixth intercostal space.

anatomical resections was the preferred technique for dissection. Major anatomic resections were always completed by systematic nodal dissection. Type and extension of lung surgery were recorded. Extended resections characteristics were registered when necessary. A chest tube was placed at the end of surgery and total blood loss and air leak after extubation were registered.

Mediastinal surgery

For the mediastinal approach, the patients were placed in supine decubitus position with partial elevation of the left hemithorax at 45°. Three robotic ports were placed according to Fig. 1 (left) and CO₂ insufflation at low pressure and flow was started (Pressure 5, Flow 5). For radical thymectomies resections started with a full dissection anterior to the left phrenic nerve and posterior to the mammary artery until identification of innominate vein. Thymic veins were divided either using clips or bipolar electrocoagulation. Dissection was completed all through the right phrenic nerve.

Conversion

We consider conversion when the surgery started by RATS but ended using another approach (VATS or open surgery) for any reason: emergent, technical or oncologic. The final surgery and outcome were registered and included in our database regardless.

Surgical time

Surgical time is the time span from skin opening to skin closure. It includes port placement, robot docking and targeting, console time, specimen retrieval and skin closure.

Statistical analysis

Results are presented as n (proportion), mean $\pm$ standard deviation or median [interquartile range] as appropriate. The IBM SPSS® statistical software was used for statistical analysis. For the analysis of the learning curves measuring the surgical time, in lobectomies and thymectomies we resorted to the CUSUM (Cumulative Sum) graphs, which are widely used for quality control and best adapt to the monitoring of clinical-care processes.^{21,22}

Results

During the study period, 51 patients underwent thoracic robotic surgery in our department. Demographics and characteristics are summarized in Table 1. The surgeries performed were: 15 (29.4%) non-pulmonary interventions including 1 pleural (2%), 2 diaphragmatic (3.9%) and 12 mediastinal (23.5%). 36 patients had pulmonary surgery (70.6%).

Lung surgery

36 patients underwent lung surgery with a median age of 65.56 (SD 11.2) and a gender distribution of 19 (52.7%) females and 17 (47.2%) males. Functional status are summarized in Table 2.

Table 1 – All RATS patients (n = 51) demographics and characteristics.

Age; mean (Std)	62.08 (12.4)
Gender	
Male; n (%)	25 (49)
Female; n (%)	26 (51)
Comorbidity	32 (62.7)
Chronic Obstructive Pulmonary Disease; n (%)	8 (15.7)
Hypertension; n (%)	22 (43.1)
Diabetes; n (%)	9 (17.6)
Chronic renal failure; n (%)	4 (7.8)
Cardiovascular; n (%)	12 (23.5)
Anticoagulation-agretation; n (%)	9 (17.6)
Surgery	
Lung; n (%)	36 (70.6)
Pleura; n (%)	1 (2)
Mediastinum; n (%)	12 (23.5)
Other; n (%)	2 (3.9)

Completed RATS resections (34; 94.4%) included: 3 wedge resections (11.1%), 2 segmentectomies (3.7%), 28 lobectomies (81.5%), and 1 sleeve lobectomy (3.7%). Conversion to thoracotomy was necessary in two cases (5.6%) for oncological reasons; 1 patient had chest wall invasions, and the other presented hilar lymph node with main pulmonary artery infiltration requiring a pneumonectomy.

After excluding wedge resections and conversion surgeries, 31 anatomic lung resections were performed by RATS during the study period. Type of surgery and distribution are summarized in Table 3. Mean surgical time for those patients who completed RATS lobectomy was 212.42 min [range 137–317], mean chest tube duration was 3.79 days [range 1–9] and mean length of stay was 4.46 days [range 2–9]. 12 patients (38.7%) spent the immediate postoperative period in the ICU. Their average stay at ICU was 1 day. When upper and non-upper lobectomies surgical times were compared no differences were detected (208 ± 56 vs 187 ± 45 min respectively; vs $P = .2$). 48% of patients (15) had a preoperative diagnostic and 16 patients (52%) needed intraoperative diagnostic; however, it did not affect surgical time (204 ± 61 min vs 195 ± 43 min, $P = .6$). Complications occurred in 4 patients (11.1%), 3 cases of air leak were registered and treated conservatively, and one patient required VATS re-interventions for bleeding. No mortality was registered at 90 days. Readmission was necessary in 3 cases (12.5%), 1 because of gastric bleeding, 1 for pain management and 1 because of wound infection.

Learning curve

The data of the CUSUM curve for the robotic surgery time of the lobectomies were adjusted to a polynomial of order 2 (Fig. 2). In our series, the learning curve was completed with 23 procedures.

Mediastinal surgery

12 (23.5%) patients underwent mediastinal surgery with a median age of 52 (SD 11.4) and a gender distribution of 6 (50%) females and 6 (50%) males. 11 surgeries were completed by RATS. Conversion was necessary in one case (8.3%) because

Table 2 – Preoperative pulmonary function tests for RATS lung resections (n = 36).

FEV1; L (SD)	2.39 (0.6)
FEV1; % (SD)	85.97 (18.01)
FVC; L (SD)	3.41 (0.8)
FVC; % (SD)	91.80 (15.7)
DLCO; % (SD)	82.17(19.4)
FEV1: Forced expiratory volume; FVC: Forced vital capacity; DLCO: Diffusion capacity of lung for carbon monoxide.	

pericardial, left lung and innominate vein involment of a complex vascular malformation of the mediastinum.

Among RATS completed surgeries the diagnoses were, 7 (58.3%) were thymomas, 3 (25%) pleuro-pericardiac cysts and 1 (8.3%) neurogenic tumor. Mean time was 141 min [104–178], mean chest tube duration was 0.9 days [0–2] and mean length of stay was 1.45 days.^{1,2} No complications were registered.

Learning curve for thymectomies

The data of the CUSUM curve for the time of robotic surgery of the thymectomies were adjusted to a polynomial of order 5 (Fig. 3).

Discussion

Within 15 months, we performed 51 surgeries, achieving an increasing level of complexity with no impact on surgical times and a low percentage of complications and conversions. Anatomical lung resections were the most common procedures with 31 surgeries, allowing us to consolidate our learning curve after 23 cases. We have also performed surgeries for mediastinal, pleural, and diaphragmatic pathologies with no complications and only one conversion.

The GEVATS group (Spanish Group of Video-Assisted Thoracic Surgery), analyzed the Spanish national database²³ and reported a rate of morbidity, mortality and conversions comparable to our first cases of RATS. When evaluating our postoperative hospital stay in pulmonary resections, it averaged 4.6 days (SD 2–9) compared to 5 days (SD 4–7) in the GEVATS series published in 2021.²⁴ In the other hand, our results are far from more experienced groups. Travis C. Geraci et al. published, a 53% rate of discharge at postoperative day one from a total of 253 robotic anatomic pulmonary resections.²⁵

Regarding the defined learning curve for RATS lobectomies, we evidenced a learning curve completed at case 23 based on

Table 3 – Patients' characteristics.

Patient	Gender (years old)	Smoker	Preoperative diagnosis	Location	Surgery	Surgery time (min)	Postoperative diagnosis	Complication	LOS
1	F; 73	Never	Yes	LLL	Lobectomy	170	ADK (pT2aN0)	No	5
2	F; 64	Active	No	LLL	Lobectomy	230	ADK (pT1bN2)	No	4
3	F; 71	Never	Yes	RUL	Lobectomy	274	ADK (pT1bN0)	PAL	9
4	M; 72	Former	No	LLL	Lobectomy	270	ADK (pT1bN0)	No	3
5	F; 83	Never	Yes	LLL	Lobectomy	225	ADK (pT2aN1)	No	3
6	M; 75	Active	No	LUL	Lobectomy	204	ADK (pT2aN0)	No	2
7	M; 75	Unknown	No	RUL	Lobectomy	180	ADK (pT1cN0)	No	3
8	F; 66	Active	Yes	LUL	Lobectomy	298	SCLC (pT1cN0)	No	3
9	M; 62	Former	Yes	RUL	Lobectomy	204	ADK (pT2aN0)	No	4
10	F; 58	Never	Yes	RLL	Lobectomy	184	Carcinoid (pT1bN0)	No	6
11	M; 79	Active	No	LUL	Lobectomy	279	ADK (pT1bN0)	Yes (Bleeding)	7
12	M; 46	Active	Yes	RUL	Lobectomy	240	Non-malignant	No	5
13	F; 47	Never	Yes	RLL	Lobectomy	165	Atypical carcinoid (T3N0)	No	2
14	M; 51	Active	Yes	RUL	Lobectomy	196	ADK (pT2aN0)	No	2
15	F; 70	Never	No	RML	Lobectomy	174	ADK (pT1bN0)	No	8
16	M; 67	Former	Yes	RLL	Lobectomy	253	ADK (pT1bN0)	PAL	9
17	M; 54	Never	No	RML	Lobectomy	144	Non-malignant	No	3
18	F; 71	Former	Yes	LLL	Lobectomy	137	SQM (pT2bN0)	No	3
19	M; 63	Active	Yes	LUL	Sleeve	317	NSCLC (pT2aN1)	No	4
20	F; 73	Active	No	RLL	Lobectomy	205	ADK (pT1bN0)	PAL	7
21	F; 54	Former	No	RUL	Lobectomy	205	ADK (pT1cN0)	No	6
22	F; 67	Never	No	RUL	Lobectomy	173	ADK (pT1bN0)	No	2
23	F; 61	Former	No	LUL	Lobectomy	227	Non malignant	No	5
24	F; 62	Active	No	RLL	Segmentectomy	144	SQM (pT1bN0)	No	2
25	F; 42	Active	Yes	RLL	Lobectomy	133	ADK (pT1cN0)	No	3
26	F; 79	Active	No	RUL	Lobectomy	164	ADK (pT1bN0)	No	3
27	F; 81	Active	No	RUL	Segmentectomy	132	ADK (pT1bN0)	No	3
28	M; 63	Former	Yes	RUL	Lobectomy	128	SQM (pT1cN0)	No	2
29	M; 59	Active	No	RUL	Lobectomy	230	SQM (pT2aN0)	PAL	7
30	F; 54	Never	No	LUL	Lobectomy	162	ADK (pT1bN0)	AF	4
31	M; 72	Active	Yes	RUL	Lobectomy	136	ADK pT1cN0	No	3

F: female; M: male; RUL: right upper lobectomy; RML: right middle lobectomy; RLL: right lower lobectomy; LUL: left upper lobectomy; LLL: left lower lobectomy; ADK: adenocarcinoma; SCLC: small cell lung cancer; SQM: squamous cell carcinoma; PAL: Persistent air leak; AF: atrial fibrillation; LOS: length of stay.




Fig. 2 – CUSUM curve, for the robotic surgery time of the lobectomies.




Fig. 3 – CUSUM curve, for the robotic surgery time of the thymectomy.

surgical time. Previous articles determined that the learning curve for anatomical lung resections is completed after 14–34 cases when analyzing the surgical time.^{21,26–28} However, the learning curve can also be measured using other parameters. Meyer et al. used mortality and surgeon comfort, and set the learning curve at 20, and 19 cases, respectively.²⁷

Most of the procedures we have performed are pulmonary resections (70.6%). From our experience with this type of procedure, we can extract three conclusions worth taking into account when starting a robotics program.

First, previous experience in minimal invasive surgery might play an important role. At the time of starting the program we had a rate of VATS near to 80% of all anatomical resections. In this direction Merritt et al. suggested that, concepts from previous VATS experience might be transferred to RATS and help to reduce surgical times in a faster and safer way.^{9,29} For example, our previous experience using the tunnel technique for anatomical lung resections, described by Decaluwe³⁰ on VATS resections permitted us to use it on RATS.³¹ Tunnel technique requires less parenchymal manipulation and reduces accidental tears happening during the first RATS cases.

Second, selection of patients is crucial to improve results. We progressively increased complexity starting with small non pulmonary tumors, continuing to thymomas and non anatomical lung resections to end up carrying out anatomical lung resections. When starting anatomical lung resections we find recommendable to avoid large (>5 cm tumors) and central (in direct contact with central structures) tumors. In contraposition, selection of previously diagnosed peripheral tumors allowed us to complete a safe and steady learning curve minimizing intraoperative time and reducing loss of CO₂ as a result of the retrieval of the specimen for diagnosis. We also preferred non-upper lobectomies for the first cases because they are less technically demanding and unexpected vascular injuries are easier to repair. Lower lobectomies required less surgical time in our series, although this data is not statistically significant.

Third, training is essential to achieve good results. Field and console surgeons trained in animals and cadaveric models before clinical application. Virtual simulation was also used adding advantage when implementing this new technique.^{17,32–34} As stated by Shahim et al., we used dual

console and the guidance of a mentor for the clinical implementation of the program. It is also important to achieve consistency and regularity in the number of surgeries to maintain high standards within the surgical team.^{33,35,36} In fact, something we experienced is that, by performing weekly sessions of robotic surgery with a close-trained team, within six months we were able to advance from minor resections (Wedge) to left lower sleeve lobectomy.

Taking a glimpse to the horizon we are convinced that new tools such as three-dimensional computed tomography reconstruction for operative planning in anatomical segmentectomies,³⁷ novel variations on the current approach (e.g., the use of a single work port³⁸) and the development of new surgical robotic systems,³⁹ announce an encouraging future for this technique and can contribute to improving results and reducing costs in the coming years.⁴⁰

Limitations

Beyond what has been described in this paper we have to point out some limitations. This is a non-randomized single center series of consecutive cases and the results and conclusions found in our study may not be completely applicable in other institutions. Furthermore, selection bias makes comparison with GEVATS series not feasible. However, as we stated before, selection is one of the cornerstones for a successful RATS program take off and GEVATS results can be a goal.

Conclusions

In our experience after starting RATS program, we achieved good clinical results and operative times in all mediastinal, pleural, pulmonary and diaphragmatic surgeries. In addition, we observed a fast learning curve that led to greater complexity in a short time for anatomical lung resections.

In our opinion, prior VATS experience, patient selection and team training have been essential to achieve these results and to successfully develop a RATS program in thoracic surgery.

Financial support

This research has not received specific aid from public sector agencies, the commercial sector or non-profit entities.

Conflict of interest

None.

REFERENCES

- Farivar AS, Cerfolio RJ, Vallières E, Knight AW, Bryant A, Lingala V, et al. Comparing robotic lung resection with thoracotomy and video-assisted thoracoscopic surgery cases entered into the society of thoracic surgeons database. *Innov: Technol Tech Cardiothorac Vasc Surg*. 2014;9:10-5. <http://dx.doi.org/10.1097/imi.0000000000000043>.
- Veronesi G, Novellis P, Voulaz E, Alloisio M. Robot-assisted surgery for lung cancer: state of the art and perspectives. *Lung Cancer*. 2016;101:28-34. <http://dx.doi.org/10.1016/j.lungcan.2016.09.004>.
- Novellis P, Alloisio M, Vanni E, Bottoni E, Cariboni U, Veronesi G. Robotic lung cancer surgery: review of experience and costs. *J Vis Surg*. 2017;3:39. <http://dx.doi.org/10.21037/jovs.2017.03.05>.
- Soliman BG, Nguyen DT, Chan EY, Chihara RK, Meisenbach LM, Graviss EA, et al. Impact of da Vinci Xi robot in pulmonary resection. *J Thorac Dis*. 2020;12:3561-72. <http://dx.doi.org/10.21037/jtd-20-720>.
- Novellis P, Bottoni E, Voulaz E, Cariboni U, Testori A, Bertolaccini L, et al. Robotic surgery, video-assisted thoracic surgery, and open surgery for early stage lung cancer: comparison of costs and outcomes at a single institute. *J Thorac Dis*. 2018;10:790-8. <http://dx.doi.org/10.21037/jtd.2018.01.123>.
- Zirafa CC, Romano G, Nesti A, Davini F, Melfi F. Nodal upstaging robotic lobectomy for non-small cell lung cancer. *Mini-Invasive Surg*. 2020;2020. <http://dx.doi.org/10.20517/2574-1225.2019.35>.
- Cerfolio RJ, Bryant AS, Minnich DJ. Complete thoracic mediastinal lymphadenectomy leads to a higher rate of pathologically proven N2 disease in patients with non-small cell lung cancer. *Ann Thorac Surg*. 2012;94:902-6. <http://dx.doi.org/10.1016/j.athoracsur.2012.05.034>.
- Melfi FMA, Menconi GF, Mariani AM, Angeletti CA. Early experience with robotic technology for thoracoscopic surgery. *Eur J Cardio-Thorac Surg*. 2002;21:864-8. [http://dx.doi.org/10.1016/S1010-7940\(02\)00102-1](http://dx.doi.org/10.1016/S1010-7940(02)00102-1).
- Merritt RE, Kneuert PJ, D'Souza DM. Successful transition to robotic-assisted lobectomy with previous proficiency in thoracoscopic lobectomy. *Innov: Technol Tech Cardiothorac Vasc Surg*. 2019;14:263-71. <http://dx.doi.org/10.1177/1556984519845672>.
- Veronesi G, Park B, Cerfolio R, Dylewski M, Toker A, Fontaine JP, et al. Robotic resection of stage III lung cancer: an international retrospective study. *Eur J Cardio-Thorac Surg*. 2018;54:912-9. <http://dx.doi.org/10.1093/ejcts/ezy166>.
- Ruffini E, Filosso PL, Guerrero F, Lausi P, Lyberis P, Oliaro A. Optimal surgical approach to thymic malignancies: new trends challenging old dogmas. *Lung Cancer*. 2018;118:161-70. <http://dx.doi.org/10.1016/j.lungcan.2018.01.025>.
- Zhang Y, Chen C, Hu J, Han Y, Huang M, Xiang J, et al. Early outcomes of robotic versus thoracoscopic segmentectomy for early-stage lung cancer: a multi-institutional propensity score-matched analysis. *J Thorac Cardiovasc Surg*. 2020;160:1363-72. <http://dx.doi.org/10.1016/j.jtcvs.2019.12.112>.
- Kneuert PJ, D'Souza DM, Moffatt-Bruce SD, Merritt RE. Robotic lobectomy has the greatest benefit in patients with marginal pulmonary function. *J Cardiothorac Surg*. 2018;13. <http://dx.doi.org/10.1186/s13019-018-0748-z>.
- Ma J, Li X, Zhao S, Wang J, Zhang W, Sun G. Robot-assisted thoracic surgery versus video-assisted thoracic surgery for lung lobectomy or segmentectomy in patients with non-small cell lung cancer: a meta-analysis. *BMC Cancer*. 2021;21. <http://dx.doi.org/10.1186/s12885-021-08241-5>.
- Kneuert PJ, D'Souza DM, Richardson M, Abdel-Rasoul M, Moffatt-Bruce SD, Merritt RE. Long-term oncologic outcomes after robotic lobectomy for early-stage non-small-cell lung cancer versus video-assisted thoracoscopic and open thoracotomy approach. *Clin Lung Cancer*. 2020;21:214-224.e2. <http://dx.doi.org/10.1016/j.clcl.2019.10.004>.
- Haruki T, Kubouchi Y, Takagi Y, Kidokoro Y, Matsui S, Nakanishi A, et al. Comparison of medium-term survival

- outcomes between robot-assisted thoracoscopic surgery and video-assisted thoracoscopic surgery in treating primary lung cancer. *Gen Thorac Cardiovasc Surg*. 2020. <http://dx.doi.org/10.1007/s11748-020-01312-7>.
17. Veluswamy RR, Whittaker Brown SA, Mhango G, Sigel K, Nicastri DG, Smith CB, et al. Comparative effectiveness of robotic-assisted surgery for resectable lung cancer in older patients. *Chest*. 2020;157:1313-21. <http://dx.doi.org/10.1016/j.chest.2019.09.017>.
 18. Li C, Hu Y, Huang J, Li J, Jiang L, Lin H, et al. Comparison of robotic-assisted lobectomy with video-assisted thoracic surgery for stage IIB-IIIa non-small cell lung cancer. *Transl Lung Cancer Res*. 2019;8:820-8. <http://dx.doi.org/10.21037/tlcr.2019.10.15>.
 19. Veronesi G, Abbas AES, Muriana P, Lembo R, Bottoni E, Perroni G, et al. Perioperative outcome of robotic approach versus manual videothoracoscopic major resection in patients affected by early lung cancer: results of a randomized multicentric study (ROMAN Study). *Front Oncol*. 2021;11. <http://dx.doi.org/10.3389/fonc.2021.726408>.
 20. Varela G, Hernando-Trancho F, Rodríguez Suárez PM, Jarabo Sarceda JR, Molins L, Azcárate L. Thoracic surgery in Spain. *J Thorac Dis*. 2022;14:779-87. <http://dx.doi.org/10.21037/jtd.21-1108>.
 21. Arnold BN, Thomas DC, Bhatnagar V, Blasberg JD, Wang Z, Boffa DJ, et al. Defining the learning curve in robot-assisted thoracoscopic lobectomy. *Surgery (United States)*. 2019;165:450-4. <http://dx.doi.org/10.1016/j.surg.2018.06.011>.
 22. Power AD, D'Souza DM, Moffatt-Bruce SD, Merritt RE, Kneuert PJ. Defining the learning curve of robotic thoracic surgery: what does it take? *Surg Endosc*. 2019;33:3880-8. <http://dx.doi.org/10.1007/s00464-019-07035-y>.
 23. Embun R, Royo-Crespo I, Recuero Díaz JL, Bolufer S, Call S, Congregado M, et al. Spanish video-assisted thoracic surgery group: method, auditing, and initial results from a national prospective cohort of patients receiving anatomical lung resections. *Arch Bronconeumol*. 2020;56:718-24. <http://dx.doi.org/10.1016/j.arbres.2020.01.005>.
 24. Gómez Hernández MT, Novoa Valentín NM, Embún Flor R, Varela Simó G, Jiménez López MF. Predictive factors of prolonged postoperative length of stay after anatomic pulmonary resection. *Cir Esp*. 2021. <http://dx.doi.org/10.1016/j.ciresp.2021.09.010>.
 25. Geraci TC, Chang SH, Chen S, Ferrari-Light D, Cerfolio RJ. Discharging patients by postoperative day one after robotic anatomic pulmonary resection. *Ann Thorac Surg*. 2022;114:234-40. <http://dx.doi.org/10.1016/j.athoracsur.2021.06.088>.
 26. Song G, Sun X, Miao S, Li S, Zhao Y, Xuan Y, et al. Learning curve for robot-assisted lobectomy of lung cancer. *J Thorac Dis*. 2019;11:2431-7. <http://dx.doi.org/10.21037/jtd.2019.05.71>.
 27. Meyer M, Gharagozloo F, Tempesta B, Margolis M, Strother E, Christenson D. The learning curve of robotic lobectomy. *Int J Med Robot Comp Assisted Surg*. 2012;8:448-52. <http://dx.doi.org/10.1002/RCS.1455>.
 28. Toker A, Özyurtkan MO, Kaba E, Ayalp K, Demirhan Ö, Uyumaz E. Robotic anatomic lung resections: the initial experience and description of learning in 102 cases. *Surg Endosc*. 2016;30:676-83. <http://dx.doi.org/10.1007/S00464-015-4259-X>.
 29. Sandri A, Filosso PL, Lausi PO, Ruffini E, Oliaro A. VATS lobectomy program: the trainee perspective. *J Thorac Dis*. 2016;8:S427-30. <http://dx.doi.org/10.21037/jtd.2016.03.82>.
 30. Decaluwe H, Sokolow Y, Deryck F, Stanzi A, Depypere L, Moons J, et al. Thoracoscopic tunnel technique for anatomical lung resections: a "fissure first, hilum last" approach with staplers in the fissureless patient. *Interact Cardiovasc Thorac Surg*. 2015;21:2-7. <http://dx.doi.org/10.1093/ICVTS/IVV048>.
 31. Quiroga Nestor, Boada Marc, Guzman Rudith, Paglialunga Pablo, Grando Leandro, Molins Laureano. Tunnel technique for robotic-assisted left upper lobectomy. *Multimed Manual Cardio-Thorac Surg*. 2022. <http://dx.doi.org/10.1510/mmcts.2022.003>.
 32. Costello DM, Huntington I, Burke G, Farrugia B, O'Connor AJ, Costello AJ, et al. A review of simulation training and new 3D computer-generated synthetic organs for robotic surgery education. *J Robot Surg*. 2022;16:749-63. <http://dx.doi.org/10.1007/s11701-021-01302-8>.
 33. Shahin GMM, Bruinsma GJBB, Stamenkovic S, Cuesta MA. Training in robotic thoracic surgery—the European way. *Ann Cardiothorac Surg*. 2019;8:202-9. <http://dx.doi.org/10.21037/ACS.2018.11.06>.
 34. Haruki T, Nakamura H. Surgical simulation in robot-assisted thoracoscopic surgery: future strategy. *Video Assist Thorac Surg*. 2018;3:44. <http://dx.doi.org/10.21037/vats.2018.10.01>.
 35. Chen S, Geraci TC, Cerfolio RJ. Techniques for lung surgery: a review of robotic lobectomy. *Expert Rev Respir Med*. 2018;12:315-22. <http://dx.doi.org/10.1080/17476348.2018.1448270>.
 36. Hu X, Wang M. Efficacy and safety of robot-assisted thoracic surgery (RATS) compare with video-assisted thoracoscopic surgery (VATS) for lung lobectomy in patients with non-small cell lung cancer. *Comb Chem High Throughput Screen*. 2019;22:169-78. <http://dx.doi.org/10.2174/1386207322666190411113040>.
 37. le Moal J, Peillon C, Dacher JN, Baste JM. Three-dimensional computed tomography reconstruction for operative planning in robotic segmentectomy: a pilot study. *J Thorac Dis*. 2018;10:196-201. <http://dx.doi.org/10.21037/jtd.2017.11.144>.
 38. Gonzalez-Rivas D, Manolache V, Bosinceanu ML, Gallego-Poveda J, Garcia-Perez A, de la Torre M, Turna A, Motas N. Uniportal pure robotic-assisted thoracic surgery—technical aspects, tips and tricks. *Ann Transl Med*. 2022. <http://dx.doi.org/10.21037/atm-22-1866>.
 39. Cepolina F, Razzoli RP. An introductory review of robotically assisted surgical systems. *Int J Med Robot Comp Assisted Surg*. 2022. <http://dx.doi.org/10.1002/RCS.2409>.
 40. Singer E, Kneuert PJ, D'Souza DM, Moffatt-Bruce SD, Merritt RE. Understanding the financial cost of robotic lobectomy: calculating the value of innovation? *Ann Cardiothorac Surg*. 2019;8:194-201. <http://dx.doi.org/10.21037/ACS.2018.05.18>.