



Images in Urology

Giant parastomal hernia in a patient with ileal urinary diversion

Hernia paraestomal gigante en paciente con derivación urinaria ileal



Figure 1 – Giant paraestomal hernia in 68-year-old male with ileal urinary diversion.



Figure 2 – Hernia reduced to the abdominal cavity. An aspiration drain is left in the hernia sac and withdrawn on the fifth postoperative day.

A. I. Gómez-Sotelo*, D. Rodríguez-Morillas, E. García-Abril and J. A. Pérez-Ramón

Department of General and Digestive Tract Surgery, Hospital Universitario San Cecilio, Granada, Spain

*Author for correspondence.

E-mail: anaisabelgomezsotelo@gmail.com (A. I. Gómez-Sotelo).

Squamous cell carcinoma of the penis after circumcision

Carcinoma epidermoide balanoprepucial de nueva aparición tras circuncisión



Figures 1 and 2 – At 4 months, a tumor is visible at the level of the balanopreputial sulcus on the right side, not previously present. After wide resection of the tumor, pathology shows a well-differentiated keratinizing squamous cell carcinoma (1.7 cm) infiltrating the subepithelial connective tissue.

J. D. Jiménez Parra*, J. L. Cebrián Lostal, S. Álvarez Bandrés, D. García García and F. Lozano Uruñuela

Department of Urology, Hospital Virgen del Camino, Pamplona, Spain

*Author for correspondence.

E-mail: jdjimenez@hotmail.es (J. D. Jiménez Parra).