


Clear cells carcinoma relapse in surgical incision

Recidiva de carcinoma de células con trombo en cava a nivel de incisión



Figures 1 and 2 – A 60-year-old patient was diagnosed with a right kidney tumor with a thrombus in the vena cava; the patient underwent a radical nephrectomy and thrombus removal with extracorporeal circulation. One year and three months later, the follow-up CT revealed subcutaneous tumoral implants in the right external oblique muscle (fig. 1) and uptake with PET (fig. 2).

Javier Barba Abad^{a,*}, Egoitz Tolosa Eizaguirre^a, Anibal Rincón Mayans^a, Luis Romero Vargas^a, Isabel Vivas Pérez^b and Jose María Berián Polo^a

^aDepartamento de Urología, Clínica Universidad de Navarra, Navarra, Spain

^bServicio de Radiodiagnóstico, Clínica Universidad de Navarra, Navarra, Spain

*Corresponding author.

E-mail: javiferbar@hotmail.com (J. Barba Abad).

Hard-boiled egg: a testicle tumor with burning sensation

Huevo duro: tumor de testículo quemándose



Figure 1 – Left radical orchiectomy piece with testicular tumor consisting of a yellowish nodule that microscopically is an ischemic necrotic area with neoplastic remains consistent with seminoma-type germ cell tumor (process of progression of “burnt-out” testicular tumor).

Angel Maximino Castro-Iglesias^{a,*}, Antonio Pazos-Riveiro^b and Antonio Selas-Perez^c

^aServicio de Urología, Hospital Xeral de Vigo, Galicia, Spain

^bServicio de Urología, Hospital Miguel Domínguez de Pontevedra, Galicia, Spain

^cServicio de Urología, Hospital Montecelo de Pontevedra, Galicia, Spain

*Corresponding author.

E-mail: amaxicas@hotmail.com (A.M. Castro-Iglesias).