

Does resistance training improve body image satisfaction among the elderly? A cross-sectional study

Roberta Luksevicius Rica,^I Eliane Florencio Gama,^I Alexandre Fernandes Machado,^I Angélica Castilho Alonso,^I Alexandre L. Evangelista,^{II} Aylton Figueira-Junior,^I Marcelo Zanetti,^I Regina Brandão,^I Maria Luiza de Jesus Miranda,^I José Vilaça Alves,^{III,IV} Marco Bergamin,^V Danilo Sales Bocalini^{VI,*}

^IDepartamento de Educação Física e Ciências do Envelhecimento, Laboratório de Percepção Corporal e Movimento, Universidade São Judas Tadeu, São Paulo, SP, BR. ^{II}Divisão de Educação Física, Departamento de Educação, Universidade Nove de Julho, São Paulo, SP, BR. ^{III}Sport Sciences Department, University of Trás-os-Montes e Alto Douro, Vila Real, Portugal. ^{IV}Research Center in Sports Sciences, Health Sciences and Human Development, Vila Real, Portugal. ^VSport and Exercise Medicine Division, Department of Medicine, University of Padova, Padova, Italy. ^{VI}Laboratório de Fisiologia e Bioquímica Experimental, Centro de Educação Física e Deportes, Universidade Federal do Espírito Santo (UFES), Vitória, ES, BR.

Rica RL, Gama EF, Machado AF, Alonso AC, Evangelista AL, Figueira-Junior A, et al. Does resistance training improve body image satisfaction among the elderly? A cross-sectional study. Clinics. 2018;73:e290

*Corresponding author. E-mail: bocaliniht@hotmail.com

OBJECTIVE: This study aims to evaluate whether body image satisfaction improves with practice in resistance training in elderly women.

METHODS: Forty women were selected and randomly divided into an untrained group (UN) group and a group trained in resistance exercises (RT). To evaluate body image satisfaction, the silhouette matching task was used.

RESULTS: No differences were found between current (5.45 ± 0.24) and ideal (4.7 ± 0.12) silhouettes in the RT group. However, the UN group showed a significant difference ($p < 0.01$) between current (10.4 ± 0.43) and ideal (5.6 ± 0.26) silhouettes. The current silhouette was significantly different between the UN and RT groups ($p < 0.002$). However, the ideal silhouette value did not differ between them. Body satisfaction was present in 90% of the participants in the RT group, and the difference was 15% less than that in the UN group (χ^2 test, $p < 0.001$).

CONCLUSION: Resistance training in elderly women can promote satisfaction with their body image.

KEYWORDS: Body Image; Aging; Resistance Training; Older Women.

INTRODUCTION

Paul Schilder addressed the concept of body image (BI) in 1935, noting that every person has an image of him/herself that comprises biological, sociological and libidinal reality (1). This portrait is how we see our bodies, including information arising from touch, smell, nerves, muscles, viscera and thermal and pain sensations. Body perception is based on BI and body schema. Body schema is the representation of different body segments in space, which is essential to guide movement, and it is unconscious. As a conscious visual representation of the body, BI can be influenced by the environment, other people, desire and emotions (2,3).

In elderly people, the relationship between body image satisfaction and aging remains unclear (4). Considering that older subjects are vulnerable to negative experiences,

an undesirable BI can be harmful for the aging process (5,6). Therefore, it is important to identify interventions that can improve BI in elderly people.

Physical exercise has been considered a key element of health promotion and improvement of overall physical fitness as well as an approach to increase body self-knowledge, self-esteem, self-perception and effectiveness, which are important factors for a positive BI (7-10).

The American College of Sports Medicine (11) recommends a regular physical exercise program for adults, including elderly people, with resistance exercises to improve muscle strength and functional capacity. However, there is little evidence (12) for a positive role of resistance training in BI; therefore, to fill this knowledge gap, this study aimed to evaluate whether resistance training (RT) can promote body satisfaction among elderly women.

MATERIAL AND METHODS

Subjects

Fifty volunteers (women over 60 years old) were recruited for this study. All participants had undergone medical screening and completed questionnaires regarding their medical history. The inclusion criteria were age > 60 years old and no use of hormone replacement therapy, drugs and vitamins that could alter calcium or bone metabolism. Participants were

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

No potential conflict of interest was reported.

Received for publication on August 1, 2017. Accepted for publication on February 15, 2018

DOI: 10.6061/clinics/2018/e290



excluded from participation in the study if they had any of the following conditions: participation in current or previous regular RT in the last 6 months (only for the control group), recent hospitalization, symptomatic cardiorespiratory disease, hypertension or metabolic disorders, severe renal or hepatic disease, cognitive impairment or progressive and debilitating conditions, marked obesity with inability to exercise, and recent bone fractures or any other medical contraindications to RT practice (for any group). According to the exclusion criteria, 10 subjects were excluded, and the remaining subjects were randomized to the untrained (UN; $n=20$) or resistance trained (RT; $n=20$) group.

Physical exercise program approach

The program used, cited in a previous publication (6), was performed for six months and consisted of three one-hour sessions of resistance exercises on nonconsecutive days. The following resistance exercises were included: leg press, chest press, leg curl, latissimus pull down, elbow flexion, elbow extension, leg extension, upper back row, military press, hip abductor, hip adductor, and abdominal curls. All exercises were performed with RT-specific devices. The load and repetitions were three sets of 10 at 75% of 1RM for all exercises. Participants alternated exercises for the upper and lower muscle groups to minimize fatigue, with one minute of rest between them. For all exercises, eccentric muscle action was emphasized. All sessions were guided by trained fitness instructors and supervised by the researchers.

Anthropometric parameters

Height (meters) was evaluated with a stadiometer (Cardiomed, WCS model; Paraná, Brazil) fixed to the wall and calibrated at a 0.1 cm scale. Body weight (kg) was assessed using an electronic scale (Filizzola Personal Line model 150; São Paulo, Brazil) accurate to 100 g. Body mass index (BMI) was calculated as follows: $BMI = \text{weight} / (\text{height})^2$.

Functional strength tests

Senior tests were applied according to the validated recommendations from Rikli and Jones (13). The arm curl test and chair stand test were used to assess upper and lower functional strength. Each subject completed two familiarization trials before the test. The scores were the total number of curls through the full range of motion (arm curl test) and the

total number of stands (chair stand test) executed correctly in 30 seconds.

Body image satisfaction evaluation

The instrument used to evaluate BI satisfaction was the silhouette matching task proposed by Stunkard, Sorenson and Schulsinger (14) and adapted by Marsh and Roche (15). This task consists of 12 silhouettes on a progressive scale (Figure 1). The discrepancy between the current (CS) and ideal silhouettes (IS) was analyzed after the figures were presented to the elderly women. The following questions were directed to the women: (1) What is the silhouette that best represents your current physical appearance? (2) What silhouette would you like to have?

Statistical analyses

All analyses were performed using SPSS for Windows software (version 22 SPSS Inc., Chicago, Illinois, USA). All data are expressed as the mean \pm standard deviation. The D'Agostino-Pearson test was applied for Gaussian distribution analysis. The X² test was used to identify category differences by frequency. A paired Student's t-test was used to compare anthropometric and functional fitness parameters. Analysis of comparisons between currently and ideal silhouette were performed with two-way ANOVA followed by Bonferroni's post hoc test. Analysis with Pearson correlations was conducted to assess the association between ideal silhouette and BMI. Statistical significance was established at $p < 0.05$.

RESULTS

The groups showed no significant differences in the anthropometric variables: height (UN: 1.60 ± 1.2 , RT: 1.60 ± 1.6 ; m), weight (UN: 74 ± 7 kg; RT: 72 ± 7 kg), and BMI (UN: 29 ± 5 , RT: 28 ± 4 ; kg/m²). The functional strength of the lower and upper limbs measured by the chair stand test (UN: 19 ± 3 , RT: 29 ± 4 ; repetitions) and arm curl test (UN: 24 ± 5 , RT: 36 ± 4 ; repetitions), respectively, was significantly higher in the RT group than in the UN group ($p < 0.001$).

In relation to BMI (Figure 1A), no significant differences were found between the CS (5.45 ± 0.24) and IS (4.7 ± 0.12) in the RT group. However, the UN group showed a significant difference ($p < 0.01$) between the CS (10.4 ± 0.43) and IS (5.6 ± 0.26). Additionally, the CS result was significantly different




Figure 1 - Panel A: values (mean \pm SEM) for the scores of current and ideal silhouettes of the UN and RT groups. * $p < 0.02$ versus current silhouettes by the UN group. * $p < 0.01$ versus ideal silhouettes by the RT group. Panel B: percentage of relative values about body satisfaction (black) and dissatisfaction (white) * $p < 0.001$. Panel C: correlation between BI satisfaction and current BMI.



($p < 0.002$) between the UN and RT groups. However, their IS values did not differ significantly, as shown in Figure 1B.

For BI satisfaction, significant differences were observed (X^2 , $p < 0.001$). Eighteen women (90%) in the RT group and 17 (85%) in the UN group showed positive and negative satisfaction with their BI, respectively. Additionally, a positive correlation was observed ($p < 0.0086$) between the BMI and CS parameters (Figure 1C).

■ DISCUSSION

The main result of our study indicates the difference between the UN and RT groups as well as the similarity between the expected silhouettes in both groups. The data from this study were similar to those in Matsuo (16) in active elderly people, reflecting the contribution of physical activity to BI through increasing physical and functional fitness and improving self-perceptions of wellness (physiological, cognitive and emotional), thus improving self-esteem. These positive effects of physical activity can change the meaning of aging and body concept for elderly people and rebuild their BI with smaller discrepancies (9,10).

The greatest determinants for negative or positive BI are satisfaction with weight, accuracy of size perception, body satisfaction, appearance evaluation and orientation, body esteem, body ideal, body schema and body awareness (17). Moreover, BI can develop in relation to other factors, such as gender, age, media, and the relationship of the body with cognitive processes such as cultural and social beliefs, values and attitudes (17,18).

Human beings are biologically and socially dependent on their appearance: body form, skills, clothes, hair and appearance reflect integration and harmony with oneself (19). In a previous study, Ricciardelli, McCabe and Banfield (20) found improvements in both self-image and self-esteem associated with physical activity. Additionally, several different physical changes were observed with exercise in the study by Ginis et al. (21), and these changes were related to different aspects of BI, such as sex, body composition and strength.

In the present study, the idealized BI of the UN group was thinner than what they considered to be realistic. This perception could be due to cultural variables reflecting societal beliefs largely influenced by the media in favor of a lean and perfect body with the appearance of youth. Moreover, this difference could be associated with the stereotype of old age: diseases and biological decline.

Additionally, we found a positive correlation between BMI and CS, supporting the use of mean BMI values grouped by actual profiles in the characterization of ideal physical types for elderly people.

A physically active lifestyle helps elderly women make their own judgments related to self-assessment and BI. Body dissatisfaction can affect their lives substantially; in Western culture, an older body is often viewed as unattractive and unproductive.

Some limitations should be considered in this study. The instrument used in this study assessed silhouettes bidimensionally and allowed for little individual identification. The distribution of subcutaneous fat and other anthropometric features also have an important role in BI (22,23), but these factors were not taken into account.

Our data indicate that resistance training can improve the BI satisfaction of elderly women. The positive evaluation of

current silhouettes in the RT group may be related to their active lifestyles, thus supporting the use of physical activity to improve BI.

■ AUTHOR CONTRIBUTIONS

Rica RL, Gama EF and Bocalini DS performed the laboratory procedures, designed the study and were the main contributors to the writing of the manuscript. Machado AF, Alonso AC and Figueira-Júnior A performed the laboratory procedures, as well as, recruiting the volunteers, conducting the anthropometric measurements and filling out the questionnaires. Zanetti M and Brandão R evaluated the data from the statistical analysis. The revision of the manuscript was performed by Evangelista AL, Alves JV and Bergamin M, who contributed to the intellectual content of the study.

■ REFERENCES

- Schilder P. A imagem do corpo: as energias construtivas da psique. São Paulo: Martins Fontes, 1999.
- Head H, Holmes G. Sensory disturbances from cerebral lesions. *Brain*. 1911;34(2-3):102-254, <http://dx.doi.org/10.1093/brain/34.2.3.102>.
- Paillard J. Body Schema and body image - A double dissociation in deafferented patients. *Motor control, today and tomorrow*. 1999;12: 197-214.
- Benedikt R, Wertheim EH, Love A. Eating attitudes and weight-loss attempts in female adolescents and their mothers. *Journal of Youth and Adolescence*. 1998;27:43-57, <http://dx.doi.org/10.1023/A:1022876715005>.
- Bocalini DS, Serra AJ, Rica RL, dos Santos L. Repercussions of training and detraining by water-based exercise on functional fitness and quality of life: a short-term follow-up in healthy older women. *Clinics*. 2010; 65(12):1305-9, <http://dx.doi.org/10.1590/S1807-59322010001200013>.
- Bocalini DS, Serra AJ, dos Santos L. Moderate resistive training maintains bone mineral density and improves functional fitness in postmenopausal women. *J Aging Res*. 2010;2010:760818, <http://dx.doi.org/10.4061/2010/760818>.
- Bocalini DS, Serra AJ, Murad N, Levy RF. Water- versus land-based exercise effects on physical fitness in older women. *Geriatr Gerontol Int*. 2008;8(4):265-71, <http://dx.doi.org/10.1111/j.1447-0594.2008.00485.x>.
- Bocalini DS, Serra AJ, dos Santos L. Physical exercise improves the functional capacity and quality of life in patients with heart failure. *Clinics*. 2008;63(4):437-42, <http://dx.doi.org/10.1590/S1807-59322008000400005>.
- Cash TF, Pruzinsky T. Body image: development, deviance, and change. 1990; New York: Guilford Press.
- Fisher S. The evolution of psychological concepts about the body. In Cash TF, Pruzinsky T. Body images: development, deviance, and change. New York: Guilford Press. 1990;3-20.
- American College of Sports Medicine. ACSM guidelines for stress testing and its prescription. Rio de Janeiro: Guanabara, 2014.
- Benton MJ, Schlaier MC, Gibson DR. Change in quality of life among breast cancer survivors after resistance training: is there an effect of age? *J Aging Phys Act*. 2014;22(2):178-85, <http://dx.doi.org/10.1123/japa.2012-0227>.
- Rikli R, Jones CJ. Development and validation of a functional fitness test for community residing older adults. *J Aging Phys Act*. 1999;7(2):129-61, <http://dx.doi.org/10.1123/japa.7.2.129>.
- Stunkard AJ, Sorenson T, Schulsinger F. Use of Danish adoption register for the study of obesity and thinness. In Kety S, Rowland LP, Sidman RL, Mattysse SW. The genetics of neurological and psychiatric disorders. New York: Raven Press. 1983;115-120.
- Marsh HW, Roche LA. Predicting self-esteem from perceptions of actual and ideal ratings of body fatness: is there only one ideal "supermodel"? *Res Q Exerc Sport*. 1996;67(1):13-23, <http://dx.doi.org/10.1080/02701367.1996.10607921>.
- Matsuo RF, Velardi M, Brandão MRF, Miranda MLJ. Older women's body image and physical activity. *Rev Mack Edu Fis Esp*. 2007;6:37-43.
- McAuley E, Mihalko SL, Bane SM. Exercise and self-esteem in middle-aged adults: multidimensional relationship and physical fitness and self-efficacy influences. *J Behav Med*. 1997;20(1):67-83, <http://dx.doi.org/10.1023/A:1025591214100>.
- McAuley E, Blissmer B, Katula J, Duncan TE, Mihalko SL. Physical activity, self-esteem, and self-efficacy relationships in older adults: a randomized controlled trial. *Ann Behav Med*. 2000;22(2):131-9, <http://dx.doi.org/10.1007/BF02895777>.
- Morant H. BMA demands more responsible media attitude on body image. *BMJ*. 2000;320(7248):1495, <http://dx.doi.org/10.1136/bmj.320.7248.1495>.
- Ricciardelli LA, McCabe MP, Banfield S. Body image and body change methods in adolescent boys. Role of parents, friends and the media.



- J Psychosom Res. 2000;49(3):189-97, [http://dx.doi.org/10.1016/S0022-3999\(00\)00159-8](http://dx.doi.org/10.1016/S0022-3999(00)00159-8).
21. Martin Ginis KA, Eng JJ, Arbour KP, Hartman JW, Phillips SM. Mind over muscle? Sex differences in the relationship between body image change and subjective and objective physical changes following a 12-week strength-training program. *Body Image*. 2005;2(4):363-72, <http://dx.doi.org/10.1016/j.bodyim.2005.08.003>.
 22. Thompson JK, Coover MD, Stormer SM. Body image, social comparison, and eating disturbance: a covariance structure modeling investigation. *Int J Eat Disord*. 1999;26(1):43-51, [http://dx.doi.org/10.1002/\(SICI\)1098-108X\(199907\)26:1<43::AID-EAT6>3.0.CO;2-R](http://dx.doi.org/10.1002/(SICI)1098-108X(199907)26:1<43::AID-EAT6>3.0.CO;2-R).
 23. Volkein K, McConatha JT. Cultural contours of the body - The impact of age and fitness. In Lidor R, Bar-Eli E. *Innovations in sport psychology: Linking theory and practice*. Tel-Aviv: ISSP. 1997;744-746.