

# Revisiting the roles of VHR/DUSP3 phosphatase in human diseases

Lilian Cristina Russo, Jéssica Oliveira Farias, Pault Yeison Minaya Ferruzo, Lucas Falcão Monteiro, Fábio Luís Forti\*

Departamento de Bioquímica, Instituto de Química, Universidade de São Paulo, São Paulo, SP, BR.

Russo LC, Farias JO, Ferruzo PY, Monteiro LF, Forti FL. Revisiting the roles of VHR/DUSP3 phosphatase in human diseases. *Clinics*. 2018;73(suppl 1):e4666

\*Corresponding author. E-mail: flforti@iq.usp.br

Protein tyrosine phosphatases have long been considered key regulators of biological processes and are therefore implicated in the origins of various human diseases. Heterozygosity, mutations, deletions, and the complete loss of some of these enzymes have been reported to cause neurodegenerative diseases, autoimmune syndromes, genetic disorders, metabolic diseases, cancers, and many other physiological imbalances. Vaccinia H1-related phosphatase, also known as dual-specificity phosphatase 3, is a protein tyrosine phosphatase enzyme that regulates the phosphorylation of the mitogen-activated protein kinase signaling pathway, a central mediator of a diversity of biological responses. It has been suggested that vaccinia H1-related phosphatase can act as a tumor suppressor or tumor-promoting phosphatase in different cancers. Furthermore, emerging evidence suggests that this enzyme has many other biological functions, such as roles in immune responses, thrombosis, hemostasis, angiogenesis, and genomic stability, and this broad spectrum of vaccinia H1-related phosphatase activity is likely the result of its diversity of substrates. Hence, fully identifying and characterizing these substrate-phosphatase interactions will facilitate the identification of pharmacological inhibitors of vaccinia H1-related phosphatase that can be evaluated in clinical trials. In this review, we describe the biological processes mediated by vaccinia H1-related phosphatase, especially those related to genomic stability. We also focus on validated substrates and signaling circuitry with clinical relevance in human diseases, particularly oncogenesis.

**KEYWORDS:** Protein tyrosine phosphatase (PTP); Vaccinia H1-related phosphatase (VHR); Dual-specificity phosphatase 3 (DUSP3); Mitogen-activated protein kinase (MAPK).

## Protein Tyrosine Phosphatases, Dual-Specificity Phosphatases, and VHR/DUSP3

Protein tyrosine phosphatases (PTPs) are highly catalytically active enzymes that counteract the actions of protein tyrosine kinases (PTKs). This pair of families has different evolutionary origins. PTPs are organized into four classes (I to IV) based on the presence of either a cysteine or an aspartate in the catalytic site (1-3).

Class I PTPs are subdivided into classical tyrosine phosphatases and non-classical or dual-specificity tyrosine phosphatases (DUSPs), the latter of which are a less common group of enzymes that dephosphorylate both tyrosine and serine/threonine residues (1). DUSPs comprise the largest (61 proteins) and most diversified group of non-classical PTPs. A highly conserved consensus or signature sequence, HC(X)<sub>5</sub>R, is present in the catalytic domain of both classical and non-classical tyrosine phosphatases (1,4). The catalytic mechanisms by which PTPs and DUSPs exert their effects are

quite evolutionarily conserved and involve substrate hydrolysis and the formation of a stable phosphoryl-cysteine intermediate (5-7).

The atypical dual-specificity phosphatases (ADUSPs) include 19 proteins and were discovered in the last 10 years. Most of these proteins are small (<27 kDa and/or <250 aa), and because they have the ability to dephosphorylate Tyr and Ser/Thr residues, they can act on substrates, such as mRNAs, lipids, and glycogen (5,8).

ADUSPs possess the consensus catalytic domain of PTPs and typical DUSPs, and this domain is responsible for their phosphatase activity. However, ADUSPs also possess regulatory and/or recognition sequences at their C- and N-termini, and these surround the catalytic active site (9,10). ADUSPs do not have the CH2 (Cdc25 homology 2) domain that is normally associated with typical DUSPs (5,8) and is responsible for their specificity for mitogen-activated protein kinases (MAPKs) (8,11,12). However, even without these recognition sequences, some ADUSPs are capable of dephosphorylating ERK, JNK, and p38 (5,8).

This is the case for Vaccinia H1-related phosphatase (VHR) or dual-specificity phosphatase 3 (DUSP3), which was first identified in higher eukaryotes and was the first phosphatase to be crystallized (13,14). It weighs approximately 21 kDa and contains 185 residues, including PTP domains with the HC(X)<sub>5</sub>R consensus sequence. The VHR crystal structure includes a catalytic site that is 6 Å deep, which is shallower than all other classic PTPs (9 Å deep), and this may explain the substrate diversity observed among this enzyme (5,13,15). VHR is constitutively active, is widely expressed in several

**Copyright** © 2018 CLINICS – This is an Open Access article distributed under the terms of the Creative Commons License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium or format, provided the original work is properly cited.

No potential conflict of interest was reported.

**Received for publication on** November 22, 2017. **Accepted for publication on** May 18, 2018

Commemorative Edition: 10 years of ICESP

**DOI:** 10.6061/clinics/2018/e4666



tissues, and can localize to either the nucleus or cytosol, with its localization important to its various functions (16,17).

Among the biological roles attributed to VHR (discussed later) is its ability to control the cell cycle, which affects cell proliferation. This makes the study of VHR important for developing therapeutic strategies for different types of cancers. VHR behaves similar to a MAPK phosphatase (MKP or a typical DUSP) in that it can dephosphorylate ERK, JNK, and p38 (16,18-22) via a CH2-independent catalytic mechanism. VHR also displays increased catalytic activity in the presence of the serine/threonine kinase VRK3 and the tyrosine kinases Zap70 and Tyk2, all of which bind and/or chemically modify VHR to enhance its activity (19,22,23). These VHR-related regulatory mechanisms warrant further investigation as a potential therapeutic targets for controlling tumorigenesis (5,12).

In addition to MAPKs, other substrates have recently been shown to be targets of VHR. These include the following: 1) the ErbB receptor of EGF2 in non-small cell lung cancers (24), 2) signal transducer and activator of transcription 5 (STAT5) in immune cells stimulated by cytokines and/or growth factors (19), and 3) focal adhesion kinase (FAK), with which VHR plays a role in the formation and disassembly of focal adhesions (25). Furthermore, VHR has been found to have a new MAPK substrate-independent biological function that affects the maintenance of genomic stability. Some authors (26,27) recently suggested that many different nuclear proteins, such as NBS1, NPM, NUCL, hnRNP C1/C2, and ATM/ATR, that are directly or indirectly involved in different steps of the DNA damage response and DNA repair pathways are very likely to be substrates of VHR. These new putative protein targets and functions of VHR will be discussed in the following sections. We argue that this dual phosphatase constitutes a potential candidate for pharmacological targeting in clinical trials.

### Biological functions of VHR/DUSP3 and its association with human diseases

A high percentage of human cancers originate from one or more activating mutations and/or genomic aberrations in members of canonical signaling pathways, such as KRAS and EGFR, that specifically activate the downstream MAPK pathway (28). Furthermore, since VHR dephosphorylates MAPK and growth factor receptors (i.e., HER2) (17,24), VHR is relevant to the study of many types of cancers in which the activities of these signaling pathways are altered. For example, cervical cancer cell lines deficient in VHR arrest at the G1-S and G2-M cell cycle transitions and therefore show massively inhibited cell proliferation followed by initial signs of senescence (17,18).

MAPK activation plays a growth-promoting role during the early G1 phase of the cell cycle (17). Rahmouni et al. (17) showed that differential VHR expression is coupled with cell cycle transitions (i.e., high in the S, G2, and M phases) and may regulate MAPK activity. They also showed that VHR is expressed at low levels in early G1 phase, during which cyclin D1 expression is required for cell cycle progression (29), and that VHR is overexpressed in many types of cancers (30,31). For example, in breast cancers, cyclin D1 expression was found to be downregulated via a variety of mechanisms (32,33). One of these mechanisms involves the VHR-mediated overexpression of BRCA1-IRIS (34,35), which is responsible for Cyclin D1 overexpression and the induction of cell proliferation. Additionally, overexpressing at least one of the components of the

BRCA1-IRIS/EGFR/ErbB2 axis reduced VHR expression in normal and tumor breast cell lines, suggesting the development of more aggressive endocrine-resistant breast cancer phenotypes. Therefore, restoring VHR expression could potentially rescue cyclin D1 overexpression-dependent cellular transformation (35).

In non-small cell lung cancer (NSCLC) tissues, VHR expression and its activity against MAPKs were lower than those observed in normal lung tissues, while VHR activity against the ErbB2 receptor, which also regulates epithelial cell growth, was increased (24). Wang et al. (24) demonstrated that VHR can dephosphorylate a specific EGFR isoform in cell lines. The EGFR-VHR interaction is weak, and the binding of the EGF ligand to its receptor makes this interaction even weaker. However, overexpressing VHR reduced the phosphorylation of EGFR at Tyr-992 and made cells less responsive to EGF.

VHR overexpression suppressed cell proliferation in two- or three-dimensional cultures of H1299 cells and reduced the tumor sizes of xenografts. In addition, the mRNA and protein levels of VHR were markedly lower in tumor tissues than in adjacent healthy tissues in patients with non-small cell lung cancers (24). Thus, decreasing VHR expression might contribute to the initiation of lung cancer pathogenesis. Wagner et al. proposed that the onset of lung tumorigenesis in NSCLC involves the enhancement of ERK1/2 signaling as a result of the epigenetic downregulation of VHR by KDM2A demethylase (36). Some histone methylation modifiers, such as KDM2A, are overexpressed in NSCLC and are important for the tumorigenicity, proliferation, and invasiveness of these cells (37). Depleting KDM2A reduced the number of cells in S phase, inhibited their ability to form colonies in soft agar, and, most interestingly, upregulated (by up to 9-fold) the mRNA and protein expression levels of VHR. These data suggest that the epigenetic repression of VHR activates the downstream effects of ERK1/2 signal-mediated proliferation, contributing to the invasive characteristics of NSCLC.

Studies have correlated VHR overexpression with the onset or development of carcinogenic phenotypes. For example, VHR levels were higher in human HeLa, SiHa, CaSki, C33, and HT3 cell lines, which are derived from cervical tissues and consist of epithelial cells representing various grades of squamous carcinomas and adenocarcinomas, than in primary cells and normal tissues (38). In cervical cancer tissues, VHR is primarily enriched in nuclei. However, in tumor cell lines, VHR is localized in both the cytoplasm and nucleus, whereas in normal primary cells, it is exclusively cytosolic (38). The enrichment of VHR in cell lines and cervical cancer tissues is not due to increased gene or protein expression or the stabilization of its mRNA. Instead, it cause by an increase in the post-translational stabilization of the VHR protein (38).

VHR is also enriched in prostate cancer tissues (39). In normal prostate epithelial cells, androgen withdrawal led to decreased cell proliferation, increased apoptosis, and atrophy, whereas tumor prostate epithelial cells overcame these deficiencies by shifting to an androgen-independent state (40,41). When androgen-responsive prostate cancer cell lines were treated with synthetic androgens, various MKPs become overexpressed, including VHR. This changed was correlated with the systematic dysregulation of MAPK signaling observed during prostate carcinogenesis (42), as was observed in LNCaP cells treated with TPA/thapsigargin plus R1881. This dysregulation decreased JNK phosphorylation but did not affect pERK expression levels (43,44). Androgen-sensitive



grafts expressing VHR showed increased resistance to castration-induced apoptosis, and tumor regression was inversely correlated with VHR expression (39). One molecular mechanism that has been suggested for inhibiting prostate cancer progression is based on the finding that VHR knockdown led to JNK activation and apoptosis, suggesting potential therapeutic applications aimed at targeting this phosphatase.

Finally, VHR is also overexpressed in dysplastic nevi (DN), which are benign melanocytic tumors with a hyper-proliferative phenotype (45). Very recently, VHR was found to interact with and dephosphorylate FAK in H1299 cells and MEF/lung epithelial cells. In VHR-null mice, these cells were more migratory, and VHR overexpression decreased pFAK levels and inhibited cell migration (25). The enhanced motility phenotype observed in these cells supports the hypothesis that the VHR phosphatase plays important roles in the development of metastatic tumors.

It seems that with regard to the mechanisms underlying tumorigenesis, this dual phosphatase plays two opposing functions. Specifically, VHR knockdown inhibits the proliferation and invasiveness of cancer cell lines, in accordance with oncogene/oncoprotein models. However, in some cellular contexts, VHR downregulation triggers hyper-proliferation, migration and invasiveness in these cells. Thus, it can also be considered a tumor suppressor (46).

Another VHR function associated with tumor onset, establishment, growth, and maintenance is the regulation of angiogenesis, which is a crucial step during tumorigenesis. As shown previously in different types of cancers, endothelial cells express higher than normal levels of VHR and other MKPs, and this alteration was found to contribute to MAPK-independent PKC-mediated tubulogenesis *in vivo*. In the same FGF2-stimulated cells, VHR knockdown blocked angiogenic sprouting. Finally, in different *in vivo* and *ex vivo* models of VHR<sup>-/-</sup> mice, neovascularization was clearly decreased (47).

Genomic instability is characteristic of a broad spectrum of cancers, and genomic alterations can occur during any cell division. These alterations or instabilities are minimized by four major mechanisms: high-fidelity DNA replication during S phase, precise chromosome segregation in M phase, accurate and error-free repair of DNA damage, and a cell cycle progression that is coordinated by cell cycle checkpoints (48). Therefore, a disruption in any step in one or more of these four mechanisms can lead to genomic instability and contribute to cancer development.

In terms of MAPK signaling, with MAPKs the best characterized substrates of VHR, over-activating ERK1/2 induced multipolar spindles and aneuploidy in cells, while inhibiting ERK1/2 did not cause defects during chromosomal events such as the spindle assembly checkpoint (SAC) and mitotic exit (49). The role of this phosphatase in the formation of multipolar spindles in cancer cells was recently investigated (50). In early mitotic mammalian cells, both VHR and ERK1/2 localized to the spindle apparatus (17,51), and transient VHR inhibition promoted the formation of multipolar spindles in human mitotic cells (50). These studies also demonstrated that depleting ERK1/2 activity but not JNK restored the multipolarity induced by a lack of VHR and that overexpressing VHR reduced ERK1/2 phosphorylation by reversing multipolar spindles. These results suggest that the VHR-mediated regulation of ERK1/2 plays multiple roles in genomic stability (50).

Many of the nuclear events that control genomic stability depend on high concentrations of proteins and/or rapid

cytoplasm-nucleoplasm translocation. Interestingly, VHR is highly enriched in the nucleus of various cell lines (17,26,38,52), especially after genotoxic stress (26). This may indicate that this phosphatase has other substrates or that it has additional roles in maintaining genomic stability that could be directly or indirectly related to MAPK functions or other substrates. In fact, recent studies using bioinformatics approaches and validation analyses have suggested that novel VHR substrates are involved in genomic stability (26). In one study, the authors showed that VHR strongly co-localized with phospho-H2AX (Ser139) in cells exposed to radiation-induced DNA damage. They applied a bioinformatics analysis approach to identify human nuclear proteins that could be putative VHR substrates. Biochemical validation techniques were performed, resulting in some very promising targets, such as pATM (S1981), pATR (S428), pBRCA1 (S1423), BRCA2, CENP-F, Cyclin A, NBS1, APE1, MRE11, RAD50, pCHK2 (T68), and pP53 (S15). These results support the hypothesis that VHR is involved in genomic maintenance and that its depletion decreases survival and proliferation in human tumor cell lines by increasing DNA damage and/or delaying or impairing DNA repair (52).

Previous authors have also employed mass spectrometry to identify novel VHR substrates under cellular genotoxic stress conditions (27). Among the proteins identified, half were involved in mechanisms that control DNA and RNA structures and functions. Based on the presence of phosphorylatable tyrosine residues and what is known of the biological processes they regulate, experimental validation studies performed using cellular and biochemical assays suggested that nucleophosmin (NPM), nucleolin (NCL), and heterogeneous ribonucleoprotein isoforms C1/C2 (hnRNP C1/C2) are very likely to be VHR substrates (27). These proteins are tyrosine-phosphorylated *in vivo* and *in vitro* and could therefore be potential targets of dephosphorylation by VHR, especially because they are involved in cell cycle regulation and genomic instability (DNA damage response and repair) processes (53-56). Thus, the phosphatase activity of VHR against these three proteins should be further investigated.

VHR also mediates other signaling events in circulatory system cells and blood-related diseases (57). The first such studies were performed in resting T cells, which constitutively express VHR. During resting, activating TCR did not induce the expression of this enzyme via positive-feedback mechanisms. In T cells, VHR dephosphorylated MAPKs (especially JNK), but the UV radiation-induced activation of p38 was not targeted by VHR phosphatase activity (16). In Jurkat T lymphocyte cells, VHR activity was regulated by ZAP-70, a tyrosine kinase that phosphorylates VHR at Tyr138 (22). Basal ZAP-70 activity was sufficient to stimulate T cells, but under some circumstances, such as when IL-2 was used to stimulate TCR, ZAP-70 activity induced more VHR-Tyr138 at the T-cell pole. These mechanisms regulated TCR and induced signaling (22).

A TCR-related signaling pathway was also affected by VHR in IFN- $\beta$ -stimulated T and B cells via a mechanism involving STAT5. STAT5 is phosphorylated by tyrosine kinase 2 (TYK2). The Src homology 2 domain (SH2) of STAT5 recognizes phospho-Tyr138 on VHR and brings these domains together so that VHR can dephosphorylate STAT5. However, in this signaling cascade, VHR does not regulate the activity (via dephosphorylation) of other upstream effectors of the pathway, such as Janus kinase 1 (JAK1) and TYK2 (19).



Another study showed that silencing VHR in *Staphylococcus aureus*-infected human and murine macrophages increased the production of pro-inflammatory cytokines via NF- $\kappa$ B signaling. The phosphatase activity of VHR seemed to affect this inflammatory response via a negative feedback component, and the observed increase in cytokine levels may have been responsible for the hyper-responsiveness observed in the host immune system (58). In contrast to the pro-inflammatory activities of VHR, VHR-deficient mice displayed tolerance to LPS-induced endotoxic shock and polymicrobial septic shock. A massive increase in anti-inflammatory M2-like macrophages, decreased TNF production, and reduced ERK1/2 activity were observed in these animals. *In vivo*, at eighteen hours after endotoxic shock was induced by LPS, DUSP3<sup>-/-</sup> mice stabilize their body temperature, while DUSP3<sup>+/+</sup> mice remained hypothermic. However, VHR knockout did not significantly affect T and B lymphocytes, neutrophils, monocytes, or platelet counts (59).

Additional studies have shown that VHR involvement in the immune response to septic shock is dependent on sex hormones produced by the host animals. Knocking out VHR did not protect male mice from LPS-induced endotoxemia or septic shock, and the percentage of M2-like macrophages was lower in these mice than in VHR<sup>-/-</sup> female mice. The body temperatures of all groups of mice were lower after LPS injection (except in VHR<sup>-/-</sup> females), while resistance to sepsis was associated with the decreased activation of ERK1/2, PI3K, and AKT, suggesting phosphatase deficiency. These data demonstrate that in the absence of VHR, female sex hormones are very likely transcriptionally involved in the acquired resistance of VHR<sup>-/-</sup> mice to LPS-induced lethality (60).

Another very recent and important report provided mechanistic clues about the roles that VHR plays in platelet aggregation (61,62). This phosphatase seems to be responsible for reducing the levels of tyrosine phosphorylation of the enzymes Syk and PLC $\gamma$ 2, which regulate signaling via collagen receptor glycoprotein VI (GPVI) and C-type lectin-like type II (CLEC-2) receptors to promote aggregation in a MAPK-independent manner. VHR knock-out mice exhibited deficiencies in thrombus formation, suggesting that VHR contributes to arterial thrombosis but is unnecessary for primary hemostasis (61,62). The drugs currently used in clinics to treat thrombosis usually cause unwanted side effects, such as increasing the risk of gastrointestinal toxicity, neutropenia, thrombocytopenia, and bleeding, in addition to increasing the incidence of arterial thrombosis. Altogether, these novel functions of VHR strongly support the notion that this enzyme is a potential target for the development of new therapies for platelet aggregation in circulatory system cells and diseases (62).

### VHR expression and patient survival

It has been shown that VHR gene expression and protein levels are constitutively high in many different cell lines and diverse tissues (38,39). The VHR protein is highly stable even under high-stress conditions, such as cellular genotoxic or oxidative pressure (26,52). VHR gene and protein expression levels are not dependent on feedback regulation, such as that associated with many other MKPs (1,5,8,63). However, most VHR-inhibited cell lines exhibit phenotypes that are clearly associated with cell cycle arrest, senescence, and reduced cell survival even though VHR knock-out mice display a normal

phenotype and are free of apparent pathological features (17,19,60,61).


However, one of the key remaining questions is that of how VHR expression varies among patients carrying different types of cancer. By mining freely available databases containing gene expression information related to human cancers, it is possible to analyze, for example, correlations between patient survival times and VHR expression. Using this approach, we investigated the VHR gene at the BioProfiling.de (64,65) website, which provides excellent assays that can be used to examine the functions, interactions, and pathway relationships of a single gene or a list of genes or to perform clinical correlations to test a gene of interest as a biomarker or to analyze its gene synergy (SynTarget tool) with relation to cancer survival (PPISURV tool).

VHR-containing gene sets are available for 13 different cancer types, including approximately 50 cancer subtypes or clinical variables, in this web tool. To illustrate the potential of this toolkit (and because these topics are very tightly correlated with previously published cellular experiments), we selected VHR gene expression and patient survival for our analysis (64,65). As shown in Figure 1, positive survival curves were observed when VHR was overexpressed in lung, kidney, sarcoma, lymphoma, and breast cancer. All five of these types of cancers have different GEO dataset IDs (mostly obtained from TCGA, <https://cancergenome.nih.gov/>) and clinical variables and are presented as data obtained from different numbers of samples (ranging from 240 to 290). The data showed that low VHR expression had a significantly negative effect on patient survival. These results, despite the fact that they are associative and predictive, when analyzed in primary and immortalized cell lines derived from the same tumor tissues, overall survival then was substantially lower in cells deficient in VHR than in control cells (17,19,52,66).


However, as shown in Figure 2, VHR overexpression resulted in negative survival curves in lung, brain, and breast cancers. Although these cancers have different GEO dataset IDs, are associated with different clinical variables, and originated from a different numbers of samples (ranging from 115 to 450), high VHR expression had a statistically significant negative effect on survival in all three cancer types. These findings are in accordance with those presented in other reports in the literature showing that in some cell lines and tissues, VHR deficiency increases proliferation and survival (24,35), while VHR overexpression leads to cell death (usually by apoptosis).

An analysis of the correlations among the expression profiles of the VHR gene in various human cancers, patient survival times, and the cellular phenotypes related to VHR loss or gain of function (LOF or GOF) experiments showed that the biological functions of VHR seem to be tissue-specific. Additionally, VHR behaves both positively and negatively to regulate or mediate different types of human diseases. Therefore, the dual-specificity phosphatase VHR, similar to many other PTPs, has emerged as an attractive drug target for future clinical interventions (11,12,62,67).

General and non-specific PTP inhibitors have long been described in the literature, starting with sodium orthovanadate (68),  $\alpha$ -bromo-4-(carboxymethoxy)acetophenone (69), phenylarsine oxide (70), and 4-hydroxy-3,3-dimethyl-2H-benz[g]indole-2,5(3H)-dione (71). Although these have been shown to be very potent inhibitory drugs, their low selectivity makes them poor candidates for clinical trials.



**Figure 1** - Survival curves generated using gene expression databases available at BioProfiling.de showing examples of different types of cancers in which VHR downregulation was correlated with a significant reduction in patient survival times.



**Figure 2** - Survival curves generated using gene expression databases available at BioProfiling.de showing examples of different types of cancer in which VHR overexpression was correlated with a reduction in patient survival times.

Because of the diversity of substrates and biological functions attributed to VHR, which, as discussed in the previous sections, can be easily extended to animal studies and possibly even to a clinical setting, a great deal of effort has been made to identify specific VHR inhibitors. Once obtained, these compounds should be tested in animal models and humans. The results obtained in these studies will stimulate the development of therapies aimed at leveraging the inhibition of VHR and/or other phosphatases in diseases where the loss of VHR function may impair the proliferative capacity

of diseased cells (67). It is especially important to facilitate the development of inhibitors that do not affect cellular processes in surrounding healthy tissues.

Some potent pharmacological inhibitors of PTPs/DUSPs are already in preparation for clinical trials, while others have received a growing amount of attention in the last few years. These include compounds that are usually obtained (identified, isolated and purified) from natural sources, such as plants, fungi, bacteria and seaweeds, and have high specificity for particular phosphatases (72). Additionally, very



recently, an alternative to targeting DUSP enzymes at the transcriptional level was presented by research into omega 3 fatty acids ingestion. For example, docosahexaenoic acid (DHA) exerts a protective effect on mice brain development, and when it was added to the diet (in mice) as DHA-enriched fish-oil, it induced astroglial hyperactivation in a dose-dependent manner that was mediated by glial fibrillary acidic protein (GFAP) and dependent on increased MKP3 activity. This is the first evidence to show that natural compounds obtained from natural sources can have potential regulatory effects on PTPs/DUSPs during different cell biology and tissue developmental processes and suggest the possibility that they could be used to treat human pathologies (73).

## ■ ACKNOWLEDGEMENTS

The authors thank Prof. Dr. Roger Chammas (School of Medicine, University of Sao Paulo) and the Instituto do Cancer do Estado de Sao Paulo (ICESP) for paying the article processing charges associated with this manuscript. The project was supported by FAPESP (Grant No 2008/58264-5 and 2015/03983-0 to FLF). LCR is a post-doctoral fellow of the CAPES-PNPD program at the Institute of Chemistry, University of Sao Paulo. JOF is a PhD student and PYFM is a master's fellow of CNPq enrolled in the post-graduation program in Biochemistry and Molecular Biology, Institute of Chemistry-University, Sao Paulo. LFM is a master's fellow of CNPq in the Biotechnology program, University of Sao Paulo. We also thank BO and JRD for technical assistance in the LSSB laboratory.

## ■ AUTHOR CONTRIBUTIONS

Russo LC and Farias JO wrote the first part of the review. Ferruzo PYM and Monteiro LF wrote the second part of this review. Forti FL wrote the third part and provided the figure data and also was the scientific mentor and supervisor of all authors.

## ■ REFERENCES

- Alonso A, Sasin J, Bottini N, Friedberg I, Friedberg I, Osterman A, et al. Protein tyrosine phosphatases in the human genome. *Cell*. 2004;117(6):699-711, <http://dx.doi.org/10.1016/j.cell.2004.05.018>.
- Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanam S. The protein kinase complement of the human genome. *Science*. 2002;298(5600):1912-34, <http://dx.doi.org/10.1126/science.1075762>.
- Mustelin T. A Brief Introduction to the Protein Phosphatase Families. In: Moorhead G, editor. *Protein Phosphatase Protocols*. Totowa, NJ: Springer; 2007. p. 9-122.
- Farooq A, Zhou MM. Structure and regulation of MAPK phosphatases. *Cell Signal*. 2004;16(7):769-79, <http://dx.doi.org/10.1016/j.cellsig.2003.12.008>.
- Bayón Y, Alonso A. Atypical DUSPs: 19 phosphatases in search of a role. In: Lazo PA editor. *Emerging Signaling Pathways in Tumor Biology*. Kerala, India: Transworld Research Network; 2010. p.185-208.
- Denu JM, Dixon JE. A catalytic mechanism for the dual-specific phosphatases. *Proc Natl Acad Sci U S A*. 1995;92(13):5910-4, <http://dx.doi.org/10.1073/pnas.92.13.5910>.
- Denu JM, Dixon JE. Protein tyrosine phosphatases: mechanisms of catalysis and regulation. *Curr Opin Chem Biol*. 1998;2(5):633-41, [http://dx.doi.org/10.1016/S1367-5931\(98\)80095-1](http://dx.doi.org/10.1016/S1367-5931(98)80095-1).
- Patterson KI, Brummer T, O'Brien PM, Daly RJ. Dual-specificity phosphatases: critical regulators with diverse cellular targets. *Biochem J*. 2009;418(3):475-89, <http://dx.doi.org/10.1042/BJ20082234>.
- Alonso A, Narisawa S, Bogetz J, Tautz L, Hadzic R, Huynh H, et al. VHY, a novel myristoylated testis-restricted dual specificity protein phosphatase related to VHx. *J Biol Chem*. 2004;279(31):32586-91, <http://dx.doi.org/10.1074/jbc.M403442200>.
- Alonso A, Rojas A, Godzik A, Mustelin T. The dual-specific protein tyrosine phosphatase family. In: Arino J, Alexander D, editors. *Protein Phosphatases*. Berlin, Heidelberg: Springer; 2004. p. 333-58.
- Bermudez O, Pagès G, Gimond C. The dual-specificity MAP kinase phosphatases: critical roles in development and cancer. *Am J Physiol Cell Physiol*. 2010;299(2):C189-202, <http://dx.doi.org/10.1152/ajpcell.00347.2009>.
- Nunes-Xavier C, Romá-Mateo C, Ríos P, Tárrega C, Cejudo-Marín R, Tabernero L, et al. Dual-specificity MAP kinase phosphatases as targets of cancer treatment. *Anticancer Agents Med Chem*. 2011;11(1):109-32, <http://dx.doi.org/10.2174/187152011794941190>.
- Ishibashi T, Bottaro DP, Chan A, Miki T, Aaronson SA. Expression cloning of a human dual-specificity phosphatase. *Proc Natl Acad Sci U S A*. 1992;89(24):12170-4, <http://dx.doi.org/10.1073/pnas.89.24.12170>.
- Yuvaniyama J, Denu JM, Dixon JE, Saper MA. Crystal structure of the dual specificity protein phosphatase VHR. *Science*. 1996;272(5266):1328-31, <http://dx.doi.org/10.1126/science.272.5266.1328>.
- Barford D, Flint AJ, Tonks NK. Crystal structure of human protein tyrosine phosphatase 1B. *Science*. 1994;263(5152):1397-404, <http://dx.doi.org/10.1126/science.8128219>.
- Alonso A, Saxena M, Williams S, Mustelin T. Inhibitory role for dual specificity phosphatase VHR in T cell antigen receptor and CD28-induced Erk and Jnk activation. *J Biol Chem*. 2001;276(7):4766-71, <http://dx.doi.org/10.1074/jbc.M006497200>.
- Rahmouni S, Cerignoli F, Alonso A, Tsutji T, Henkens R, Zhu C, et al. Loss of the VHR dual-specific phosphatase causes cell-cycle arrest and senescence. *Nat Cell Biol*. 2006;8(5):524-31, <http://dx.doi.org/10.1038/ncb1398>.
- Cerignoli F, Rahmouni S, Ronai Z, Mustelin T. Regulation of MAP kinases by the VHR dual-specific phosphatase: implications for cell growth and differentiation. *Cell Cycle*. 2006;5(19):2210-5, <http://dx.doi.org/10.4161/cc.5.19.3267>.
- Hoyt R, Zhu W, Cerignoli F, Alonso A, Mustelin T, David M. Cutting edge: selective tyrosine dephosphorylation of interferon-activated nuclear STAT5 by the VHR phosphatase. *J Immunol*. 2007;179(6):3402-6, <http://dx.doi.org/10.4049/jimmunol.179.6.3402>.
- Kondoh K, Nishida E. Regulation of MAP kinases by MAP kinase phosphatases. *Biochim Biophys Acta*. 2007;1773(8):1227-37, <http://dx.doi.org/10.1016/j.bbamer.2006.12.002>.
- Todd JL, Tanner KG, Denu JM. Extracellular regulated kinases (ERK) 1 and ERK2 are authentic substrates for the dual-specificity protein-tyrosine phosphatase VHR. A novel role in down-regulating the ERK pathway. *J Biol Chem*. 1999;274(19):13271-80, <http://dx.doi.org/10.1074/jbc.274.19.13271>.
- Alonso A, Rahmouni S, Williams S, van Stipdonk M, Jaroszewski L, Godzik A, et al. Tyrosine phosphorylation of VHR phosphatase by ZAP-70. *Nat Immunol*. 2003;4(1):44-8, <http://dx.doi.org/10.1038/ni856>.
- Kang TH, Kim KT. Negative regulation of ERK activity by VRK3-mediated activation of VHR phosphatase. *Nat Cell Biol*. 2006;8(8):863-9, <http://dx.doi.org/10.1038/ncb1447>.
- Wang JY, Yeh CL, Chou HC, Yang CH, Fu YN, Chen YT, et al. Vaccinia H1-related phosphatase is a phosphatase of ErbB receptors and is down-regulated in non-small cell lung cancer. *J Biol Chem*. 2011;286(12):10177-84, <http://dx.doi.org/10.1074/jbc.M110.163295>.
- Chen YR, Chou HC, Yang CH, Chen HY, Liu YW, Lin TY, et al. Deficiency in VHR/DUSP3, a suppressor of focal adhesion kinase, reveals its role in regulating cell adhesion and migration. *Oncogene*. 2017;36(47):6509-17, <http://dx.doi.org/10.1038/onc.2017.255>.
- Forti FL. Combined experimental and bioinformatics analysis for the prediction and identification of VHR/DUSP3 nuclear targets related to DNA damage and repair. *Integr Biol (Camb)*. 2015;7(1):73-89.
- Panico K, Forti FL. Proteomic, cellular, and network analyses reveal new DUSP3 interactions with nucleolar proteins in HeLa cells. *J Proteome Res*. 2013;12(12):5851-66, <http://dx.doi.org/10.1021/pr400867j>.
- Herbst RS, Heymach JV, Lippman SM. Lung cancer. *N Engl J Med*. 2008;359(13):1367-80, <http://dx.doi.org/10.1056/NEJMra0802714>.
- Baldin V, Lukas J, Marcote MJ, Pagano M, Draetta G. Cyclin D1 is a nuclear protein required for cell cycle progression in G1. *Genes Dev*. 1993;7(5):812-21, <http://dx.doi.org/10.1101/gad.7.5.812>.
- Sicinski P, Donaher JL, Parker SB, Li T, Fazeli A, Gardner H, et al. Cyclin D1 provides a link between development and oncogenesis in the retina and breast. *Cell*. 1995;82(4):621-30, [http://dx.doi.org/10.1016/0092-8674\(95\)90034-9](http://dx.doi.org/10.1016/0092-8674(95)90034-9).
- Tetsu O, McCormick F. Beta-catenin regulates expression of cyclin D1 in colon carcinoma cells. *Nature*. 1999;398(6726):422-6, <http://dx.doi.org/10.1038/18884>.
- Ahnström M, Nordenskjöld B, Rutqvist LE, Skoog L, Stål O. Role of cyclin D1 in ErbB2-positive breast cancer and tamoxifen resistance. *Breast Cancer Res Treat*. 2005;91(2):145-51, <http://dx.doi.org/10.1007/s10549-004-6457-4>.
- Sicinski P, Weinberg RA. A specific role for cyclin D1 in mammary gland development. *J Mammary Gland Biol Neoplasia*. 1997;2(4):335-42, <http://dx.doi.org/10.1023/A:1026391128117>.
- ElShamy WM, Livingston DM. Identification of BRCA1-IRIS, a BRCA1 locus product. *Nat Cell Biol*. 2004;6(10):954-67, <http://dx.doi.org/10.1038/ncb1171>.
- Hao L, ElShamy WM. BRCA1-IRIS activates cyclin D1 expression in breast cancer cells by downregulating the JNK phosphatase DUSP3/VHR. *Int J Cancer*. 2007;121(1):39-46, <http://dx.doi.org/10.1002/ijc.22597>.
- Wagner KW, Alam H, Dhar SS, Giri U, Li N, Wei Y, et al. KDM2A promotes lung tumorigenesis by epigenetically enhancing ERK1/2 signaling. *J Clin Invest*. 2013;123(12):5231-46, <http://dx.doi.org/10.1172/JCI68642>.
- Shi Y. Histone lysine demethylases: emerging roles in development, physiology and disease. *Nat Rev Genet*. 2007;8(11):829-33, <http://dx.doi.org/10.1038/nrg2218>.



38. Henkens R, Delvenne P, Arafa M, Moutschen M, Zeddou M, Tautz L, et al. Cervix carcinoma is associated with an up-regulation and nuclear localization of the dual-specificity protein phosphatase VHR. *BMC Cancer*. 2008;8:147, <http://dx.doi.org/10.1186/1471-2407-8-147>.
39. Arnoldussen YJ, Lorenzo PI, Pretorius ME, Waehre H, Risberg B, Maelandsmo GM, et al. The mitogen-activated protein kinase phosphatase vaccinia H1-related protein inhibits apoptosis in prostate cancer cells and is overexpressed in prostate cancer. *Cancer Res*. 2008;68(22):9255-64, <http://dx.doi.org/10.1158/0008-5472.CAN-08-1224>.
40. Mercader M, Sengupta S, Bodner BK, Manecke RG, Cosar EF, Moser MT, et al. Early effects of pharmacological androgen deprivation in human prostate cancer. *BJU Int*. 2007;99(1):60-7, <http://dx.doi.org/10.1111/j.1464-410X.2007.06538.x>.
41. Feldman BJ, Feldman D. The development of androgen-independent prostate cancer. *Nat Rev Cancer*. 2001;1(1):34-45, <http://dx.doi.org/10.1038/35094009>.
42. Bakin RE, Gioeli D, Sikes RA, Bissonette EA, Weber MJ. Constitutive activation of the Ras/mitogen-activated protein kinase signaling pathway promotes androgen hypersensitivity in LNCaP prostate cancer cells. *Cancer Res*. 2003;63(8):1981-9.
43. Engedal N, Korkmaz CG, Saatcioglu F. C-Jun N-terminal kinase is required for phorbol ester- and thapsigargin-induced apoptosis in the androgen responsive prostate cancer cell line LNCaP. *Oncogene*. 2002;21(7):1017-27, <http://dx.doi.org/10.1038/sj.onc.1205167>.
44. Lorenzo PI, Saatcioglu F. Inhibition of apoptosis in prostate cancer cells by androgens is mediated through downregulation of c-Jun N-terminal kinase activation. *Neoplasia*. 2008;10(5):418-28, <http://dx.doi.org/10.1593/neo.07985>.
45. Mitsui H, Kiecker F, Shemer A, Cannizzaro MV, Wang CQF, Gulati N, et al. Discrimination of Dysplastic Nevi from Common Melanocytic Nevi by Cellular and Molecular Criteria. *J Invest Dermatol*. 2016;136(10):2030-40, <http://dx.doi.org/10.1016/j.jid.2015.11.035>.
46. Weinstein IB, Joe A. Oncogene addiction. *Cancer Res*. 2008;68(9):3077-80, <http://dx.doi.org/10.1158/0008-5472.CAN-07-3293>.
47. Amand M, Erpicum C, Bajou K, Cerignoli F, Blacher S, Martin M, et al. DUSP3/VHR is a pro-angiogenic atypical dual-specificity phosphatase. *Mol Cancer*. 2014;13:108, <http://dx.doi.org/10.1186/1476-4598-13-108>.
48. Shen Z. Genomic instability and cancer: an introduction. *J Mol Cell Biol*. 2011;3(1):1-3, <http://dx.doi.org/10.1093/jmcb/mjq057>.
49. Eves EM, Shapiro P, Naik K, Klein UR, Trakul N, Rosner MR. Raf kinase inhibitory protein regulates aurora B kinase and the spindle checkpoint. *Mol Cell*. 2006;23(4):561-74, <http://dx.doi.org/10.1016/j.molcel.2006.07.015>.
50. Tambe MB, Narvi E, Kallio M. Reduced levels of Dusp3/Vhr phosphatase impair normal spindle bipolarity in an Erk1/2 activity-dependent manner. *FEBS Lett*. 2016;590(16):2757-67, <http://dx.doi.org/10.1002/1873-3468.12310>.
51. Willard FS, Crouch MF. MEK, ERK, and p90RSK are present on mitotic tubulin in Swiss 3T3 cells: a role for the MAP kinase pathway in regulating mitotic exit. *Cell Signal*. 2001;13(9):653-64, [http://dx.doi.org/10.1016/S0898-6568\(01\)00185-1](http://dx.doi.org/10.1016/S0898-6568(01)00185-1).
52. Torres TE, Russo LC, Santos A, Marques GR, Magalhaes YT, Tabassum S, et al. Loss of DUSP3 activity radiosensitizes human tumor cell lines via attenuation of DNA repair pathways. *Biochim Biophys Acta*. 2017;1861(7):1879-94, <http://dx.doi.org/10.1016/j.bbagen.2017.04.004>.
53. Srivastava M, Pollard HB. Molecular dissection of nucleolin's role in growth and cell proliferation: new insights. *FASEB J*. 1999;13(14):1911-22, <http://dx.doi.org/10.1096/fasebj.13.14.1911>.
54. Miranda GA, Chokler I, Aguilera RJ. The murine nucleolin protein is an inducible DNA and ATP binding protein which is readily detected in nuclear extracts of lipopolysaccharide-treated splenocytes. *Exp Cell Res*. 1995;217(2):294-308, <http://dx.doi.org/10.1006/excr.1995.1090>.
55. Christian KJ, Lang MA, Raffalli-Mathieu F. Interaction of heterogeneous nuclear ribonucleoprotein C1/C2 with a novel cis-regulatory element within p53 mRNA as a response to cytostatic drug treatment. *Mol Pharmacol*. 2008;73(5):1558-67, <http://dx.doi.org/10.1124/mol.107.042507>.
56. Yang C, Maiguel DA, Carrier F. Identification of nucleolin and nucleophosmin as genotoxic stress-responsive RNA-binding proteins. *Nucleic Acids Res*. 2002;30(10):2251-60, <http://dx.doi.org/10.1093/nar/30.10.2251>.
57. Vang T, Miletic AV, Arimura Y, Tautz L, Rickert RC, Mustelin T. Protein tyrosine phosphatases in autoimmunity. *Annu Rev Immunol*. 2008;26:29-55, <http://dx.doi.org/10.1146/annurev.immunol.26.021607.090418>.
58. Yan Q, Sharma-Kuinkel BK, Deshmukh H, Tsalkis EL, Cyr DD, Lucas J, et al. Dusp3 and Psme3 are associated with murine susceptibility to *Staphylococcus aureus* infection and human sepsis. *PLoS Pathog*. 2014;10(6):e1004149, <http://dx.doi.org/10.1371/journal.ppat.1004149>.
59. Singh P, Dejager L, Amand M, Theatre E, Vandereyken M, Zurashvili T, et al. DUSP3 Genetic Deletion Confers M2-like Macrophage-Dependent Tolerance to Septic Shock. *J Immunol*. 2015;194(10):4951-62, <http://dx.doi.org/10.4049/jimmunol.1402431>.
60. Vandereyken MM, Singh P, Wathieu CP, Jacques S, Zurashvili T, Dejager L, et al. Dual-Specificity Phosphatase 3 Deletion Protects Female, but Not Male, Mice from Endotoxemia-Induced and Polymicrobial-Induced Septic Shock. *J Immunol*. 2017;199(7):2515-27, <http://dx.doi.org/10.4049/jimmunol.1602092>.
61. Musumeci L, Kuijpers MJ, Gilio K, Hego A, Théâtre E, Maurissen L, et al. Dual-specificity phosphatase 3 deficiency or inhibition limits platelet activation and arterial thrombosis. *Circulation*. 2015;131(7):656-68, <http://dx.doi.org/10.1161/CIRCULATIONAHA.114.010186>.
62. Tautz L, Senis YA, Oury C, Rahmouni S. Perspective: Tyrosine phosphatases as novel targets for antiplatelet therapy. *Bioorg Med Chem*. 2015;23(12):2786-97, <http://dx.doi.org/10.1016/j.bmc.2015.03.075>.
63. Owens DM, Keyse SM. Differential regulation of MAP kinase signalling by dual-specificity protein phosphatases. *Oncogene*. 2007;26(22):3203-13, <http://dx.doi.org/10.1038/sj.onc.1210412>.
64. Antonov AV. BioProfiling.de: analytical web portal for high-throughput cell biology. *Nucleic Acids Res*. 2011;39(Web Server issue):W323-7, <http://dx.doi.org/10.1093/nar/gkr372>.
65. Amelio I, Tsvetkov PO, Knight RA, Lisitsa A, Melino G, Antonov AV. SynTarget: an online tool to test the synergistic effect of genes on survival outcome in cancer. *Cell Death Differ*. 2016;23(5):912, <http://dx.doi.org/10.1038/cdd.2016.12>.
66. Arnoldussen YJ, Saatcioglu F. Dual specificity phosphatases in prostate cancer. *Mol Cell Endocrinol*. 2009;309(1-2):1-7, <http://dx.doi.org/10.1016/j.mce.2009.05.019>.
67. Stanford SM, Bottini N. Targeting Tyrosine Phosphatases: Time to End the Stigma. *Trends Pharmacol Sci*. 2017;38(6):524-40, <http://dx.doi.org/10.1016/j.tips.2017.03.004>.
68. Tamura S, Brown TA, Whipple JH, Fujita-Yamaguchi Y, Dubler RE, Cheng K, et al. A novel mechanism for the insulin-like effect of vanadate on glycogen synthase in rat adipocytes. *J Biol Chem*. 1984;259(10):6650-8.
69. Arabaci G, Yi T, Fu H, Porter ME, Beebe KD, Pei D. alpha-bromoacetophenone derivatives as neutral protein tyrosine phosphatase inhibitors: structure-Activity relationship. *Bioorg Med Chem Lett*. 2002;12(21):3047-50, [http://dx.doi.org/10.1016/S0960-894X\(02\)00681-9](http://dx.doi.org/10.1016/S0960-894X(02)00681-9).
70. Levenson RM, Blackshear PJ. Insulin-stimulated protein tyrosine phosphorylation in intact cells evaluated by giant two-dimensional gel electrophoresis. *J Biol Chem*. 1989;264(33):19984-93.
71. Liljebris C, Baranczewski P, Björkstrand E, Byström S, Lundgren B, Tjernberg A, et al. Oxidation of protein tyrosine phosphatases as a pharmaceutical mechanism of action: a study using 4-hydroxy-3,3-dimethyl-2H-benzof[glindole-2,5(3H)-dione. *J Pharmacol Exp Ther*. 2004;309(2):711-9, <http://dx.doi.org/10.1124/jpet.103.062745>.
72. Zhang ZY. Drugging the Undruggable: Therapeutic Potential of Targeting Protein Tyrosine Phosphatases. *Acc Chem Res*. 2017;50(1):122-9, <http://dx.doi.org/10.1021/acs.accounts.6b00537>.
73. Tripathi S, Kushwaha R, Mishra J, Gupta MK, Kumar H, Sanyal S, et al. Docosahexaenoic acid up-regulates both PI3K/AKT-dependent FABP7-PPARγ interaction and MKP3 that enhance GFAP in developing rat brain astrocytes. *J Neurochem*. 2017;140(1):96-113, <http://dx.doi.org/10.1111/jnc.13879>.