

Epidemiology and biology of cutaneous human papillomavirus

Emily M. Nunes,* Valéria Talpe-Nunes, Laura Sichero

Centro de Investigação Translacional em Oncologia, Instituto do Cancer do Estado de Sao Paulo (ICESP), Hospital das Clinicas HCFMUSP, Faculdade de Medicina, Universidade de Sao Paulo, Sao Paulo, SP, BR.

Nunes EM, Talpe-Nunes V, Sichero L. Epidemiology and biology of cutaneous human papillomavirus. Clinics. 2018;73(suppl 1):e489s

*Corresponding author. E-mail: emontosa.biotec@gmail.com

Cutaneous human papillomaviruses (HPVs) include β - and γ -HPVs, in addition to a small fraction of α -HPVs. β -HPVs were first isolated from patients with the rare genetic disorder *Epidermodysplasia verruciformis*, and they are associated with the development of nonmelanoma skin cancer at sun-exposed skin sites in these individuals. Organ transplant recipients also have greater susceptibility to β -HPV infection of the skin and an increased risk of developing nonmelanoma skin cancer. In both immunosuppressed and immunocompromised individuals, cutaneous HPVs are ubiquitously disseminated throughout healthy skin and may be an intrinsic part of the commensal flora. Functional analysis of E6 and E7 proteins of specific cutaneous HPVs has provided a mechanistic comprehension of how these viruses may induce carcinogenesis. Nevertheless, additional research is crucial to better understand the pathological implications of the broad distribution of these HPVs.

KEYWORDS: Human Papillomavirus; Cutaneous; Prevalence; Nonmelanoma Skin Cancer.

■ INTRODUCTION

Human papillomavirus (HPV) represents a diverse group of viruses infecting mainly epithelial and mucosal tissues (1). Based on the identity of the *L1* major capsid gene sequence, the majority of the over 200 viral types characterized to date cluster within the alpha (α)-, beta (β)-, or gamma (γ)-HPV genus (2, 3). While the great majority of α -HPVs are mucosal types isolated from the anogenital epithelia, some viral types in this genus (e.g., HPVs 2, 3 and 10) and β - and γ -HPVs were originally designated cutaneous types (Figure 1). To date, 54 β -HPVs (subdivided into 5 species, β 1-5) and 98 γ -HPVs (subdivided into 27 species, γ 1-27) have been fully sequenced and characterized (http://www.nordicehealth.se/hpv-center/reference_clones/), and it is expected that these numbers will further increase once partial sequences of putative novel types are described (4). Some cutaneous HPVs are clearly associated with the development of various skin lesions, from warts to carcinomas, in restricted populations (1,5). Nevertheless, it has proven difficult to determine the role of particular β -HPVs in cutaneous malignancies because of the high viral diversity and ubiquity of multiple types throughout healthy skin, the oral cavity, the nasal mucosa and the anogenital region (6-11).

Prevalence and distribution of cutaneous HPVs among immunosuppressed individuals

The first record concerning the association of HPV with papillomatous skin lesions that harbored carcinogenic potential dates to the early 1920s, when Lewandowsky and Lutz (12) first described a hereditary condition named *Epidermodysplasia verruciformis* (EV) that is characterized by extensive warts throughout the body (Table 1). Later, Jablonska et al. (13) observed that EV individuals infected with β -HPV 5 and 8 had a higher risk of developing nonmelanoma skin cancer (NMSC), particularly in ultraviolet (UV)-exposed sites. Together, both viral types are detected in approximately 90% of skin squamous cell carcinomas (SCCs) in EV patients. In these cases, β -HPVs are actively transcribed and generally persistent at high copy number (14). Currently, β -HPV 5 and 8 are accepted as possible etiological agents (carcinogen group 2B) of cutaneous SCC (cSCC) in immunosuppressed EV individuals by the International Agency for Research on Cancer (IARC) (15).

In the following years, several studies focused on analyzing the oncogenic potential of cutaneous HPVs in other immunosuppressed individuals, among which organ transplant recipients (OTRs) were the most extensively investigated. OTRs slightly resemble EV patients in that they are often covered with wart-like skin lesions and actinic keratosis (AK). Furthermore, OTRs have up to a 100-fold increased risk of developing NMSC compared to the general population (16-17). Importantly, the increased risk of NMSC is mostly associated with a higher incidence of cSCC (18-20). Clinical and histological features of these lesions suggest that cSCCs occasionally develop from viral warts or other precursor lesions (21,22). NMSC in OTRs often presents as multiple lesions and is usually confined to UV-exposed anatomical sites, most likely associated with local immunosuppression (23,24). These tumors are also more aggressive

Copyright © 2018 CLINICS – This is an Open Access article distributed under the terms of the Creative Commons License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium or format, provided the original work is properly cited.

No potential conflict of interest was reported.

Received for publication on December 1, 2017. **Accepted for publication on** February 6, 2018

Commemorative Edition: 10 years of ICESP

DOI: 10.6061/clinics/2018/e489s




Figure 1 - Distribution of cutaneous viral types within different HPV genera. The number of cutaneous viral types within each genus is indicated.

Table 1 - Select studies on the prevalence and distribution of cutaneous HPVs among immunosuppressed individuals.

Year	Author (s) (reference number)	Data
1922	Lewandowsky and Lutz (12)	First description of epidermodysplasia verruciformis (EV).
1972	Jablonska et al. (13)	β -HPVs 5 and 8 infected EV individuals had a higher risk of developing NMSC (after UV exposition).
1974	Koranda et al. (26)	Cutaneous warts were detected in 43% of OTR individuals after 3 months to 9 years following transplant.
1976	Mullen et al. (20)	Increased risk of NMSC is mostly associated to higher incidence of cSCC.
1977	Hoxtell et al. (19)	
1980	Hardie et al. (18)	
1978	Sbano et al. (22)	cSCC occasionally develop from viral warts or other precursor lesions.
1989	Barr et al. (21)	
1980	Hardie et al. (18)	The incidence of skin cancer increases 5% per year after the first year of transplant, with a cumulative risk of 44% after 9 years.
1980	Hardie et al. (18)	Tumors are more aggressive in OTRs than in the general population.
1984	Boyle et al. (27)	18% of renal transplant patients who were highly exposed to UV developed carcinogenic lesion in the skin.
1995	Birkeland et al. (16)	OTRs have until 100 fold increased risk of developing NMSC as compared to the general population.
2000	Lindelöf et al. (17)	
1997	Boxman et al. (35)	
2000	Harwood et al. (36)	β -HPV is more prevalent in skin warts biopsies than in both the normal skin and plucked hairs among OTRs.
2003	Meyer et al. (37)	
2000	Antonsson et al. (6)	Among OTRs, dialysis patients, and healthy controls, solely the first group reported ever having skin cancer (11.5%).
2000	Lindelöf et al. (17)	Within 15 years of transplantation, up to 90% of OTRs develop warts and/or cSCC.
2000	Berkhout et al. (29)	Infections of cutaneous HPVs frequently persist in OTRs.
2007	Hazard et al. (30)	
2003	Feltkamp et al. (40)	There is a positive epidemiological association between β -HPV seroreactivity and cSCC development.
2004	Termorshuizen et al. (24)	NMSCs among OTRs are often multiple and usually confined to UV-exposed anatomical sites.
2004	Harwood et al. (34)	
2007	Forslund et al. (23)	
2005	Weissenborn et al. (46)	Data regarding the association between cutaneous HPV infection and cSCC is still inconclusive.
2008	Rollison et al. (45)	
2011	Arron et al. (43)	
2016	Chahoud et al. (44)	
2007	Nindl et al. (28)	OTRs have higher cutaneous HPV prevalence rate up to 90% in cSCC compared to the normal skin (11-32%).
2007	Hazard et al. (30)	Older ages and history of sunburn are associated to an elevated risk of β -HPV persistent infection.
2014	Hampras et al. (31)	
2008	Michael et al. (39)	Seroconversion to β -HPV increases with age.
2010	Antonsson et al. (38)	
2009	Bouvard et al. (15)	β -HPVs 5 and 8 are accepted as possible etiological agents (carcinogens group 2B) of cSCC in immunosuppressed EV individuals.
2011	Proby et al. (33)	Individuals with concordant β -HPV DNA in plucked eyebrow hairs and serologic tests had a significantly increased risk of developing SCC.
2013	Neale et al. (32)	There is a significant association between the number of β -HPVs detected at eyebrow hair follicles and the increased risk of cSCC among OTRs.



in OTRs than in the general population and form metastases more readily (18,25).

Among OTRs, cutaneous warts were detected in 43% of individuals at 3 months to 9 years following transplant (25,26). Additionally, within 15 years of transplantation, up to 90% of OTRs develop warts and/or cSCC (17). Upon analyzing skin smears from OTRs, dialysis patients and healthy controls, Antonsson et al. (6) observed that 11.5% of OTRs reported ever having skin cancer, whereas no cases of NMSC were observed in the other groups studied. NMSC incidence in OTRs varies depending on the duration of immunosuppression; Hardie et al. (18) demonstrated that the incidence of skin cancer increased 5% per year after the first year of transplant, with a cumulative risk of 44% after 9 years.

The incidence of NMSC is also related to long sun exposure. Boyle et al. (27) observed that 18% of renal transplant patients with high exposure levels to sunshine (>3 months in a tropical or subtropical climate or >5 years in an outdoor occupation) developed carcinogenic alterations in their skin: two patients were diagnosed with cSCC, and seven were diagnosed with AK, whereas neither lesion type was noted in the other patients or in the control group.

Although cumulative sun exposure is the major risk factor for NMSC, recent studies have revealed a role for HPV as a cofactor in association with UV radiation in cSCC in OTRs. OTRs have a higher cutaneous HPV prevalence rate in cSCC (up to 90%) than in normal skin (11-32%) (28). These infections frequently persist (29,30), and it has been observed that older age and a history of sunburn are associated with an elevated risk of persistent β -HPV infection (30,31). Furthermore, a significant association between the number of β -HPVs detected in eyebrow hair follicles and an increased risk of cSCC was reported among OTRs from Europe (the Netherlands, the United Kingdom, France and Italy) (32).

Moreover, individuals with concordant β -HPV DNA in plucked eyebrow hairs and serologic tests had a significantly increased risk of developing SCC (33).

Among immunosuppressed OTRs, the occurrence of multiple cutaneous HPV infections is common (34), but high viral loads were shown to be associated with an increased risk of SCC development, with total load seemingly more important than the individual load of any specific type (32). It should be noted, however, that β -HPV is more highly prevalent in skin wart biopsies from OTRs than in either normal skin or plucked hairs from these patients (29,35-37). Serological studies have also demonstrated that seroconversion to β -HPV increases with age (38,39) and have revealed a positive epidemiological association between β -HPV seroreactivity and cSCC development (40), even though not every infection is accompanied by a detectable or relevant seroresponse.

Nevertheless, given the high incidence of cSCCs in OTRs, identifying a clear link between β -HPV infection and cSCC would have important implications for therapy and prevention (41,42). Therefore, more recent case-control studies are ongoing, focusing on the association of cutaneous HPV in the early stages of NMSC carcinogenesis in immunosuppressed individuals. However, the data are inconclusive (42-46).

Prevalence and distribution of cutaneous HPV's among immunocompetent individuals

Cutaneous β - and γ -HPV DNA can be detected beginning in early infancy and may be detected in 70% of children by 4 years of age (47). Additionally, β -HPV types detected on parents are more commonly found on their babies (47,48). Viral transmission seems to occur inevitably through direct skin contact (10,49-51), and these viruses have been suggested to be commensal to humans (38,52,53) (Table 2).

Table 2 - Select studies on the prevalence and distribution of cutaneous HPV's among immunocompetent individuals.

Year	Author (s) (reference number)	Data
1997	Boxman et al. (35)	Cutaneous HPV's detection in eyebrow hairs seems to reflect infections in other parts of the body (useful in epidemiological studies).
2000	Antonsson et al. (6)	Sun exposure and history of skin cancer are risk factors associated to β -HPV's detection in IC individuals.
2000	Harwood et al. (36)	Cutaneous HPV's prevalence was higher among individuals who reported ever having skin lesions.
2000 2003 2007 2009 2009	Antonsson et al. (6) Struijk et al. (56) Hazard et al. (30) Weissenborn et al. (48) de Koning et al. (55)	β -HPV DNA detection and seroprevalence increases with age.
2003	Antonsson et al. (47)	The presence of cutaneous β - and γ -HPV's DNA is observed since early infancy.
2003 2009	Antonsson et al. (47) Weissenborn et al. (48)	β -HPV's types detected on parents are also more commonly found in their babies.
2003 2010 2014	Antonsson et al. (52) Antonsson et al. (38) Bzhalava et al. (53)	β - and γ -HPV's may be commensal to humans.
2004	Termorshuizen et al. (24)	Severe sunburns are associated with the presence of β -HPV DNA.
2004	Stockfleth et al. (60)	HPV's 5 and 8 were the most frequently found in premalignant lesions, SCC and BCC.
2004 2006 2011 2017 2017 2017	Smith et al. (75) Fakhry et al. (73) Termine et al. (77) Hampras et al. (74) Steinau et al. (76) Nunes et al. (10)	Simultaneous oral-genital type-specific β -HPV infections are relatively rare.

**Table 2 - Continued.**

Year	Author (s) (reference number)	Data
2005 2011	Weissenborn et al. (46) Arron et al. (43)	Higher viral loads are detected within pre-malignant skin lesions as compared to SCC.
2007	Forslund et al. (23)	Most viruses detected on the external skin surface may reflect HPV deposition.
2007 2009	Köhler et al. (54) de Koning et al. (55)	The prevalence of β -HPVs DNA in plucked hairs from different body sites of IC individuals is approximately 90%.
2008	Patel et al. (59)	β -HPVs may play a role in the pathogeny of NMSC also in healthy individuals.
2008 2017 2017	Feltkamp et al. (49) Moscicki et al. (51) Nunes et al. (10)	Viral transmission seems to occur through direct skin contact.
2009	Bouvard et al. (15)	Epidemiological evidence concerning the association between specific β - and γ -HPVs and the development of skin cancer in IC is inconclusive.
2009	Weissenborn et al. (48)	UV radiation may be a putative viral detection-related risk factor.
2010 2010	Antonsson et al. (38) Iannaccone et al. (62)	High overall seropositivity (>90%) to at least one viral β -HPV is observed within healthy individuals.
2010 2011	Michael et al. (63) Proby et al. (33)	Low overall seropositivity to at least one viral β -HPV is observed within healthy individuals.
2011	Proby et al. (33)	β -HPV types most commonly detected in the skin also have the highest seroprevalence.
2011 2013 2013 2013 2013 2014 2014 2015 2015 2016 2017	Bottalico et al. (7) Forslund et al. (8) Pierce Campbell et al. (68) Sichero et al. (65) Paolini et al. (70) Hampras et al. (31) Sichero et al. (66) Sichero et al. (69) Donà et al. (67) Nunes et al. (9) Moscicki et al. (51)	Analysis of β - and γ -HPVs DNA and antibodies prevalence among series of samples (anogenital, oral, skin, nasal cavity from women and men).
2013	Neale et al. (32)	Cutaneous HPVs target the hair follicle bulge, which is probably the reservoir of these viruses.
2013	Sichero et al. (65)	The majority of male genital samples could not be classified using technologies widely used for typing of α -HPVs.
2013 2016	Pierce Campbell et al. (68) Rahman et al. (71)	Male external genital lesions (EGL) are not associated to β -HPVs infections.
2013 2014	Sichero et al. (65) Sichero et al. (66)	Most samples from the male genitals were positive for multiples cutaneous HPV DNA.
2013 2014 2015	Sichero et al. (65) Sichero et al. (66) Sichero et al. (69)	The detection of β -HPVs in one anatomic site may also represent deposition of virions shed from other anatomic sites.
2013 2014 2015 2015 2016	Sichero et al. (65) Sichero et al. (66) Donà et al. (67) Torres et al. (72) Nunes et al. (9)	The detection of cutaneous HPVs DNA is not associated to sexual risk factors. Other routes of transmission such as autoinoculation and non-penetrative sexual activities could be associated.
2016	Chahoud et al. (44)	Analyses of association between the detection of antibodies to β -HPVs with SCC risk development.
2017	Hampras et al. (74)	The occurrence of concordant β -HPV infections seem to be higher across keratinized tissues than across mucosal sites.
2017	Moscicki et al. (51)	The transmission rate of β -HPVs between anogenital sites from men-to-women and women-to-men was similar, suggesting these are sexually transmitted.

β -HPVs are widespread in immunocompetent (IC) individuals within the general population: when plucked hairs from different body sites are tested, the prevalence is approximately 90% (54,55). It is believed that cutaneous HPVs target the hair follicle bulge, which is probably the reservoir of these viruses (32). Therefore, eyebrow hairs have served as an easily obtained material for marker analysis in several epidemiological studies and seem to reflect infection in other parts of the body (35).

Advanced age has been shown to be the most important factor influencing the presence of β -HPV DNA in IC individuals (56,57). Furthermore, sun exposure and a history of skin cancer are risk factors associated with β -HPV detection in these individuals (6). For OTRs, some studies have investigated the prevalence of cutaneous HPVs among IC individuals of different ethnicities and residing under different climate conditions (52). It was reported that the prevalence of HPV DNA was lower in samples from Zambia than in those



collected in Sweden ($p < 0.01$) and Bangladesh ($p < 0.05$) (52). β -HPV prevalence and distribution studies have shown that viral positivity was, on average, higher on the forehead (36%) and back of the hand (38%) than on the buttocks (26%), indicating that UV radiation may be a putative risk factor for viral infection, even though sun exposure data were not collected in this specific study (48). In fact, severe sunburns have been associated with the presence of β -HPV DNA (24). Second-degree burns and repetitive sunburns, with skin regeneration of the underlying capillary bulb, may result in the amplification of β -HPV DNA by activating the HPV life cycle (24). In fact, the risk for SCC development among Australian or Netherlander IC individuals is higher for those in which β -HPV DNA was detected at high loads in plucked eyebrow hairs (32,56,58).

β -HPV persistence was more commonly observed in adults (92%) than in children (66%), and although multiple β -HPVs can infect persistently, no specific type seems to predominate in such infections (48). Nevertheless, it must be highlighted that the prevalence of β -HPVs significantly decreases after tape stripping, indicating that only a small number of epithelial cells are in fact infected, and most of the detected viruses may reflect deposition throughout the external skin surface (23).

The IARC recognizes the need for further research on cutaneous HPVs to better understand the widespread distribution of these viruses. β -HPVs may also play a role in the pathogeny of NMSC in healthy individuals (59); however, to date, epidemiological evidence is inconclusive concerning the association between specific β - and γ -HPVs and the development of skin cancer in IC individuals (14,15).

Regarding the HPV status in NMSCs from immunosuppressed and IC individuals, viral prevalence was higher in the former group for all lesion types analyzed: premalignant lesions (88% among immunosuppressed *vs* 54% among IC), SCC (84% *vs* 27%) and basal cell carcinoma (BCC, 75% *vs* 36%) (36). Nevertheless, the prevalence and spectrum of HPV types detected within the two populations were equivalent among premalignant lesions, SCC and BCC, and HPVs 5 and 8 were the most frequently identified types (60). Interestingly, β -HPV prevalence was reported to be higher in premalignant AK than in cSCC, and real-time PCR analysis indicated higher viral loads in premalignant lesions (46) than in SCC, in which viral load rarely reaches the level of one viral copy per cell (43,46). This scenario is compatible with a carcinogenic role for HPV at the early stages of skin carcinogenesis. Because cSCC most commonly develops in sun-exposed anatomical sites, it is reasonable to suppose that UV radiation may impede HPV antigen presentation by suppressing local cell immunity (61). In addition, several studies have suggested the importance of β -HPVs as cofactors to UV radiation in the development of SCC by facilitating the accumulation of UV-induced mutations, which can ultimately lead to cell transformation.

In addition to HPV DNA detection, the detection of antibodies to β -HPVs and their association with SCC risk development have been evaluated in several studies (44). Nevertheless, overall, serological studies show considerable heterogeneity in the results: whereas high overall seropositivity (>90%) to at least one viral β -HPV has been described (38,62), a lower prevalence is observed in other studies (33,63). The divergence in the data obtained in these studies could be attributed not only to differences in serological methods but also to the range of cutaneous HPVs tested. With β -HPV DNA detection, the seroprevalence has been shown to increase with age (6,30,55). Notably, β -HPV

types most commonly detected in the skin have the highest seroprevalence worldwide (33).

Due to the wide distribution of cutaneous HPVs in the skin, several groups, including ours, have recently focused on investigating the distribution of cutaneous HPVs in other anatomical sites, including the anogenital area. Within the HIM (HPV Infection in Men) cohort study (64), we initially observed that most of the ~15% of male genital samples that could not be classified with widely used α -HPV typing technologies harbored β - and γ -HPVs, as evidenced by using a PCR sequencing protocol (65). We further observed that most samples were positive for HPV DNA of multiple cutaneous types using a sensitive Luminex-based methodology, suggesting that the former protocol could underestimate the true prevalence of cutaneous β - and γ -HPVs in the male genital region (65,66). In order to better understand the prevalence and distribution of cutaneous HPVs, we and others further analyzed β - and γ -HPV DNA and antibodies by Luminex methodology in a series of samples obtained from the anogenital region of both men and women (9,51,65-69), the oral cavity (7,9,70), the skin (31), and the nasal cavity (8). Taken together, these studies corroborate that cutaneous HPVs are ubiquitously disseminated throughout healthy skin and may be an intrinsic part of the commensal flora.

We further observed that male external genital lesions (EGLs) are not associated with β -HPV infections (68,71) and that the detection of DNA from these viruses is not associated with sexual risk factors, indicating other routes of transmission, such as autoinoculation and nonpenetrative sexual activities (9,65-67,72). Alternatively, the detection of β -HPVs at one anatomic site may indicate the deposition of virions shed from other anatomic sites (50,65,66,69). Nevertheless, it was recently reported that among heterosexual couples, the transmission rate of β -HPVs between anogenital sites was 15.9 per 100 person-months from men-to-women, with a similar risk for women-to-men transmission, suggesting that β -HPVs can be sexually transmitted (51). Lastly, the few reports in which the oral and anogenital regions were analyzed concurrently indicated that simultaneous oral-genital type-specific β -HPV infections are relatively rare (10,73-77) but seem to be higher across keratinized tissues than across mucosal sites (74). Nevertheless, given the large number of samples and β -HPV types analyzed, it is unlikely that the same HPV type will be found at a distant skin site by chance.

Functional analysis of cutaneous HPVs

HPVs are small, nonenveloped viruses with circular double-stranded DNA of approximately 8000 bp. The viral genome is divided into three regions: the long control region (LCR) contains cis-responsive elements for viral and cellular proteins that regulate viral gene expression and replication; the early region (E), which encodes proteins crucial to viral transcription and replication; and the late region (L), which encodes the viral capsid structural proteins (78-80). Although the viral genome structure and organization are highly conserved among HPVs, the LCR of β -HPVs is shorter than that of α -HPVs, and the E5 gene is absent from the β -HPV genome (81-83).

The HPV life cycle is tightly associated with the differentiation of the stratified squamous epithelium. HPV infection begins with entrance of the virus into the basal layer of the epithelia due to microtrauma (79,84,85). At least for high-risk α -HPV-infected tissues, the differentiation process is altered




Figure 2 - Cellular targets of the E6 and E7 proteins from specific β -HPVs.

Table 3 - Highlights regarding the epidemiology and biology of cutaneous human papillomavirus in immunosuppressed and immunocompetent individuals.

- β - and γ -HPVs, in addition to few α -HPVs are originally designated "cutaneous types".
- The IARC classified β -HPVs 5 and 8 as possible etiological agents of skin SCC in EV individuals.
- OTRs not only have a higher susceptibility to β -HPVs but further attain an ~100 fold increased risk of developing NMSC as compared to the general population.
- Among OTRs and IC individuals cutaneous HPVs are ubiquitously spread throughout the body and may be an intrinsic part of the commensal flora.
- Although cumulative sun exposure is the major risk factor for NMSC, studies points towards β -HPV infections as co-factors in skin SCC in association with UV radiation.
- β -HPVs most probably play a role in the initiation of skin SCC rather than in the maintenance of the transformed phenotype.
- Function analysis of E6 and E7 proteins of specific cutaneous HPVs indicate that the biology involved in β -HPV mediated skin carcinogenesis differ from that induced by high-risk α -HPV types.

by expression of the E6 and E7 viral oncoproteins, which interact principally with the TP53 and pRb suppressor proteins, respectively, but also interact with a broad spectrum of other cellular proteins, altering the biological properties of the host cell (79,83,86-89).

As previously discussed in this review, β -HPVs most likely play a role in the initiation of cSCC rather than in the maintenance of the transformed phenotype (14). It is hypothesized that β -HPV infections destabilize the host genome, allowing tumors to further develop in the absence of the viral genome (90-92). Some studies have provided a mechanistic comprehension of how these viruses induce carcinogenesis and have indicated that the biology involved in β -HPV-mediated skin carcinogenesis differs from that induced by high-risk α -HPVs (80,91,93,94).

Studies have shown that β -HPVs 38 and 49 are able to immortalize primary human keratinocytes, whereas HPVs 10, 14, 22, 23, 24 and 36 do not have this ability (95-97). It has also been reported that transgenic mice expressing the HPV 38 E6 and E7 proteins under control of a keratinocyte-specific promoter exhibit epidermal hyperplasia and are susceptible to the development of cutaneous tumors promoted by chemical carcinogens and UV radiation (98-101). Although the E6 protein from HPVs 8, 24 and 38 binds *in vitro* to E6-AP (E6-associated protein), p53 degradation was observed in the presence of only HPV 49 (95-97,102) (Figure 2). The E6 protein from β -HPVs 5, 8, and 38 attenuates p53 phosphorylation and ubiquitination in response to UV exposure,

resulting in less efficient repair of damaged cellular DNA (90,96). Additionally, HPV 38 induces telomerase by a mechanism dependent on E6-AP (95). HPV 38 E6 also alters the capacity of p53 to activate proteins involved in apoptosis and suppress proliferation by inducing the accumulation of Δ Np73, a p53 isoform that antagonizes p53 (96). The E6 protein from HPVs 5, 8 and 38 was shown to bind p300, preventing p53 acetylation and p53 -induced repair and transcriptional transactivation, thus contributing to the accumulation of mutations and chromosomal abnormalities (90). Furthermore, as with α -HPV, the E6 protein of some β -HPVs induces BAK degradation, thus preventing the release of pro-apoptotic mitochondrial factors (103). The interaction of E6 with E6-AP is required not only for BAK degradation but also for hTERT (human telomerase reverse transcriptase) induction (104,105).

The E7 proteins from the cutaneous HPVs 154, 22, 23, 24, 36, 38 and 49 bind *in vitro* to pRb but are unable to induce pRb degradation when expressed in human keratinocytes (97,106-108) (Figure 2). Nevertheless, in human keratinocytes transduced with HPV 38 and 39 E6 and E7 proteins, E2F-induced transcription is likely activated because these viral proteins induce pRb hyperphosphorylation (80,95,97).

Although most research on oncogenic potential and disease association has focused on α -HPVs, there is interest in identifying a role of non- α HPV types in the pathogenesis of benign and malignant lesions (Table 3). Challenges in finding relevant associations between cutaneous HPV infection and NMSC



Table 4 - Challenges/critical open questions regarding the epidemiology and biology of cutaneous human papillomavirus in immunosuppressed and immunocompetent individuals.

- Are cutaneous HPVs associated to the development of non-melanoma skin cancer among immunosuppressed and immunocompetent individuals?
- Why are cutaneous HPVs more diverse than their mucosal counterparts?
- Do cutaneous HPVs contribute to carcinogenesis associated to other carcinogens at skin and non-skin sites?
- Do cutaneous HPVs contribute to high-risk mucosal carcinogenesis in cases of co-infections?
- Why are cutaneous HPVs more prevalent in precursor lesions compared to malignant lesions?
- Why do specific β -HPVs deregulate fundamental cellular events intimately linked to transformation *in vitro*, but do not cause cancer in humans?

development include the multiplicity and ubiquity of these viruses throughout the human body, the high probability of viral transmission (including autoinoculation), and differences observed in the carcinogenic potential of individual β -HPVs (Table 4). The last IARC monograph (100B) was unable to identify consistent epidemiological evidence for an etiological role attributable to any specific cutaneous HPV type or species in NMSC development. In addition, biological mechanisms explaining the oncogenicity of these viruses have not been fully elucidated.

AUTHORS' CONTRIBUTIONS

Nunes EM, Talpe-Nunes V and Sichero L critically discussed and wrote the manuscript.

REFERENCES

- IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Biological agents. Volume 100 B. A review of human carcinogens. IARC Monogr Eval Carcinog Risks Hum. 2012;100(Pt B):1-441.
- de Villiers EM, Fauquet C, Broker TR, Bernard HU, zur Hausen H. Classification of papillomaviruses. *Virology*. 2004;324(1):17-27, <http://dx.doi.org/10.1016/j.virol.2004.03.033>.
- de Villiers EM. Cross-roads in the classification of papillomaviruses. *Virology*. 2013;445(1-2):2-10, <http://dx.doi.org/10.1016/j.virol.2013.04.023>.
- Chouhy D, Bolatti EM, Pérez GR, Giri AA. Analysis of the genetic diversity and phylogenetic relationships of putative human papillomavirus types. *J Gen Virol*. 2013;94(Pt 11):2480-8, <http://dx.doi.org/10.1099/vir.0.055137-0>.
- IARC Working Group. IARC Monograph on the Evaluation of Carcinogenic Risk to Humans: Human Papillomaviruses. Lyon: International Agency for Research on Cancer; 2007.
- Antonsen A, Forslund O, Ekberg H, Sterner G, Hansson BG. The ubiquity and impressive genomic diversity of human skin papillomaviruses suggest a commensal nature of these viruses. *J Virol*. 2000;74(24):11636-41, <http://dx.doi.org/10.1128/JVI.74.24.11636-11641.2000>.
- Bottalico D, Chen Z, Dunne A, Ostolozza J, McKinney S, Sun C, et al. The oral cavity contains abundant known and novel human papillomaviruses from the Betapapillomavirus and Gammapapillomavirus genera. *J Infect Dis*. 2011;204(5):787-92, <http://dx.doi.org/10.1093/infdis/jir383>.
- Forslund O, Johansson H, Madsen KG, Kofoed K. The nasal mucosa contains a large spectrum of human papillomavirus types from the Betapapillomavirus and Gammapapillomavirus genera. *J Infect Dis*. 2013;208(8):1335-41, <http://dx.doi.org/10.1093/infdis/jit326>.
- Nunes EM, Sudenga SL, Gheit T, Tommasino M, Baggio ML, Ferreira S, et al. Diversity of beta-papillomavirus at anogenital and oral anatomic sites of men: The HIM Study. *Virology*. 2016;495:33-41, <http://dx.doi.org/10.1016/j.virol.2016.04.031>.
- Nunes EM, López RVM, Sudenga SL, Gheit T, Tommasino M, Baggio ML, et al. Concordance of Beta-papillomavirus across anogenital and oral anatomic sites of men: The HIM Study. *Virology*. 2017;510:55-9, <http://dx.doi.org/10.1016/j.virol.2017.07.006>.
- Sichero L, El-Zein M, Nunes EM, Ferreira S, Franco EL, Villa LL; Ludwig-McGill Cohort Study. Cervical Infection with Cutaneous Beta and Mucosal Alpha Papillomaviruses. *Cancer Epidemiol Biomarkers Prev*. 2017;26(8):1312-20, <http://dx.doi.org/10.1158/1055-9965.EPI-17-0081>.
- Lewandowsky F, Lutz W. Ein Fall einer bisher nicht beschriebenen Hauterkrankung (Epidermodysplasia verruciformis). *Archiv für Dermatologie und Syphilis*. 1922;141(2):193-203.
- Jablonska S, Dabrowski J, Jakubowicz K. Epidermodysplasia verruciformis as a model in studies on the role of papovaviruses in oncogenesis. *Cancer Res*. 1972;32(3):583-9.
- Pfister H. Chapter 8: Human papillomavirus and skin cancer. *J Natl Cancer Inst Monogr*. 2003;(31):52-6, <http://dx.doi.org/10.1093/oxfordjournals.jncimonographs.a003483>.
- Bouvard V, Baan R, Straif K, Grosse Y, Secretan B, El Ghissassi F, et al. WHO International Agency for Research on Cancer Monograph Working Group. A review of human carcinogens-Part B: biological agents. *Lancet Oncol*. 2009;10(4):321-2, [http://dx.doi.org/10.1016/S1470-2045\(09\)70096-8](http://dx.doi.org/10.1016/S1470-2045(09)70096-8).
- Birkeland SA, Storm HH, Lamm LU, Barlow L, Blohmé I, Forsberg B, et al. Cancer risk after renal transplantation in the Nordic countries, 1964-1986. *Int J Cancer*. 1995;60(2):183-9, <http://dx.doi.org/10.1002/ijc.2910600209>.
- Lindelöf B, Sigurgeirsson B, Gäbel H, Stern RS. Incidence of skin cancer in 5356 patients following organ transplantation. *Br J Dermatol*. 2000;143(3):513-9.
- Hardie IR, Strong RW, Hartley LC, Woodruff PW, Clunie GJ. Skin cancer in Caucasian renal allograft recipients living in a subtropical climate. *Surgery*. 1980;87(2):177-83.
- Hoxtell EO, Mandel JS, Murray SS, Schuman LM, Goltz RW. Incidence of skin carcinoma after renal transplantation. *Arch Dermatol*. 1977;113(4):436-8, <http://dx.doi.org/10.1001/archderm.1977.01640040044005>.
- Mullen DL, Silverberg SG, Penn I, Hammond WS. Squamous cell carcinoma of the skin and lip in renal homograft recipients. *Cancer*. 1976;37(2):729-34, [http://dx.doi.org/10.1002/1097-0142\(197602\)37:2<729::AID-CNCR2820370220>3.0.CO;2-R](http://dx.doi.org/10.1002/1097-0142(197602)37:2<729::AID-CNCR2820370220>3.0.CO;2-R).
- Barr BB, Benton EC, McLaren K, Bunney MH, Smith IW, Blessing K, et al. Human papilloma virus infection and skin cancer in renal allograft recipients. *Lancet*. 1989;1(8630):124-9, [http://dx.doi.org/10.1016/S0140-6736\(89\)91143-4](http://dx.doi.org/10.1016/S0140-6736(89)91143-4).
- Sbano E, Andreassi L, Fimiani M, Valentino A, Baiocchi R. DNA-repair after UV-irradiation in skin fibroblasts from patients with actinic keratosis. *Arch Dermatol Res*. 1978;262(1):55-61, <http://dx.doi.org/10.1007/BF00455573>.
- Forslund O, Iftner T, Andersson K, Lindelöf B, Hradil E, Nordin P, et al. Cutaneous human papillomaviruses found in sun-exposed skin: Beta-papillomavirus species 2 predominates in squamous cell carcinoma. *J Infect Dis*. 2007;196(6):876-83, <http://dx.doi.org/10.1086/521031>.
- Termorshuizen F, Feltkamp MC, Struijk L, de Grijl FR, Bavinck JN, van Loveren H. Sunlight exposure and (sero) prevalence of epidermodysplasia verruciformis-associated human papillomavirus. *J Invest Dermatol*. 2004;122(6):1456-62, <http://dx.doi.org/10.1111/j.0022-202X.2004.22617.x>.
- Pfister H. Papillomaviruses. In: Salzman NP, Howley PM, editors. *The Papovaviridae. The Viruses*. Boston: Springer; 1987. p. 1-38.
- Koranda FC, Dehmel EM, Kahn G, Penn I. Cutaneous complications in immunosuppressed renal homograft recipients. *JAMA*. 1974;229(4):419-24, <http://dx.doi.org/10.1001/jama.1974.03230420031020>.
- Boyle J, MacKie RM, Briggs JD, Junor BJ, Aitchison TC. Cancer, warts, and sunshine in renal transplant patients. A case-control study. *Lancet*. 1984;1(8379):702-5, [http://dx.doi.org/10.1016/S0140-6736\(84\)92221-9](http://dx.doi.org/10.1016/S0140-6736(84)92221-9).
- Nindl I, Köhler A, Gottschling M, Förschner T, Lehmann M, Meijer CJ, et al. Extension of the typing in a general-primer-PCR reverse-line-blotting system to detect all 25 cutaneous beta human papillomaviruses. *J Virol Methods*. 2007;146(1-2):1-4, <http://dx.doi.org/10.1016/j.jviromet.2007.05.022>.
- Berkhout RJ, Bouwes Bavinck JN, ter Schegget J. Persistence of human papillomavirus DNA in benign and (pre)malignant skin lesions from renal transplant recipients. *J Clin Microbiol*. 2000;38(6):2087-96.
- Hazard K, Karlsson A, Andersson K, Ekberg H, Dillner J, Forslund O. Cutaneous human papillomaviruses persist on healthy skin. *J Invest Dermatol*. 2007;127(1):116-9, <http://dx.doi.org/10.1038/sj.jid.5700570>.
- Hampras SS, Giuliano AR, Lin HY, Fisher KJ, Abrahamsen ME, Sirak BA, et al. Natural history of cutaneous human papillomavirus (HPV) infection in men: the HIM study. *PLoS One*. 2014;9(9):e104843, <http://dx.doi.org/10.1371/journal.pone.0104843>.
- Neale RE, Weissenborn S, Abeni D, Bavinck JN, Euvrard S, Feltkamp MC, et al. Human papillomavirus load in eyebrow hair follicles and risk of cutaneous squamous cell carcinoma. *Cancer Epidemiol Biomarkers Prev*. 2013;22(4):719-27, <http://dx.doi.org/10.1158/1055-9965.EPI-12-0917-T>.
- Proby CM, Harwood CA, Neale RE, Green AC, Euvrard S, Naldi L, et al. A case-control study of betapapillomavirus infection and cutaneous



- squamous cell carcinoma in organ transplant recipients. *Am J Transplant.* 2011;11(7):1498-508, <http://dx.doi.org/10.1111/j.1600-6143.2011.03589.x>.
34. Harwood CA, Suretheran T, Sasieni P, Proby CM, Bordea C, Leigh IM, et al. Increased risk of skin cancer associated with the presence of epidermodysplasia verruciformis human papillomavirus types in normal skin. *Br J Dermatol.* 2004;150(5):949-57, <http://dx.doi.org/10.1111/j.1365-2133.2004.05847.x>.
35. Boxman IL, Berkhout RJ, Mulder LH, Wolkers MC, Bouwes Bavinck JN, Vermeer BJ, et al. Detection of human papillomavirus DNA in plucked hairs from renal transplant recipients and healthy volunteers. *J Invest Dermatol.* 1997;108(5):712-5, <http://dx.doi.org/10.1111/1523-1747.ep12292090>.
36. Harwood CA, Suretheran T, McGregor JM, Spink PJ, Leigh IM, Breuer J, et al. Human papillomavirus infection and non-melanoma skin cancer in immunosuppressed and immunocompetent individuals. *J Med Virol.* 2000;61(3):289-97, [http://dx.doi.org/10.1002/1096-9071\(200007\)61:3<289::AID-JMV2>3.0.CO;2-Z](http://dx.doi.org/10.1002/1096-9071(200007)61:3<289::AID-JMV2>3.0.CO;2-Z).
37. Meyer T, Arndt R, Nindl I, Ulrich C, Christophers E, Stockfleth E. Association of human papillomavirus infections with cutaneous tumors in immunosuppressed patients. *Transpl Int.* 2003;16(3):146-53, <http://dx.doi.org/10.1111/j.1432-2277.2003.tb00278.x>.
38. Antonsson A, Green AC, Mallitt KA, O'Rourke PK, Pandeya N, Pawlita M, et al. Prevalence and stability of antibodies to 37 human papillomavirus types-a population-based longitudinal study. *Virology.* 2010;407(1):26-32, <http://dx.doi.org/10.1016/j.virol.2010.07.046>.
39. Michael KM, Waterboer T, Sehr P, Rother A, Reidel U, Boeing H, et al. Seroprevalence of 34 human papillomavirus types in the German general population. *PLoS Pathog.* 2008;4(6):e1000091, <http://dx.doi.org/10.1371/journal.ppat.1000091>.
40. Feltkamp MC, Broer R, di Summa FM, Struijk L, van der Meijden E, Verlaan BP, et al. Seroreactivity to epidermodysplasia verruciformis-related human papillomavirus types is associated with nonmelanoma skin cancer. *Cancer Res.* 2003;63(10):2695-700.
41. Bhat P, Mattarollo SR, Gosmann C, Frazer IH, Leggatt GR. Regulation of immune responses to HPV infection and during HPV-directed immunotherapy. *Immunol Rev.* 2011;239(1):85-98, <http://dx.doi.org/10.1111/j.1600-065X.2010.00966.x>.
42. Aldabagh B, Angeles JG, Cardones AR, Arron ST. Cutaneous squamous cell carcinoma and human papillomavirus: is there an association? *Dermatol Surg.* 2013;39(1 Pt 1):1-23, <http://dx.doi.org/10.1111/j.1524-4725.2012.02558.x>.
43. Arron ST, Ruby JG, Dybbro E, Ganem D, Derisi JL. Transcriptome sequencing demonstrates that human papillomavirus is not active in cutaneous squamous cell carcinoma. *J Invest Dermatol.* 2011;131(8):1745-53, <http://dx.doi.org/10.1038/jid.2011.91>.
44. Chahoud J, Semaan A, Chen Y, Cao M, Rieber AG, Rady P, et al. Association Between β -Genus Human Papillomavirus and Cutaneous Squamous Cell Carcinoma in Immunocompetent Individuals-A Meta-analysis. *JAMA Dermatol.* 2016;152(12):1354-64, <http://dx.doi.org/10.1001/jamadermatol.2015.4530>.
45. Rollison DE, Pawlita M, Giuliano AR, Iannacone MR, Sondak VK, Messina JL, et al. Measures of cutaneous human papillomavirus infection in normal tissues as biomarkers of HPV in corresponding non-melanoma skin cancers. *Int J Cancer.* 2008;123(10):2337-42, <http://dx.doi.org/10.1002/ijc.23795>.
46. Weissenborn SJ, Nindl I, Purdie K, Harwood C, Proby C, Breuer J, et al. Human papillomavirus-DNA loads in actinic keratoses exceed those in non-melanoma skin cancers. *J Invest Dermatol.* 2005;125(1):93-7, <http://dx.doi.org/10.1111/j.0022-202X.2005.23733.x>.
47. Antonsson A, Karanfilovska S, Lindqvist PG, Hansson BG. General acquisition of human papillomavirus infections of skin occurs in early infancy. *J Clin Microbiol.* 2003;41(6):2509-14, <http://dx.doi.org/10.1128/JCM.41.6.2509-2514.2003>.
48. Weissenborn SJ, De Koning MN, Wieland U, Quint WG, Pfister HJ. Intrafamilial transmission and family-specific spectra of cutaneous betapapillomaviruses. *J Virol.* 2009;83(2):811-6, <http://dx.doi.org/10.1128/JVI.01338-08>.
49. Feltkamp MC, de Koning MN, Bavinck JN, Ter Schegget J. Betapapillomaviruses: innocent bystanders or causes of skin cancer. *J Clin Virol.* 2008;43(4):353-60, <http://dx.doi.org/10.1016/j.jcv.2008.09.009>.
50. Liu Z, Rashid T, Nyitray AG. Penises not required: a systematic review of the potential for human papillomavirus horizontal transmission that is non-sexual or does not include penile penetration. *Sex Health.* 2016;13(1):10-21.
51. Moscicki AB, Ma Y, Gheit T, McKay-Chopin S, Farhat S, Widdice LE, et al. Prevalence and Transmission of Beta and Gamma Human Papillomavirus in Heterosexual Couples. *Open Forum Infect Dis.* 2017;4(1): ofw216, <http://dx.doi.org/10.1093/ofid/ofw216>.
52. Antonsson A, Erfurt C, Hazard K, Holmgren V, Simon M, Kataoka A, et al. Prevalence and type spectrum of human papillomaviruses in healthy skin samples collected in three continents. *J Gen Virol.* 2003;84(Pt 7):1881-6, <http://dx.doi.org/10.1099/vir.0.18836-0>.
53. Bzhalava D, Mühr LS, Lagheden C, Ekström J, Forslund O, Dillner J, et al. Deep sequencing extends the diversity of human papillomaviruses in human skin. *Sci Rep.* 2014;4:5807, <http://dx.doi.org/10.1038/srep05807>.
54. Köhler A, Forschner T, Meyer T, Ulrich C, Gottschling M, Stockfleth E, et al. Multifocal distribution of cutaneous human papillomavirus types in hairs from different skin areas. *Br J Dermatol.* 2007;156(5):1078-80, <http://dx.doi.org/10.1111/j.1365-2133.2007.07809.x>.
55. de Koning MN, Weissenborn SJ, Abeni D, Bouwes Bavinck JN, Euvrard S, Green AC, et al. Prevalence and associated factors of betapapillomavirus infections in individuals without cutaneous squamous cell carcinoma. *J Gen Virol.* 2009;90(Pt7):1611-21, <http://dx.doi.org/10.1099/vir.0.010017-0>.
56. Struijk L, Bouwes Bavinck JN, Wanningen P, van der Meijden E, Westendorp RG, Ter Schegget J, et al. Presence of human papillomavirus DNA in plucked eyebrow hairs is associated with a history of cutaneous squamous cell carcinoma. *J Invest Dermatol.* 2003;121(6):1531-5, <http://dx.doi.org/10.1046/j.1523-1747.2003.12632.x>.
57. Bouwes Bavinck JN, Plasmeijer EI, Feltkamp MC. Beta-papillomavirus infection and skin cancer. *J Invest Dermatol.* 2008;128(6):1355-8, <http://dx.doi.org/10.1038/jid.2008.123>.
58. Iannacone MR, Gheit T, Pfister H, Giuliano AR, Messina JL, Fenske NA, et al. Case-control study of genus-beta human papillomaviruses in plucked eyebrow hairs and cutaneous squamous cell carcinoma. *Int J Cancer.* 2014;134(9):2231-44, <http://dx.doi.org/10.1002/ijc.28552>.
59. Patel AS, Karagas MR, Perry AE, Nelson HH. Exposure profiles and human papillomavirus infection in skin cancer: an analysis of 25 genus beta-types in a population-based study. *J Invest Dermatol.* 2008;128(12):2888-93, <http://dx.doi.org/10.1038/jid.2008.162>.
60. Stockfleth E, Nindl I, Sterry W, Ulrich C, Schmook T, Meyer T. Human papillomaviruses in transplant-associated skin cancers. *Dermatol Surg.* 2004;30(4 Pt 2):604-9, <http://dx.doi.org/10.1111/j.1524-4725.2004.00144.x>.
61. Norval M. The mechanisms and consequences of ultraviolet-induced immunosuppression. *Prog Biophys Mol Biol.* 2006;92(1):108-18, <http://dx.doi.org/10.1016/j.pbiomolbio.2006.02.009>.
62. Iannacone MR, Michael KM, Giuliano AR, Waterboer T, Pawlita M, Rollison DE. Risk factors for cutaneous human papillomavirus seroreactivity among patients undergoing skin cancer screening in Florida. *J Infect Dis.* 2010;201(5):760-9, <http://dx.doi.org/10.1086/650466>.
63. Michael KM, Waterboer T, Pfister H, Gariglio M, Majewski S, Favre M, et al. Seroreactivity of 38 human papillomavirus types in epidermodysplasia verruciformis patients, relatives, and controls. *J Invest Dermatol.* 2010;130(3):841-8, <http://dx.doi.org/10.1038/jid.2009.356>.
64. Giuliano AR, Lazcano-Ponce E, Villa LL, Flores R, Salmeron J, Lee JH, et al. The human papillomavirus infection in men study: human papillomavirus prevalence and type distribution among men residing in Brazil, Mexico, and the United States. *Cancer Epidemiol Biomarkers Prev.* 2008;17(8):2036-43, <http://dx.doi.org/10.1158/1055-9965.EPI-08-0151>.
65. Sichero L, Pierce Campbell CM, Ferreira S, Sobrinho JS, Luiza Baggio M, Galan L, et al. Broad HPV distribution in the genital region of men from the HPV infection in men (HIM) study. *Virology.* 2013;443(2):214-7, <http://dx.doi.org/10.1016/j.virol.2013.04.024>.
66. Sichero L, Pierce Campbell CM, Fulp W, Ferreira S, Sobrinho JS, Baggio M, et al. High genital prevalence of cutaneous human papillomavirus DNA on male genital skin: the HPV Infection in Men Study. *BMC Infect Dis.* 2014;14:677, <http://dx.doi.org/10.1186/s12879-014-0677-y>.
67. Donà MG, Gheit T, Latini A, Benevolo M, Torres M, Smelov V, et al. Alpha, beta and gamma Human Papillomaviruses in the anal canal of HIV-infected and uninfected men who have sex with men. *J Infect.* 2015;71(1):74-84, <http://dx.doi.org/10.1016/j.jinf.2015.02.001>.
68. Pierce Campbell CM, Messina JL, Stoler MH, Jukic DM, Tommasino M, Gheit T, et al. Cutaneous human papillomavirus types detected on the surface of male external genital lesions: a case series within the HPV Infection in Men Study. *J Clin Virol.* 2013;58(4):652-9, <http://dx.doi.org/10.1016/j.jcv.2013.10.011>.
69. Sichero L, Nyitray AG, Nunes EM, Nepal B, Ferreira S, Sobrinho JS, et al. Diversity of human papillomavirus in the anal canal of men: the HIM Study. *Clin Microbiol Infect.* 2015;21(5):502-9, <http://dx.doi.org/10.1016/j.cmi.2014.12.023>.
70. Paolini F, Rizzo C, Sperduti I, Pichi B, Mafera B, Rahimi SS, et al. Both mucosal and cutaneous papillomaviruses are in the oral cavity but only alpha genus seems to be associated with cancer. *J Clin Virol.* 2013;56(1):72-6, <http://dx.doi.org/10.1016/j.jcv.2012.09.016>.
71. Rahman S, Rollison DE, Pierce Campbell CM, Waterboer T, Michel A, Pawlita M, et al. Seroprevalence of Cutaneous Human Papillomaviruses and the Risk of External Genital Lesions in Men: A Nested Case-Control Study. *PLoS One.* 2016;11(11):e0167174, <http://dx.doi.org/10.1371/journal.pone.0167174>.
72. Torres M, Gheit T, McKay-Chopin S, Rodríguez C, Romero JD, Filotico R, et al. Prevalence of beta and gamma human papillomaviruses in the



- anal canal of men who have sex with men is influenced by HIV status. *J Clin Virol.* 2015;67:47-51, <http://dx.doi.org/10.1016/j.jcv.2015.04.005>.
73. Fakhry C, D'souza G, Sugar E, Weber K, Goshu E, Minkoff H, et al. Relationship between prevalent oral and cervical human papillomavirus infections in human immunodeficiency virus-positive and negative women. *J Clin Microbiol.* 2006;44(12):4479-85, <http://dx.doi.org/10.1128/JCM.01321-06>.
 74. Hampras SS, Rollison DE, Giuliano AR, McKay-Chopin S, Minoni L, Sereday K, et al. Prevalence and Concordance of Cutaneous Beta Human Papillomavirus Infection at Mucosal and Cutaneous Sites. *J Infect Dis.* 2017;216(1):92-96, <http://dx.doi.org/10.1093/infdis/jix245>.
 75. Smith EM, Ritchie JM, Yankowitz J, Wang D, Turek LP, Haugen TH. HPV prevalence and concordance in the cervix and oral cavity of pregnant women. *Infect Dis Obstet Gynecol.* 2004;12(2):45-56, <http://dx.doi.org/10.1080/1064744040009896>.
 76. Steinau M, Gorbach P, Gratzler B, Braxton J, Kerndt PR, Crosby RA, et al. Concordance between Anal and Oral Human Papillomavirus (HPV) Infections Among Young Men Who Have Sex with Men. *J Infect Dis.* 2017; 215(12):1832-5, <http://dx.doi.org/10.1093/infdis/jix232>.
 77. Termine N, Giovannelli L, Matranga D, Caleca MP, Bellavia C, Perino A, et al. Oral human papillomavirus infection in women with cervical HPV infection: new data from an Italian cohort and a metaanalysis of the literature. *Oral Oncol.* 2011;47(4):244-50, <http://dx.doi.org/10.1016/j.oraloncology.2011.02.011>.
 78. Gissmann L, zur Hausen H. Physical characterization of deoxy ribonucleic acids of different human papilloma viruses (HPV). *Med Microbiol Immunol.* 1978;166(1-4):3-11, <http://dx.doi.org/10.1007/BF02121128>.
 79. zur Hausen H. Papillomavirus infections-a major cause of human cancers. *Biochim Biophys Acta.* 1996;1288(2):F55-78, [http://dx.doi.org/10.1016/0304-419X\(96\)00020-0](http://dx.doi.org/10.1016/0304-419X(96)00020-0).
 80. Tommasino M. The biology of beta human papillomaviruses. *Virus Res.* 2017;231:128-38, <http://dx.doi.org/10.1016/j.virusres.2016.11.013>.
 81. Stubenrauch F, Malejczyk J, Fuchs PG, Pfister H. Late promoter of human papillomavirus type 8 and its regulation. *J Virol.* 1992;66(6):3485-93.
 82. Sankovski E, Männik A, Geimanen J, Ustav E, Ustav M. Mapping of betapapillomavirus human papillomavirus 5 transcription and characterization of viral-genome replication function. *J Virol.* 2014;88(2):961-73, <http://dx.doi.org/10.1128/JVI.01841-13>.
 83. Tommasino M. The human papillomavirus family and its role in carcinogenesis. *Semin Cancer Biol.* 2014;26:13-21, <http://dx.doi.org/10.1016/j.semcancer.2013.11.002>.
 84. Oriel JD. Natural history of genital warts. *Br J Vener Dis.* 1971;47(1):1-13.
 85. Doorbar J. Host control of human papillomavirus infection and disease. *Best Pract Res Clin Obstet Gynaecol.* 2018;47:27-41, <http://dx.doi.org/10.1016/j.bpobgyn.2017.08.001>.
 86. Werness BA, Levine AJ, Howley PM. Association of human papillomavirus types 16 and 18 E6 proteins with p53. *Science.* 1990;248(4951):76-9, <http://dx.doi.org/10.1126/science.2157286>.
 87. Dyson N, Howley PM, Münger K, Harlow E. The human papilloma virus-16 E7 oncoprotein is able to bind to the retinoblastoma gene product. *Science.* 1989;243(4893):934-7, <http://dx.doi.org/10.1126/science.2537532>.
 88. Scheffner M, Werness BA, Huibregtse JM, Levine AJ, Howley PM. The E6 oncoprotein encoded by human papillomavirus types 16 and 18 promotes the degradation of p53. *Cell.* 1990;63(6):1129-36, [http://dx.doi.org/10.1016/0092-8674\(90\)90409-8](http://dx.doi.org/10.1016/0092-8674(90)90409-8).
 89. Yim EK, Park JS. The role of HPV E6 and E7 oncoproteins in HPV-associated cervical carcinogenesis. *Cancer Res Treat.* 2005;37(6):319-24, <http://dx.doi.org/10.4143/crt.2005.37.6.319>.
 90. Wallace NA, Robinson K, Galloway DA. Beta human papillomavirus E6 expression inhibits stabilization of p53 and increases tolerance of genomic instability. *J Virol.* 2014;88(11):6112-27, <http://dx.doi.org/10.1128/JVI.03808-13>.
 91. Wallace NA, Robinson K, Howie HL, Galloway DA. β -HPV 5 and 8 E6 disrupt homology dependent double strand break repair by attenuating BRCA1 and BRCA2 expression and foci formation. *PLoS Pathog.* 2015; 11(3):e1004687, <http://dx.doi.org/10.1371/journal.ppat.1004687>.
 92. Howley PM, Pfister HJ. Beta genus papillomaviruses and skin cancer. *Virology.* 2015;479-480:290-6, <http://dx.doi.org/10.1016/j.virol.2015.02.004>.
 93. Wallace NA, Robinson K, Howie HL, Galloway DA. HPV 5 and 8 E6 abrogate ATR activity resulting in increased persistence of UVB induced DNA damage. *PLoS Pathog.* 2012;8(7):e1002807, <http://dx.doi.org/10.1371/journal.ppat.1002807>.
 94. Wallace NA, Gasior SL, Faber ZJ, Howie HL, Deininger PL, Galloway DA. HPV 5 and 8 E6 expression reduces ATM protein levels and attenuates LINE-1 retrotransposition. *Virology.* 2013;443(1):69-79, <http://dx.doi.org/10.1016/j.virol.2013.04.022>.
 95. Caldeira S, Zehbe I, Accardi R, Malanchi I, Dong W, Giarrè M, et al. The E6 and E7 proteins of the cutaneous human papillomavirus type 38 display transforming properties. *J Virol.* 2003;77(3):2195-206, <http://dx.doi.org/10.1128/JVI.77.3.2195-2206.2003>.
 96. Accardi R, Dong W, Smet A, Cui R, Hautefeuille A, Gabet AS, et al. Skin human papillomavirus type 38 alters p53 functions by accumulation of deltaNp73. *EMBO Rep.* 2006;7(3):334-40, <http://dx.doi.org/10.1038/sj.embo.7400615>.
 97. Cornet I, Bouvard V, Campo MS, Thomas M, Banks L, Gissmann L, et al. Comparative analysis of transforming properties of E6 and E7 from different beta human papillomavirus types. *J Virol.* 2012;86(4):2366-70, <http://dx.doi.org/10.1128/JVI.06579-11>.
 98. Dong W, Klotz U, Accardi R, Caldeira S, Tong WM, Wang ZQ, et al. Skin hyperproliferation and susceptibility to chemical carcinogenesis in transgenic mice expressing E6 and E7 of human papillomavirus type 38. *J Virol.* 2005;79(23):14899-908, <http://dx.doi.org/10.1128/JVI.79.23.14899-14908.2005>.
 99. Viariso D, Mueller-Decker K, Klotz U, Aengeneyndt B, Kopp-Schneider A, Gröne HJ, et al. E6 and E7 from beta HPV38 cooperate with ultraviolet light in the development of actinic keratosis-like lesions and squamous cell carcinoma in mice. *PLoS Pathog.* 2011;7(7):e1002125, <http://dx.doi.org/10.1371/journal.ppat.1002125>.
 100. Viariso D, Decker KM, Aengeneyndt B, Flechtenmacher C, Gissmann L, Tommasino M. Human papillomavirus type 38 E6 and E7 act as tumour promoters during chemically induced skin carcinogenesis. *J Gen Virol.* 2013;94(Pt 4):749-52, <http://dx.doi.org/10.1099/vir.0.048991-0>.
 101. Viariso D, Müller-Decker K, Hassel JC, Alvarez JC, Flechtenmacher C, Pawlita M, et al. The BRAF Inhibitor Vemurafenib Enhances UV-Induced Skin Carcinogenesis in Beta HPV38 E6 and E7 Transgenic Mice. *J Invest Dermatol.* 2017;137(1):261-4, <http://dx.doi.org/10.1016/j.jid.2016.08.030>.
 102. Viariso D, Müller-Decker K, Zanna P, Klotz U, Aengeneyndt B, Accardi R, et al. Novel β -HPV49 Transgenic Mouse Model of Upper Digestive Tract Cancer. *Cancer Res.* 2016;76(14):4216-25, <http://dx.doi.org/10.1158/0008-5472.CAN-16-0370>.
 103. Simmonds M, Storey A. Identification of the regions of the HPV 5 E6 protein involved in Bak degradation and inhibition of apoptosis. *Int J Cancer.* 2008;123(10):2260-6, <http://dx.doi.org/10.1002/ijc.23815>.
 104. Bedard KM, Underbrink MP, Howie HL, Galloway DA. The E6 oncoproteins from human betapapillomaviruses differentially activate telomerase through an E6AP-dependent mechanism and prolong the lifespan of primary keratinocytes. *J Virol.* 2008;82(8):3894-902, <http://dx.doi.org/10.1128/JVI.01818-07>.
 105. Shterzer N, Heyman D, Shapiro B, Yaniv A, Jackman A, Serour F, et al. Human papillomavirus types detected in skin warts and cancer differ in their transforming properties but commonly counteract UVB induced protective responses in human keratinocytes. *Virology.* 2014;468-470:647-659, <http://dx.doi.org/10.1016/j.virol.2014.09.015>.
 106. Schmitt A, Harry JB, Rapp B, Wettstein FO, Iftner T. Comparison of the properties of the E6 and E7 genes of low- and high-risk cutaneous papillomaviruses reveals strongly transforming and high Rb-binding activity for the E7 protein of the low-risk human papillomavirus type 1. *J Virol.* 1994;68(11):7051-9.
 107. Akgül B, García-Escudero R, Ghali L, Pfister HJ, Fuchs PG, Navsaria H, et al. The E7 protein of cutaneous human papillomavirus type 8 causes invasion of human keratinocytes into the dermis in organotypic cultures of skin. *Cancer Res.* 2005;65(6):2216-23, <http://dx.doi.org/10.1158/0008-5472.CAN-04-1952>.
 108. Akgül B, Lemme W, García-Escudero R, Storey A, Pfister HJ. UV-irradiation stimulates the promoter activity of the high-risk, cutaneous human papillomavirus 5 and 8 in primary keratinocytes. *Arch Virol.* 2005;150(1):145-51, <http://dx.doi.org/10.1007/s00705-004-0398-4>.