

# Association between interleukin-22 genetic polymorphisms and bladder cancer risk

Tao Zhao,<sup>1</sup> XiaoHou Wu,<sup>1,\*</sup> JiaJi Liu<sup>1</sup>

<sup>1</sup>Chongqing Medical University, YongChuan Hospital, Department of Urology, YongChuan, Chongqing, China. <sup>2</sup>The First Affiliated Hospital of Chongqing, Medical University, Department of Urology, Chongqing, China.

**OBJECTIVE:** The cytokine interleukin-22 (IL-22), which is produced by T cells and natural killer cells, is associated with tumorigenesis and tumor progression in cancers. However, the role of IL-22 in bladder cancer has not been investigated.

**MATERIALS AND METHODS:** A prospective hospital-based case-control study comprising 210 patients with pathologically proven bladder cancer and 210 age- and gender-matched healthy controls was conducted. The genotypes of 3 common polymorphisms (-429 C/T, +1046 T/A and +1995 A/C) of the *IL-22* gene were determined with fluorogenic 5' exonuclease assays.

**RESULTS:** Patients with bladder cancer had a significantly higher frequency of the *IL-22* -429 TT genotype [odds ratio (OR)=2.04, 95% confidence interval (CI)=1.19, 3.49;  $p=0.009$ ] and -429 T allele (OR=1.42, 95% CI=1.08, 1.87;  $p=0.01$ ) than the healthy controls. These findings were still significant after a Bonferroni correction. When stratifying according to the stage of bladder cancer, we found that patients with superficial bladder cancer had a significantly lower frequency of the *IL-22* -429 TT genotype (OR=0.48, 95% CI=0.23, 0.98;  $p=0.04$ ). When stratifying according to the grade and histological type of bladder cancer, we found no statistical association. The *IL-22* +1046 T/A and *IL-22* +1995 A/C gene polymorphisms were not associated with the risk of bladder cancer.

**CONCLUSION:** To the authors' knowledge, this is the first report documenting that the *IL-22* -429 C/T gene polymorphism is associated with bladder cancer risk. Additional studies are required to confirm this finding.

**KEYWORDS:** Interleukin-22; Gene Polymorphism; Bladder Cancer; Case-Control Study.

Zhao T, Wu X, Liu J. Association between interleukin-22 genetic polymorphisms and bladder cancer risk. Clinics. 2015;70(10):686-690

Received for publication on May 29, 2015; First review completed on June 08, 2015; Accepted for publication on July 21, 2015

E-mail: wuxiaohou@outlook.com

\*Corresponding author

## ■ INTRODUCTION

Bladder cancer is the most common cancer of the urinary tract and is the fourth most commonly diagnosed malignancy in men in the United States. It is estimated that 74,000 new cases of bladder cancer are expected to occur in the United States in 2015 (1). Smoking tobacco and occupational exposure to chemical carcinogens have been established as the strongest risk factors for developing bladder cancer (2-4). It is now commonly accepted that the cause of bladder cancer is the multi-factorial interaction of environmental triggers with genetic susceptibility (5-9). Genome-wide association studies have identified multiple susceptibility loci associated with bladder cancer risk (10-13).

Interleukin-22 (IL-22) is a member of the *IL-10* family or *IL-10* superfamily, a class of potent mediators of cellular

inflammatory responses (14). IL-22 is synthesized by different cell types, including T- and natural killer (NK)-cells and has been reported to mediate crosstalk between inflammatory cells and keratinocytes (15-17). IL-22 is also associated with tumorigenesis and tumor progression in cancers (18,19). The human *IL-22* gene is located on the long arm of chromosome 12, on 12q15, approximately 52 and 99 kbp upstream from the *IL-26* and interferon loci, respectively and has the same transcriptional orientation as these two adjoining genes (15). Several single nucleotide polymorphisms (SNPs) have previously been identified in the *IL-22* gene (20-27).

However, the role of IL-22 in bladder cancer has not been investigated. The aim of this study was to investigate the association between *IL-22* gene polymorphisms (-429 C/T, +1046 T/A and +1995 A/C) and the risk of bladder cancer in a Chinese population.

## ■ MATERIALS AND METHODS

### Study population

A prospective hospital-based case-control study of 210 patients with pathologically proven bladder cancer and 210

**Copyright** © 2015 CLINICS – This is an Open Access article distributed under the terms of the Creative Commons License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium or format, provided the original work is properly cited.

No potential conflict of interest was reported.

**DOI:** 10.6061/clinics/2015(10)05

**Table 1** - Primer and probe sequences for the 5'-exonuclease assays of *IL-22* polymorphisms.

SNPs	rs2227485	rs1182844	rs1179246
Exchange	-429 C/T	+ 1046 T/A	+ 1995 A/C
Forward Primer	AAAATGAGTCCGTGACCAAAATGC	CCACCTATGAGACTTCCTATCAGT	GAAAAAGCCTTCCTGCCTAATGG
Reverse Primer	ACACAATTGTTTGTCTTAGTAGAGTTCAGAT	CACTAAAGGAAAAGGAAAGCTGTGTTT	GGTGTGCCTAAAGGTCAGA
Wild-type Probe	FAM-CTCCTATAGTGACTGAGTAA-NFQ	VIC-AAACTTACTAGTAGGTATGACTC-NFQ	VIC-TGAACAGAGTTATCTGCCTC-NFQ
Mutant Probe	VIC-CTCTATAGTGGCTGAGTAA-NFQ	FAM-CTTACTAGTAGGAATGACTC-NFQ	FAM-AACAGAGTTAGTGCCTC-NFQ

SNPs: single nucleotide polymorphisms.

**Table 2** - Distribution of the characteristics of bladder cancer cases and healthy control subjects.

	Cases (n=210)	Controls (n=210)	p
Sex (Male/Female)	134/76	127/83	0.48
Age (Years)	61.3 ± 9.8	60.7 ± 9.4	0.52
Smoking (Ever/Never)	111/99	85/125	0.01
Tumor stage (%)			
Invasive (T2-T4)	134(63.8)		
Superficial (Tis-T1)	76(36.2)		
Tumor grade (%)			
High (G2 + G3)	145(69.0)		
Low (G1)	65(31.0)		
Tumor histological type (%)			
Papillary	157(74.8)		
Nonpapillary	53(25.2)		

age- and gender-matched healthy controls was conducted in the Department of Urology of the YongChuan Hospital of ChongQing Medical University. The healthy control subjects were randomly selected when they attended a clinic for a routine examination. All the bladder cancer cases were staged according to the TNM staging system of the Union Internationale Contre le Cancer. Bladder tumors were graded using the World Health Organization (WHO) classification. All the individuals were interviewed by trained nurse-interviewers using a structured questionnaire that asked for information regarding the patients' gender, age, smoking status and occupational and other exposure histories. All parts of the study were approved by the Institutional Ethical Committee of the ChongQing Medical University and informed consent according to the Declaration of Helsinki

was obtained from all the participants or their families/surrogates.

### DNA extraction and genotyping

DNA was extracted from peripheral blood lymphocytes using a commercially available Qiagen kit (Qiagen Inc., Valencia, CA, USA). The genotypes of 3 common polymorphisms (-429 C/T, +1046 T/A and +1995 A/C) of the *IL-22* gene were determined with fluorogenic 5' exonuclease assays (TaqMan, Applied Biosystems, Foster City, CA, USA). The primer and probe sequences for the 5'-exonuclease assays of the *IL-22* polymorphisms are listed in Table 1. The polymerase chain reaction (PCR) was performed in a Primus 96 plus thermal cycler using a total volume of 5 µl containing 2.5 µl of Universal-MasterMix, 0.125 µl of 40x Assay-by-Design mix, 0.375 µl of H<sub>2</sub>O and 2 µl of DNA. The reactions were overlaid with 15 µl of mineral oil. The cycling parameters were as follows: 10 min at 94°C for primary denaturation, followed by 40 cycles of 20 s at 92°C and 1 min at 60°C. Fluorescence was measured in a Lambda Fluoro 320 Plus plate reader (MWG Biotech AG, Germany).

### Statistical analysis

The data were presented as the means ± standard deviation (SD) or as percentages for categorical variables. Differences between continuous variables were assessed using Student's *t* test, while those between categorical variables were evaluated using Pearson's  $\chi^2$  test. A multivariate logistic regression analysis was used to calculate crude and adjusted odds ratios (OR) and 95% confidence intervals (CI) for the association between each *IL-22* polymorphism and bladder cancer risk. Statistical significance

**Table 3** - Genotype and allele frequencies of *IL-22* gene polymorphisms among bladder cancer cases and healthy controls.

Genotypes	Cases (n=210)	Controls (n=210)	OR (95% CI)	p
-429 CC	68(32.4)	79(37.6)	1.00(Reference)	
-429 CT	84(40.0)	98(46.7)	1.00(0.64, 1.54)	0.99
-429 TT	58(27.6)	33(15.7)	2.04(1.19, 3.49)	0.009
-429 C allele frequency	220(52.4)	256(60.9)	1.00(Reference)	
-429 T allele frequency	200(47.6)	164(39.1)	1.42(1.08, 1.87)	0.01
+ 1046 TT	90(42.8)	97(46.2)	1.00(Reference)	
+ 1046 TA	77(36.7)	74(35.2)	1.12(0.73, 1.72)	0.60
+ 1046 AA	43(20.5)	39(18.6)	1.19(0.71, 2.00)	0.52
+ 1046 T allele frequency	257(61.2)	268(63.8)	1.00(Reference)	
+ 1046 A allele frequency	163(38.8)	152(36.2)	1.12(0.85, 1.48)	0.43
+ 1995 AA	68(32.4)	65(30.9)	1.00(Reference)	
+ 1995 AC	101(48.1)	98(46.7)	0.99(0.64, 1.53)	0.95
+ 1995 CC	41(19.5)	47(22.4)	0.83(0.49, 1.43)	0.51
+ 1995 A allele frequency	237(56.4)	228(54.3)	1.00(Reference)	
+ 1995 C allele frequency	183(43.6)	192(45.7)	0.92(0.70, 1.20)	0.53

OR: odds ratio; CI: confidence interval.



**Table 4** - Stratification analysis of the *IL-22* -429 C/T polymorphism in bladder cancer cases.

		CC		
	Cases (n=210)	n (%)	OR (95% CI)	p
Tumor stage	210	68(32.4)	1(Reference)	
Invasive	134	40(29.9)	0.92(0.59, 1.44)	0.72
Superficial	76	28(36.8)	1.14(0.68, 1.90)	0.62
Tumor grade	210	68(32.4)	1(Reference)	
High	145	48(33.1)	1.02(0.67, 1.57)	0.92
Low	65	20(30.8)	0.95(0.54, 1.68)	0.86
Tumor histological type	210	68(32.4)	1(Reference)	
Papillary	157	49(31.2)	0.96(0.63, 1.47)	0.86
Nonpapillary	53	19(35.8)	1.11(0.61, 2.00)	0.74
		CT		
	Cases (n=210)	n (%)	OR (95% CI)	P
Tumor stage	210	84(40.0)	1(Reference)	
Invasive	134	46(34.3)	0.86(0.56, 1.31)	0.48
Superficial	76	38(50.0)	1.25(0.79, 1.99)	0.35
Tumor grade	210	84(40.0)	1(Reference)	
High	145	55(37.9)	0.95(0.64, 1.42)	0.80
Low	65	29(44.6)	1.12(0.67, 1.85)	0.67
Tumor histological type	210	84(40.0)	1(Reference)	
Papillary	157	66(42.0)	1.05(0.72, 1.54)	0.80
Nonpapillary	53	18(34.0)	0.85(0.47, 1.53)	0.59
		TT		
	Cases (n=210)	n (%)	OR (95% CI)	p
Tumor stage	210	58(27.6)	1(Reference)	
Invasive	134	48(35.8)	1.30(0.84, 2.01)	0.25
Superficial	76	10(13.2)	0.48(0.23, 0.98)	0.04
Tumor grade	210	58(27.6)	1(Reference)	
High	145	42(29.0)	1.05(0.67, 1.65)	0.84
Low	65	16(24.6)	0.89(0.48, 1.66)	0.72
Tumor histological type	210	58(27.6)	1(Reference)	
Papillary	157	42(26.8)	0.97(0.62, 1.52)	0.89
Nonpapillary	53	16(30.2)	1.09(0.58, 2.05)	0.78

OR: odds ratio; CI: confidence interval.

was set at a nominal *p*-value <0.05 for all comparisons. SAS version 9.1 (SAS Institute, Cary, NC) was used for all the statistical tests.

## RESULTS

The characteristics of the bladder cancer cases and healthy control subjects are presented in Table 2. A univariate analysis was performed and the results indicated that smoking (*p*=0.01) was associated with bladder cancer (Table 2). No significant difference was found between the bladder cancer cases and healthy controls regarding sex or age (Table 2). Regarding the tumor stage of the cases, 134 (63.8%) cases were invasive (T2-T4) bladder cancer and 76 (36.2%) were superficial (Tis-T1) bladder cancer. Regarding the tumor grades of the cases, 65 (31.0%) cases were low grade (G1) bladder cancer and 145 (69.0%) cases were high grade (G2+G3) bladder cancer. The analysis of the histological types of the cases found that 53 (25.2%) cases were nonpapillary bladder cancer and 157 (74.8%) were papillary bladder cancer.

The genotype frequencies were in agreement with the Hardy-Weinberg equilibrium. The patients with bladder cancer had a significantly higher frequency of the *IL-22* -429 TT genotype (OR=2.04, 95% CI=1.19, 3.49; *p*=0.009) and -429 T allele (OR=1.42, 95% CI=1.08, 1.87; *p*=0.01) than the

healthy control subjects (Table 3). The findings remained significant after a Bonferroni correction was implemented. When stratifying according to the stage of bladder cancer, we found that superficial bladder cancer had a significantly lower frequency of the *IL-22* -429 TT genotype (OR=0.48, 95% CI=0.23, 0.98; *p*=0.04) (Table 4). When stratifying according to the grade and histological type of bladder cancer, we found no statistical association (Table 4). The *IL-22* +1046 T/A and *IL-22* +1995 A/C gene polymorphisms were not associated with the risk of bladder cancer (Table 3).

## DISCUSSION

In this study, we investigated the association between three common polymorphisms (-429 C/T, +1046 T/A and +1995 A/C) of the *IL-22* gene and the risk of bladder cancer in a Chinese population. This prospective hospital-based case-control study revealed that the *IL-22* -429 C/T gene polymorphism is associated with bladder cancer risk. To the best of our knowledge, this is the first report in the literature that evaluated the association between *IL-22* gene polymorphisms and the risk of bladder cancer.

There is accumulating evidence that genetics plays a key role in the susceptibility to and clinicopathologic characteristics of



bladder cancer. Nine meta-analyses, each of which analyzed between four and twenty-four studies, have provided evidence that the following polymorphisms are associated with increased bladder cancer risk: *NQO1* Pro187Ser; *PSCA* rs2294008 (C>T); *XRCC1* Arg399Gln (especially in non-Asian populations); *MMP-2* -1306 C/T; *MMP-9* -1562 C/T; *XPD* Lys751Gln; *ERCC2* Arg156Arg; Asp312Asn; Lys751Gln; *CCND1* G870A (which may modulate the risk of bladder cancer in conjunction with tobacco smoking); *XPC* A499V (in Caucasian populations); *CYP1A1* polymorphisms (especially the 11599G>C, 2455A>G, 3810T>C, and 113T>C polymorphisms in Asians); *MDM2* SNP309T>G (among Caucasians) (28-36).

The mechanisms of action for the *IL-22* -429 TT genotype and T allele as risk factors for bladder cancer are still unclear. *IL-22* may play a role in controlling tumor growth and tumor progression by inhibiting signaling pathways that promote tumor cell proliferation, such as ERK1/2 and AKT phosphorylation (37). The *IL-22* -429 C/T gene polymorphism has been associated with the risks for various cancers. Genetic polymorphisms and plasma levels of *IL-22* contribute to the development of non-small cell lung cancer (38). Recently, a case-control study found that the *IL-22* -429 C/T gene polymorphism might be associated with the risk and multifocality of papillary thyroid cancer (26). In 561 colon cancer cases and 722 population controls, an association study suggested that the rs1179251 SNP in *IL-22* was associated with the risk of colon cancer (23).

Several limitations to this study should be mentioned. First, these results should be interpreted with caution because the study subjects were Chinese; therefore, the study does not permit extrapolation of the results to different ethnic populations. Second, this is a hospital-based case-control study. Therefore, a selection bias could not be avoided and the subjects may not be representative of the general population. Third, the sample size of our study was relatively small and may not have had adequate statistical power in detecting significant differences. Finally, we did not evaluate the relationships of the 3 common polymorphisms (-429 C/T, +1046 T/A and +1995 A/C) to the plasma levels of *IL-22*, which may potentially reflect the disease state of patients. Therefore, the association of the *IL-22* polymorphisms with plasma levels of *IL-22* should be further investigated.

In conclusion, to the authors' knowledge, this is the first report documenting that the *IL-22* -429 C/T gene polymorphism is associated with bladder cancer risk. Additional studies are required to confirm this finding.

## ACKNOWLEDGMENTS

Thanks are given to all coinvestigators, local project coordinators, research assistants, laboratory technicians, and secretaries/administrative assistants.

## AUTHOR CONTRIBUTIONS

Zhao T and Wu X carried out the molecular genetic studies and drafted the manuscript. Liu J carried out the genotyping. Zhao T and Liu J participated in the design of the study and performed the statistical analysis. Zhao T, Wu X and Liu J conceived the study, participated in its design and coordination and helped to draft the manuscript. All the authors read and approved the final manuscript.

## REFERENCES

- Siegel R, Ma J, Zou Z, Jemal A. Cancer statistics, 2014. *CA Cancer J Clin*. 2014;64(1):9-29, <http://dx.doi.org/10.3322/caac.21208>.
- Dowdy D. Tobacco smoking and bladder cancer. *JAMA*. 2011;306(20):2216-7; author reply 7.
- Bachand A, Mundt KA, Mundt DJ, Carlton LE. Meta-analyses of occupational exposure as a painter and lung and bladder cancer morbidity and mortality 1950-2008. *Crit Rev Toxicol*. 2010;40(2):101-25, <http://dx.doi.org/10.3109/10408440903352826>.
- Mommensen S, Aagaard J. Tobacco as a risk factor in bladder cancer. *Carcinogenesis*. 1983;4(3):335-8, <http://dx.doi.org/10.1093/carcin/4.3.335>.
- Wronski S. Genetic polymorphism intermingled with environmental factors substantially contributes to the bladder cancer progression. *Cent European J Urol*. 2013;66(1):12-3, <http://dx.doi.org/10.5173/cej.2013.01.art4>.
- Volanis D, Kadiyska T, Galanis A, Delakas D, Logotheti S, Zoumpourlis V. Environmental factors and genetic susceptibility promote urinary bladder cancer. *Toxicol Lett*. 2010;193(2):131-7, <http://dx.doi.org/10.1016/j.toxlet.2009.12.018>.
- Horikawa Y, Gu J, Wu X. Genetic susceptibility to bladder cancer with an emphasis on gene-gene and gene-environmental interactions. *Curr Opin Urol*. 2008;18(5):493-8, <http://dx.doi.org/10.1097/MOU.0b013e32830b88ff>.
- Hung RJ, Boffetta P, Brennan P, Malaveille C, Hautefeuille A, Donato F, et al. GST, NAT, SULT1A1, CYP1B1 genetic polymorphisms, interactions with environmental exposures and bladder cancer risk in a high-risk population. *Int J Cancer*. 2004;110(4):598-604, <http://dx.doi.org/10.1002/ijc.20157>.
- Hung RJ, Boffetta P, Brennan P, Malaveille C, Gelatti U, Placidi D, et al. Genetic polymorphisms of MPO, COMT, MnSOD, NQO1, interactions with environmental exposures and bladder cancer risk. *Carcinogenesis*. 2004;25(6):973-8, <http://dx.doi.org/10.1093/carcin/bgh080>.
- Rothman N, Garcia-Closas M, Chatterjee N, Malats N, Wu X, Figueroa JD, et al. A multi-stage genome-wide association study of bladder cancer identifies multiple susceptibility loci. *Nat Genet*. 2010;42(11):978-84, <http://dx.doi.org/10.1038/ng.687>.
- Garcia-Closas M, Ye Y, Rothman N, Figueroa JD, Malats N, Dinney CP, et al. A genome-wide association study of bladder cancer identifies a new susceptibility locus within SLC14A1, a urea transporter gene on chromosome 18q12.3. *Hum Mol Genet*. 2011;20(21):4282-9, <http://dx.doi.org/10.1093/hmg/ddr342>.
- Gu J, Chen M, Shete S, Amos CI, Kamat A, Ye Y, et al. A genome-wide association study identifies a locus on chromosome 14q21 as a predictor of leukocyte telomere length and as a marker of susceptibility for bladder cancer. *Cancer Prev Res (Phila)*. 2011;4(4):514-21.
- Rafnar T, Vermeulen SH, Sulem P, Thorleifsson G, Aben KK, Witjes JA, et al. European genome-wide association study identifies SLC14A1 as a new urinary bladder cancer susceptibility locus. *Hum Mol Genet*. 2011;20(21):4268-81, <http://dx.doi.org/10.1093/hmg/ddr303>.
- Andoh A, Zhang Z, Inatomi O, Fujino S, Deguchi Y, Araki Y, et al. Interleukin-22, a member of the IL-10 subfamily, induces inflammatory responses in colonic subepithelial myofibroblasts. *Gastroenterology*. 2005;129(3):969-84, <http://dx.doi.org/10.1053/j.gastro.2005.06.071>.
- Wolk K, Sabat R. Interleukin-22: a novel T- and NK-cell derived cytokine that regulates the biology of tissue cells. *Cytokine Growth Factor Rev*. 2006;17(5):367-80, <http://dx.doi.org/10.1016/j.cytogfr.2006.09.001>.
- Wolk K, Kunz S, Witte E, Friedrich M, Asadullah K, Sabat R. IL-22 increases the innate immunity of tissues. *Immunity*. 2004;21(2):241-54, <http://dx.doi.org/10.1016/j.immuni.2004.07.007>.
- Boniface K, Bernard FX, Garcia M, Gurney AL, Lecron JC, Morel F. IL-22 inhibits epidermal differentiation and induces proinflammatory gene expression and migration of human keratinocytes. *J Immunol*. 2005;174(6):3695-702, <http://dx.doi.org/10.4049/jimmunol.174.6.3695>.
- Nardinocchi L, Sonogo G, Passarelli F, Avitabile S, Scarponi C, Falla CM, et al. Interleukin-17 and interleukin-22 promote tumor progression in human nonmelanoma skin cancer. *Eur J Immunol*. 2015;45(3):922-31, <http://dx.doi.org/10.1002/eji.201445052>.
- Kim K, Kim G, Kim JY, Yun HJ, Lim SC, Choi HS. Interleukin-22 promotes epithelial cell transformation and breast tumorigenesis via MAP3K8 activation. *Carcinogenesis*. 2014;35(6):1352-61, <http://dx.doi.org/10.1093/carcin/bgu044>.
- Hennig BJ, Frodsham AJ, Hellier S, Knapp S, Yee LJ, Wright M, et al. Influence of IL-10RA and IL-22 polymorphisms on outcome of hepatitis C virus infection. *Liver Int*. 2007;27(8):1134-43, <http://dx.doi.org/10.1111/j.1478-3231.2007.01518.x>.
- Endam LM, Bosse Y, Filali-Mouhim A, Cormier C, Boisvert P, Boulet LP, et al. Polymorphisms in the interleukin-22 receptor alpha-1 gene are associated with severe chronic rhinosinusitis. *Otolaryngol Head Neck Surg*. 2009;140(5):741-7, <http://dx.doi.org/10.1016/j.otohns.2008.12.058>.
- Weger W, Hofer A, Wolf P, El-Shabrawi Y, Renner W, Kerl H, et al. Common polymorphisms in the interleukin-22 gene are not associated with chronic plaque psoriasis. *Exp Dermatol*. 2009;18(9):796-8, <http://dx.doi.org/10.1111/j.1600-0625.2009.00840.x>.
- Thompson CL, Plummer SJ, Tucker TC, Casey G, Li L. Interleukin-22 genetic polymorphisms and risk of colon cancer. *Cancer Causes Control*. 2010;21(8):1165-70, <http://dx.doi.org/10.1007/s10552-010-9542-5>.
- Ding GG, Zhang GL, Chen XC, Zhang MX, Yang L, Wang Z. [A study on the relationship between single nucleotide polymorphisms of interleukin-22





- and susceptibility to pulmonary tuberculosis]. *Zhonghua Jie He He Hu Xi Za Zhi*. 2012;35(8):596-600.
25. Musolino C, Allegra A, Ferraro M, Aguenouz M, Russo S, Alonci A, et al. Involvement of T2677T multidrug resistance gene polymorphism in Interleukin 22 plasma concentration in B-chronic lymphocytic leukemia patients. *Acta Oncol*. 2012;51(3):406-8, <http://dx.doi.org/10.3109/0284186X.2011.631577>.
  26. Eun YG, Shin IH, Lee YC, Shin SY, Kim SK, Chung JH, et al. Interleukin 22 polymorphisms and papillary thyroid cancer. *J Endocrinol Invest*. 2013; 36(8):584-7.
  27. Suh JS, Cho SH, Chung JH, Moon A, Park YK, Cho BS. A polymorphism of interleukin-22 receptor alpha-1 is associated with the development of childhood IgA nephropathy. *J Interferon Cytokine Res*. 2013;33(10):571-7, <http://dx.doi.org/10.1089/jir.2012.0097>.
  28. Yang S, Jin T, Su HX, Zhu JH, Wang DW, Zhu SJ, et al. The Association between NQO1 Pro187Ser Polymorphism and Bladder Cancer Susceptibility: A Meta-Analysis of 15 Studies. *PLoS One*. 2015;10(1):e0116500, <http://dx.doi.org/10.1371/journal.pone.0116500>.
  29. Zhao Y, Gui ZL, Liao S, Gao F, Ge YZ, Jia RP. Prostate stem cell antigen rs2294008 (C>T) polymorphism and bladder cancer risk: a meta-analysis based on cases and controls. *Genet Mol Res*. 2014;13(3):5534-40, <http://dx.doi.org/10.4238/2014.July.25.7>.
  30. Yang D, Liu C, Shi J, Wang N, Du X, Yin Q, et al. Association of XRCC1 Arg399Gln polymorphism with bladder cancer susceptibility: a meta-analysis. *Gene*. 2014;534(1):17-23, <http://dx.doi.org/10.1016/j.gene.2013.10.038>.
  31. Yan Y, Liang H, Li T, Li M, Li R, Qin X, et al. The MMP-1, MMP-2, and MMP-9 gene polymorphisms and susceptibility to bladder cancer: a meta-analysis. *Tumour Biol*. 2014;35(4):3047-52, <http://dx.doi.org/10.1007/s13277-013-1395-6>.
  32. Xiong T, Yang J, Wang H, Wu F, Liu Y, Xu R, et al. The association between the Lys751Gln polymorphism in the XPD gene and the risk of bladder cancer. *Mol Biol Rep*. 2014;41(4):2629-34, <http://dx.doi.org/10.1007/s11033-014-3121-x>.
  33. Wu Y, Yang Y. Complex association between ERCC2 gene polymorphisms, gender, smoking and the susceptibility to bladder cancer: a meta-analysis. *Tumour Biol*. 2014;35(6):5245-57, <http://dx.doi.org/10.1007/s13277-014-1682-x>.
  34. Guan Z, Zeng J, Wang Z, Xie H, Lv C, Ma Z, et al. Urine tenascinC is an independent risk factor for bladder cancer patients. *Mol Med Rep*. 2014; 9(3):961-6.
  35. Wang Y, Kong CZ, Zhang Z, Yang CM, Li J. Relationships between CYP1A1 genetic polymorphisms and bladder cancer risk: a meta-analysis. *DNA Cell Biol*. 2014;33(3):171-81, <http://dx.doi.org/10.1089/dna.2013.2298>.
  36. Wang HG, Wu QY, Zhou H, Peng XS, Shi MJ, Li JM, et al. The MDM2 SNP309T>G polymorphism increases bladder cancer risk among Caucasians: a meta-analysis. *Asian Pac J Cancer Prev*. 2014;15(13):5277-81, <http://dx.doi.org/10.7314/APJCP.2014.15.13.5277>.
  37. Weber GF, Gaertner FC, Erl W, Janssen KP, Bleichert B, Holzmann B, et al. IL-22-mediated tumor growth reduction correlates with inhibition of ERK1/2 and AKT phosphorylation and induction of cell cycle arrest in the G2-M phase. *J Immunol*. 2006;177(11):8266-72, <http://dx.doi.org/10.4049/jimmunol.177.11.8266>.
  38. Liu F, Pan X, Zhou L, Zhou J, Chen B, Shi J, et al. Genetic polymorphisms and plasma levels of interleukin-22 contribute to the development of non-small cell lung cancer. *DNA Cell Biol*. 2014;33(10):705-14, <http://dx.doi.org/10.1089/dna.2014.2432>.