

ORIGINAL RESEARCH

DOES ENDOTHELIAL DYSFUNCTION CORRELATE BETTER WITH WAIST-TO-HIP RATIO THAN WITH BODY MASS INDEX OR WAIST CIRCUMFERENCE AMONG OBESE PATIENTS?

Nivaldo Ribeiro Villela, Luiz Guilherme Kraemer Aguiar, Luciana Bahia, Daniel Bottino, and Eliete Bouskela

Villela NR, Aguiar LGK, Bahia L, Bottino D, Bouskela E. Does endothelial dysfunction correlate better with waist-to-hip ratio than with body mass index or waist circumference among obese patients? Clinics. 2006;61(1):53-8.

PURPOSE: Obesity is associated with cardiovascular disease, affecting large arteries and the microcirculation. Waist circumference and body mass index are routinely employed as measures for assessing obesity-related health risk, whereas waist-to-hip ratio is not. We aimed to investigate the association between brachial vascular reactivity and body mass index, waist circumference, and waist-to-hip ratio.

METHODS: Eighty-five volunteers (21 men/66 women), aged between 20 and 55 years, underwent determination of waist circumference, body mass index, waist-to-hip ratio, and endothelial function by venous occlusion plethysmography. Forearm blood flow was measured in response to intrabrachial artery infusions of 3 different concentrations of endothelium-dependent (acetylcholine 7.5, 15, and 30 mg/min) and endothelium-independent (sodium nitroprusside 2, 4, and 8 mg/min) vasodilators.

RESULTS: There was an inverse correlation of body mass index and waist circumference with forearm blood flow increments after acetylcholine and sodium nitroprusside infusions, while waist-to-hip ratio showed an inverse correlation with forearm blood flow increments only after acetylcholine. When subjects older than 40 years ($n = 25$) were excluded from the analysis, the inverse correlation of body mass index with forearm blood flow increments after acetylcholine infusion no longer existed, while waist circumference and waist-to-hip ratio showed the same results observed before.

CONCLUSION: The waist-to-hip ratio is probably a better estimator of endothelial dysfunction and possibly of cardiovascular risk than body mass index. These findings underscore the importance of routinely collecting hip circumference as an obesity index and risk estimator.

KEYWORDS: Endothelium. Vascular Reactivity. Waist-to-Hip Ratio. Body Mass Index. Waist Circumference.

INTRODUCTION

Obesity is a chronic disease, the incidence of which is growing alarmingly, affecting not only adults but also children and adolescents. Obesity is a risk factor for a host of pathologies from arthritis to cardiovascular diseases. In cardiovascular disease, obesity increases the risk not only of

large artery disease, such as myocardial infarction and stroke,^{1,2} but also of disease entities that are in part caused by microangiopathy, notably retinopathy, nephropathy, and heart failure.³⁻⁵ These associations cannot be entirely explained by obesity-associated insulin resistance, hypertension, or dyslipidemia.^{3,5,6} Responsible mechanisms are likely to include metabolic and inflammatory responses to the increased amount of stored fat⁷ that may deeply and negatively influence endothelial physiology, a condition which may lead to the formation of the atherosclerotic plaque.⁸

Waist circumference (WC) is widely viewed as a sim-

Laboratório de Pesquisas em Microcirculação, Universidade do Estado do Rio de Janeiro, RJ, Brazil.

Email: eliete_bouskela@yahoo.com.br

Received for publication on December 30, 2005.

Accepted for publication on January 12, 2006.

ple but effective measure for assessing obesity-related health risks, whereas the measurement of hip circumference is not currently a priority. In a population-based study, Lisner et al⁹ showed that hip circumference was a significant independent inverse risk estimator for diabetes and cardiovascular endpoints. Recently Yusuf et al¹⁰ demonstrated, in the Interheart study, that waist-to-hip ratio (WHR) was a strong predictor of myocardial infarction, again suggesting that hip measurement should not be excluded from health surveys.

Endothelial dysfunction precedes and predicts clinical macrovascular disease.¹¹ Abnormal vasomotor responses occur in obesity and insulin resistance states.¹² Severe endothelial dysfunction even in the absence of obstructive coronary disease is associated with increased cardiac events,¹³ and the impairment of brachial artery endothelium-dependent vasodilation correlates with the same alteration in coronary arteries¹⁴.

The purpose of the present study was to define the relationship between brachial vascular reactivity and body mass index (BMI), WC, and WHR in Brazilian subjects.

METHODS

Subjects

Eighty-five volunteers (21 men – 66 women), aged between 20 and 55 years, from the Cardiometabolic Clinic for outpatient care at the State University of Rio de Janeiro participated in this study after giving written informed consent. The exclusion criteria were current tobacco use and any established cardiovascular and respiratory disease.

Study Protocol

The study was approved by the Ethical Committee of the State University of Rio de Janeiro.

The same trained examiner collected the anthropometric measurements in duplicate: waist at its smallest point with the abdomen relaxed; hip circumference at the level of the widest diameters around the buttocks; and body mass (weight) using a digital scale (Filizola®, São Paulo, SP, Brazil). Body mass index was defined as weight (in kg) divided by squared height (in meters).

Vascular reactivity was evaluated by measuring forearm blood flow (FBF) responses to intra-arterial infusions of acetylcholine (Ach; Acetilcolina®, USP, São Paulo, SP, Brazil) and sodium nitroprusside (SNP; NPS®, Lebon, Porto Alegre, RS, Brazil) to evaluate endothelium-dependent and -independent vasodilation, respectively. The study was performed in the morning after an 8 to 10-hour overnight fast

and in a temperature controlled room (20° to 22°C). A 27-gauge steel spinal needle (Becton, Dickinson Indústrias Cirúrgicas Ltda, Juiz de Fora, MG, Brazil) was inserted into the left brachial artery, anterior to the elbow, under sterile conditions and after local anesthesia with 1% lidocaine. A mercury-filled silastic strain-gauge was placed on the upper third of the forearm, which rested slightly above the level of the heart. Venous occlusion was accomplished using a blood pressure cuff applied proximal to the elbow and inflated to 40 mm Hg by a rapid cuff inflator. Wrist cuff was inflated to at least 50 mm Hg above systolic blood pressure, 1 minute before each flow measurement, to avoid hand shunt. Blood pressure was noninvasively measured simultaneously in the right arm at each period, and a 1-lead electrocardiogram was recorded continuously. Baseline blood flow was recorded after a 20 min infusion of NaCl 0.9%, at a rate of 0.8 mL/min, delivered by a Harvard pump (Harvard Apparatus Inc, South Natick, Massachusetts, USA). Acetylcholine and SNP were always infused in progressive dosages: 7.5, 15, and 30 mg/min and 2, 4, and 8 mg/min, respectively, for 5 minutes at each dose. Acetylcholine was always infused first, with an interval of 20 minutes between each drug. Blood flow was measured during the last 2 minutes for each dose and recorded for 10 seconds in every 15 seconds. The plethysmograph (Hokanson EC6, D.E. Honkanson Inc, Bellevue, WA, USA) was connected to an analog-to-digital converter (8SP, AD Instruments Pty Ltd, Castle Hill, Australia), and data were analyzed by Power Lab software on an IBM PC. Evaluations of FBF were made by calculation of the mean of at least 4 consecutive recordings.

Analysis

Forearm blood flow was expressed as mL/min per 100 mL forearm tissue. Responses to Ach and SNP were quantified as the increase in FBF above baseline value.

Data were analyzed using SPSS 8.0.0. Correlation analysis was performed by Spearman rank order test, and significant differences were assumed to be present at $P < .05$.

RESULTS

Table 1 displays the main characteristics of the subjects enrolled in the study.

Basal FBF before Ach correlated with basal FBF before SNP ($r = 0.61$, $P < .001$), which is an index of the good reliability of the method. It should be noted that 5 subjects received only Ach infusions, so FBF after SNP could not be evaluated in these subjects.

Table 1- General characteristics of the sample of the 85 subjects investigated

Variables	Quartiles	n
Age (years)	18 – 27	22
	27 – 37	26
	37 – 41	16
	41 – 55	21
BMI (kg/m ²)	19.8 – 30.8	20
	30.8 – 35.4	23
	35.4 – 40.4	21
	40.4 – 54.1	21
Waist (cm)	60 – 94	24
	94 – 101	19
	101 – 109	23
	109 – 129	19
WHR	0.68 – 0.83	23
	0.83 – 0.88	26
	0.88 – 0.92	17
	0.92 – 1.05	19

We observed significant inverse correlations of BMI and WC with FBF increments after Ach and SNP infusions ($P < .05$) (Table 2). The inverse correlation of WHR with FBF increment was significant after Ach ($P < .05$), but not after SNP infusion (Table 2).

When subjects older than 40 years were excluded ($n = 25$) from the analysis, a significant inverse correlation of WC and WHR, but not BMI, with the FBF increment after Ach was observed, with no significant correlation of WHR with SNP infusion (Table 3).

DISCUSSION

Obesity is associated with increased cardiovascular morbidity and mortality and is now regarded as a major independent risk factor.¹⁵ Obesity is still routinely diagnosed when the BMI equals or exceeds 30 kg/m². Our findings suggest that there is a stronger positive correlation of WHR with impaired nitric oxide-dependent vasodilation than of BMI or WC.

Data from long-term follow-up studies in adults show that impaired endothelium-dependent arterial vasorelaxation in response to muscarinic stimulation with acetylcholine precedes cardiovascular events by years.¹¹ Fichtlscherer et al¹⁶ found a good correlation between impaired endothelium-dependent FBF and recurrence of instability and cardiovascular event rates in patients with acute coronary syndrome. Numerous studies have evaluated lifestyle and pharmacological interventions to improve endothelial function and to limit cardiovascular risk.¹⁷ Woo et al¹⁸ reported an improvement of endothelial function in young obese after dietary modification alone or in combination with exercise training, showing that a primary intervention can change the cardiovascular risk.

Atherosclerosis is an inflammatory process, with metabolic disturbances such as obesity, hyperglycemia, and dyslipidemia considered to be proinflammatory triggers that initiate a chronic subclinical inflammatory status.¹⁹ Excessive fat accumulation plays a major role in initiating and maintaining this inflammatory process. The adipocyte is a

Table 2 - Correlations (Spearman r) between forearm blood flow and obesity indexes in all patients

	Body mass index		Waist circumference		Waist-to-hip ratio	
	r	P	r	P	r	P
Ach (7.5 µg/min) (n = 85)	-0.26	0.010	-0.32	0.003	-0.16	0.080
Ach (15 µg/min) (n = 85)	-0.22	0.020	-0.28	0.010	-0.24	0.020
Ach (30 µg/min) (n = 85)	-0.26	0.010	-0.27	0.012	-0.22	0.020
SNP (2 µg/min) (n = 80)	-0.31	0.001	-0.27	0.014	-0.16	0.100
SNP (4 µg/min) (n = 80)	-0.37	0.001	-0.32	0.005	-0.14	0.120
SNP (8 µg/min) (n = 80)	-0.37	0.001	-0.31	0.006	-0.16	0.090

Ach = acetylcholine; SNP = sodium nitroprusside

Table 3 - Correlations (Spearman r) between forearm blood flow and obesity indexes in patients aged ≤ 40 years

	Body mass index		Waist circumference		Waist-to-hip ratio	
	r	P	r	P	r	P
Ach (7.5 µg/min) (n = 60)	-0.19	0.180	-0.29	0.040	-0.35	0.010
Ach (15 µg/min) (n = 60)	-0.23	0.110	-0.27	0.050	-0.28	0.050
Ach (30 µg/min) (n = 60)	-0.28	0.060	-0.32	0.020	-0.28	0.050
SNP (2 µg/min) (n = 55)	-0.27	0.070	-0.30	0.030	-0.25	0.090
SNP (4 µg/min) (n = 55)	-0.35	0.010	-0.34	0.010	-0.17	0.260
SNP (8 µg/min) (n = 55)	-0.39	0.008	-0.36	0.010	-0.19	0.200

Ach = acetylcholine; SNP = sodium nitroprusside

complex and active endocrine cell, releasing numerous products, including free fatty acids, leptin, adiponectin, PAI-1, interleukin-6, tumor necrosis factor alpha, resistin, and angiotensin.²⁰

Although there is no doubt that intra-abdominal fat has the greatest impact as a cardiovascular risk factor, the metabolic role of peripheral fat has not been completely clarified. Ferreira et al²¹ showed that only visceral fat is adversely associated with large artery stiffness, whereas some degree of protection is conferred by peripheral fat and lean mass. There is a good positive correlation of intra-abdominal or visceral adipose tissue with insulin resistance, and it should be noted that patients of normal weight can also be insulin resistant.²² These findings call into question the usefulness of BMI as a risk marker, especially in lean persons.

Although accurate quantification of body fat compartments requires expensive techniques such as magnetic resonance imaging or computed tomography, clinical anthropometric measurements such as WC or WHR can be used to assess regional fat distributions. Yusuf et al,¹⁰ in a case-control study of acute myocardial infarction enrolling 27098 participants from 52 countries, showed a strong correlation between WHR and myocardial infarction risk, while BMI showed only a modest relation with it.

Impairment in endothelium-dependent vasodilation was also found by Higashi et al²³ in patients with a BMI higher than 30 kg/m². We found that BMI and WC had a negative correlation with FBF under the condition of endothelial-dependent vasodilation, but also under the condition of en-

dothelial-independent vasodilation. When only young patients were included in the analysis we found a significant inverse correlation only of WHR with FFB under the condition of endothelial-dependent vasodilation; this finding could be reflecting the probability that older subjects had higher rates of established arterial disease.

Our data show a stronger correlation of blunted responses to endothelium-dependent arterial vasorelaxation with WHR than with BMI, suggesting that WHR could be a precocious and more specific obesity index for endothelial dysfunction in this group. Findings in conduit arteries were also observed by Brook et al²⁴ who studied obese subjects and found a significantly blunted response in brachial artery flow-mediated dilation in subjects with a WHR ³ 0.85.

These findings underscore the importance of the collection of hip circumference as an obesity index and risk estimator. Considering that there is a good correlation of WHR with endothelial dysfunction of coronary and brachial arteries, it is possible that redefining obesity based on WHR, as proposed by Yusuf et al,¹⁰ would greatly increase the accuracy of estimation of myocardial infarction risk from obesity.

ACKNOWLEDGEMENTS

This work was supported by CNPq (505860/2004-0) and FAPERJ (E-26/152.020/2004).

The authors wish to thank Miss Kelly Silva de Andrade for technical assistance.

RESUMO

Villela NR, Aguiar LGK, Bahia L, Bottino D, Bouskela E. Em obesos, a disfunção endotelial correlaciona melhor com a relação cintura-quadril do que com a medida da cintura ou índice de massa corpórea. Clinics. 2006;61(1):53-8.

OBJETIVO: A obesidade é associada a doenças cardiovasculares e compromete tanto a macro como a micro-circulação. As medidas da cintura e do índice de massa corpórea são rotineiramente empregadas para avaliação do risco cardiovascular em obesos, enquanto a relação cintura-quadril é pouco utilizada. O objetivo do trabalho foi determinar que medida antropométrica, entre as rotineiramente usadas, avalia melhor o risco cardiovascular em obesos.

MATERIAL E MÉTODO: Oitenta e quatro voluntários (21 homens/ 66 mulheres), idade entre 20 e 55 anos foram

avaliados quanto ao diâmetro da cintura, ao índice de massa corpórea, à relação cintura-quadril e à função endotelial pela técnica de pletismografia com oclusão venosa para medida do fluxo sanguíneo braquial, em resposta a injeção intrabraquial de três doses de acetilcolina (7,5; 15 e 30 mg/min) ou de nitroprussiato de sódio (2; 4 e 8 mg/min), para avaliação da vasodilatação endotélio-dependente e -independente.

RESULTADO: Houve correlação inversa entre o índice de massa corpórea, diâmetro da cintura e aumento do fluxo sanguíneo após injeção de acetilcolina e nitroprussiato de sódio, enquanto que a relação cintura-quadril mostrou uma correlação negativa apenas com o aumento no fluxo de sangue no antebraço, após as infusões de acetilcolina. Quando os indivíduos com mais de 40 anos foram retirados da análise, não observamos mais a relação inversa entre índi-

ce de massa corpórea e aumento do fluxo sanguíneo após injeção de acetilcolina, enquanto que a cintura e a relação cintura-quadril mantiveram os resultados observados anteriormente.

CONCLUSÃO: A relação cintura-quadril é provavelmente um melhor índice para estimar a disfunção endotelial, e consequentemente o risco cardiovascular, que o índice de

massa corpórea e esses achados reforçam a importância da aferição da circunferência do quadril como um índice de obesidade e para estimativa do risco cardiovascular.

UNITERMOS: Disfunção endotelial. Vasodilatação. Diâmetro da cintura. Índice de massa corpórea. Relação cintura-quadril.

REFERENCES

1. Manson JE, Willett WC, Stampfer MJ, Colditz GA, Hunter DJ, Hankinson SE, et al. Body weight and mortality among women. *N Engl J Med*. 1995;333:677-85.
2. Bray GA. Medical consequences of obesity. *J Clin Endocrinol Metab*. 2004;89:2583-89.
3. van Leiden HA, Dekker JM, Moll AC, Nijpels G, Heine RJ, Bouter LM, et al. Blood pressure, lipids, and obesity are associated with retinopathy: the hoorn study. *Diabetes Care*. 2002;25:1320-5.
4. Serne EH, Ijzerman RG, de Jongh RT, Stehouwer CD. Blood pressure and insulin resistance: role for microvascular function? *Cardiovasc Res*. 2002;55:418-9.
5. Kenchaiah S, Evans JC, Levy D, Wilson PW, Benjamin EJ, Larson MG, et al. Obesity and the risk of heart failure. *N Engl J Med*. 2002;347:305-13.
6. de Jongh RT, Serne EH, Ijzerman RG, de Vries G, Stehouwer CD. Impaired microvascular function in obesity: implications for obesity-associated microangiopathy, hypertension, and insulin resistance. *Circulation*. 2004;109:2529-35.
7. Shankar SS, Steinberg HO. Obesity and endothelial dysfunction. *Semin Vasc Med*. 2005;5:56-64.
8. Avogaro A, de Kreutzenberg SV. Mechanisms of endothelial dysfunction in obesity. *Clin Chim Acta*. 2005;360:9-26.
9. Lissner L, Björkelund C, Heitmann BL, Seidell JC, Bengtsson C. Larger hip circumference independently predicts health and longevity in Swedish female cohort. *Obes Research*. 2001;9:644-46.
10. Yusuf S, Hawken S, Ôunupu S, Bautista L, Franzoli MG, Commerford P, et al. Obesity and the risk of myocardial infarction in 27,000 participants from 52 countries: a case-control study. *Lancet*. 2005;366:1640-49.
11. Suwaidi JA, Hamasaki S, Higano ST, Nishimura RA, Holmes DR, Lerman A. Long-term follow-up of patients with mild coronary artery disease and endothelial dysfunction. *Circulation*. 2000;101:948-54.
12. Steinberg HO, Chaker H, Leaming R, Johnson A, Brechtel G, Baron AD. Obesity/insulin resistance is associated with endothelial dysfunction. Implications for the syndrome of insulin resistance. *J Clin Invest*. 1996;97:2601-10.
13. Schächinger V, Britten MB, Zeiher AM. Prognostic impact of coronary vasodilator dysfunction on adverse long-term outcome of coronary heart disease. *Circulation*. 2000;101:1899-1906.
14. Neunteufl T, Katzenschlager R, Hassen A, Klaar U, Schwarzacher S, Glogar D, Bauer P, Weidinger F. Systemic endothelial dysfunction is related to the extent and severity of coronary artery disease. *Atherosclerosis*. 1997;129:111-18.
15. Eckel RH, Kraus RM. American Heart Association call to action: obesity as a major risk factor for coronary heart disease. *Circulation*. 1998;97:2099-100.
16. Fichtlscherer S, Breuer S, Zeiher AM. Prognostic value of systemic endothelial dysfunction in patients with acute coronary syndromes further evidence for the existence of the "vulnerable" patient. *Circulation*. 2004;110:1926-32.
17. Widlansky ME, Gokce N, Keaney JF, Vita JA. The clinical implications of endothelial dysfunction. *J Am Coll Cardiol*. 2003; 42:1149-60.
18. Woo KS, Chook P, Yu CW, Sung RY, Qiao M, Leung SS, et al. Effects of diet and exercise on obesity-related vascular dysfunction in children. *Circulation*. 2004;109:1981-6.
19. Soodini GR, Hamdy O. Obesity and endothelial function. *Curr Opin Endocrinol Diabetes*. 2004;11:186-91.

20. Yudkin JS, Stehouwer CDA, Emeis JJ, Coppack SW. C-reactive protein in healthy subjects: associations with obesity, insulin resistance, and endothelial dysfunction: a potential role for cytokines originating from adipose tissue? *Arterioscl Thromb Vasc Biol.* 1999;19:972-8.
21. Ferreira I, Snijder MB, Twisk JW, van Mechelen W. Central fat mass versus peripheral fat and lean mass: opposite (adverse versus favorable) associations with arterial stiffness? The Amsterdam Growth and Health Longitudinal Study. *J Clin Endocrinol Metab.* 2004;89: 2632-9.
22. Eckel RH, Grundy SM, Zimmet PZ. The metabolic syndrome. *Lancet.* 2005;365:1415-28.
23. Higashi Y, Sasaki S, Nakagawa K, Kimura M, Noma K, Sasaki S, et al. Low body mass index is a risk factor for impaired endothelium-dependent vasodilation in humans: role of nitric oxide and oxidative stress. *J Am Coll Cardiol.* 2003;42:256-63.
24. Brook RD, Bard RL, Rubenfire M, Ridker PM, Rajagopalan S. Usefulness of visceral obesity (waist/hip ratio) in predicting vascular endothelial function in healthy overweight adults. *Am J Cardiol.* 2001;88:1264-9.