



Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



Scientific letters

Usefulness of the antigen test for diagnosing SARS-CoV-2 infection in patients with and without symptoms[☆]



Utilidad del antígeno para el diagnóstico de infección por SARS-CoV-2 en pacientes con y sin síntomas

Dear Editor:

The COVID-19 pandemic has been accompanied by uncertainty. The need for tests to diagnose it in real time has led to tests being developed in record time, without the validation that would have been required in other circumstances.

Techniques based on protein detection offer theoretical advantages over those based on RNA amplification (which are more costly, slow and require sophisticated equipment)¹ and were included in the COVID-19 diagnostic protocols.²

SARS-CoV-2 antigen (Ag) determination (2019-nCoV Ag Test Fluorescence IC Assay[®] Shenzhen Bioeasy Biotechnology CO LTD, China, read with Bioeasy Immunofluorescence Analyzer mod EASY-11 Ref YRLE1105), which detects nucleocapsid protein, became available at the beginning of April. According to the manufacturer, the sample needed to be transported in a specific medium and preferably processed within 30 min of being taken, although the summary of product characteristics allows it to be carried out within 24 h and describes a sensitivity (SE) and specificity (SP) of 92% and 100%, respectively.

Our centre's COVID-19 working group decided to include this determination in patients with respiratory symptoms of less than 7 days' evolution, following expert recommendations.³ Nasal samples were collected and processed within a maximum of 30 min. If the result was negative, a polymerase chain reaction (PCR) test was performed. After one week of using the Ag test, doubts arose concerning its accuracy (positive results in asymptomatic patients for whom Ag tests had been ordered in spite of their not being included in the protocol's indications). It was therefore decided *a posteriori* to confirm all results with PCR.

Between 6 and 17 April, Ag detection was performed in 193 patients (44 asymptomatic, not anticipated in the protocol). In 172, samples were available for PCR (Table 1).

Taking the PCR as a benchmark, the SE, SP, positive predictive value (PPV) and negative predictive value (NPV) of the Ag test in symptomatic patients were 76%, 88.33%, 61.3% and 93.8%, respectively. The concordance in asymptomatic patients was much lower.

Cases with discordant results were analysed using clinical/radiological assessment, earlier/later PCR results and serological data where available. Of the 21 patients with positive Ag/negative PCR, 13 were considered Ag false positives (FPs) (9

Table 1
Ag detection.

		Symptomatic		Asymptomatic	
		PCR		PCR	
		Positive	Negative	Positive	Negative
Ag	Positive	19	12	0	9
	Negative	6	91	1	34
	Total	25	103	1	43

asymptomatic and 4 symptomatic with subsequent negative serology). In the remaining 8 patients, the clinical and epidemiological picture was compatible with a COVID-19 diagnosis, no other microbiological tests were carried out and they were considered PCR false negatives (FNs).

Of the 7 patients with negative Ag/positive PCR, one case was asymptomatic and in the rest the duration of the symptoms ranged from one to 18 days (median 4 days). In 4 patients, subsequent positive serology was available. All were considered to be Ag FNs.

If we consider clinical and radiological criteria, serology and earlier/later PCRs to define COVID cases, the SE, SP, PPV and NPV of Ag in symptomatic patients were 79.4%, 95.7%, 87.1% and 92.8%.

The discrepancies in the results may be because the Ag/PCR determinations were performed on different samples (although obtained on the same day) and because neither of the techniques has 100% SE and SP, with PCR, according to Börger's meta-analysis, having an SE of 73.3% (95% CI: 68.1–78.0).⁴

At our centre, the Ag determination was withdrawn due to the discordance of the results with PCR, the difficulties entailed by the limited time for processing and the need for a different transport medium than PCR, which required two samples to be taken. These disadvantages should be reconsidered in light of recent studies: Porte et al.⁵ kept samples refrigerated for 48 h with excellent results, thereby overcoming the time limitation, and other studies used a universal viral transport medium^{5,6} instead of that proposed by the manufacturer, thereby saving material, time and discomfort for the patient, there being no need to obtain a second sample for PCR.

The WHO currently recommends PCR as the technique of choice for COVID-19 diagnosis,⁷ but knowledge of the performance, advantages and limitations of tests taken at different times/on different samples/in different pandemic settings may help in developing suitable strategies for future situations. Based on our results and those of recent publications, Ag determination may be useful in symptomatic patients.

Funding

This study was partially funded through a call for proposals by the Department of Health of the Autonomous Government of Catalonia, code 6-17, principal investigator: Francesc Vidal.

References

[☆] Please cite this article as: Parada-Ricart E, Gomez-Bertomeu F, Picó-Plana E, Olona-Cabases M. Utilidad del antígeno para el diagnóstico de infección por SARS-CoV-2 en pacientes con y sin síntomas. *Enferm Infecc Microbiol Clin*. 2021;39:357–358.

1. Diao B, Wen K, Chen J, Liu Y, Yuan Z, Han C, et al. Diagnosis of acute respiratory syndrome coronavirus 2 infection by detection of nucleocapsid protein. medRxiv. 2020. <http://dx.doi.org/10.1101/2020.03.07.20032524>.
2. Mateo M. Procediment d'actuació enfront de casos d'infecció pel nou coronavirus SARS-CoV-2. Generalitat de Catalunya; p. 39 (Sub-direcció General de Vigilància i Resposta a Emergències de Salut Pública).
3. Grupo de expertos SEIMC para el análisis del diagnóstico microbiológico del COVID-19. Recomendaciones institucionales. Documento de posicionamiento de la SEIMC sobre el diagnóstico microbiológico de COVID-19. Soc Esp Enfermedades Infecc Microbiol Clín. 2020 https://seimc.org/documentos/documentoscnicos/recomendaciones/seimc-2020-Posicionamiento_SEIMC_diagnostico_microbiologico_COVID19.pdf
4. Böger B, Fachi MM, Vilhena RO, Cobre AF, Tonin FS, Pontarolo R. Systematic review with meta-analysis of the accuracy of diagnostic tests for COVID-19. Am J Infect Control. 2020 <https://linkinghub.elsevier.com/retrieve/pii/S0196655320306933>
5. Porte L, Legarraga P, Vollrath V, Aguilera X, Munita JM, Araos R, et al. Evaluation of novel antigen-based rapid detection test for the diagnosis of SARS-CoV-2 in respiratory samples. Int J Infect Dis IJID Off Publ Int Soc Infect Dis. 2020;99:328–33.
6. Weitzel T, Legarraga P, Iruretagoyena M, Pizarro G, Vollrath V, Araos R, et al. Head-to-head comparison of four antigen-based rapid detection tests for the diagnosis of SARS-CoV-2 in respiratory samples. bioRxiv. 2020 <https://www.biorxiv.org/content/10.1101/2020.05.27.119255v1>
7. Advice on the use of point-of-care immunodiagnostic tests for COVID-19: scientific brief [Accessed 16 July 2020]. Available from: <https://www.who.int/publications-detail-redirect/advice-on-the-use-of-point-of-care-immunodiagnostic-tests-for-covid-19-scientific-brief>.

Ester Parada-Ricart^{a,b,*}, Frederic Gomez-Bertomeu^{b,c},
Ester Picó-Plana^c, Montserrat Olona-Cabases^{b,d}

^a Servicio de Pediatría, Hospital Universitari de Tarragona Joan XXIII, Tarragona, Spain

^b Facultat Medicina y Cirugía, Universitat Rovira i Virgili, Tarragona, Spain

^c Àrea de Microbiologia, Laboratori Clínico ICS Camp de Tarragona-Terres de l'Ebre Hospital Universitari de Tarragona Joan XXIII, Tarragona, Spain

^d Unidad de Medicina Preventiva y Epidemiología, Hospital Universitari de Tarragona Joan XXIII, Tarragona, Spain

* Corresponding author.

E-mail address: eparada.hj23.ics@gencat.cat (E. Parada-Ricart).

<https://doi.org/10.1016/j.eimce.2021.05.003>

2529-993X/ © 2020 Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. Published by Elsevier España, S.L.U. All rights reserved.

Malignant external otitis by *Aspergillus flavus*



Otitis externa maligna por *Aspergillus flavus*

Malignant or invasive external otitis (MEO) is a rare disease. It is mainly caused by *Pseudomonas aeruginosa* and typically affects elderly patients with diabetes mellitus (DM) or other immunocompromised individuals.¹

Aspergillus MEO frequently involves necrosis of the epithelium and subepithelial tissue of the external auditory canal, bone erosion, and perforation of the tympanic membrane.²

An 80-year-old patient with a history of high blood pressure and DM was referred to the ENT department for persistent pain in his left ear for 4 months. At the initial evaluation, only medial otorrhea was found and antibiotic drops were prescribed. The ear pain persisted and extended to the left side of the patient's face and left buccal mucosa. He was treated with carbamazepine and tramadol, with no response. At the next visit, he presented with edema of the external auditory canal, left hearing loss, and a tumor-

fraction at the left temporomandibular joint with normal cranial nerve examination results.

A cerebral magnetic resonance imaging (MRI) was performed, with images suggestive of left MEO, inflammatory changes in the middle ear and deep cervical spaces, and a small abscess in the chewing space, osteomyelitis in the left clivus and left occipital condyle (Fig. 1A).

Treatment with intravenous piperacillin/tazobactam and vancomycin was prescribed. A computed tomography (CT) scan of the skull base showed that soft tissue occupied the external auditory canal and middle ear, and it also showed erosion and dehiscence of the tegmen timpani, mastoid sclerosis, and bone erosion in the left jugular foramen, front and lateral wall of occipital condyle, and carotid duct.

A biopsy of the external auditory canal was conducted. Histologically, granulation tissue by non-specific inflammation was observed. *A. flavus* were found in culture, with no malignancy findings. *A. flavus* CMI to voriconazole was 0.25 µg/mL. Antibiotic treatment was simplified to ciprofloxacin and voriconazole, which




Fig. 1. (A) Coronal T1 sequence, gadolinium. Thickening of the left external auditory canal (blue star) and soft tissues (orange star), parotid gland (green star), temporomandibular joint (straight arrow), occipital condyle (red star) and middle ear (curved arrow). (B) Similar section after 18 m. Clear improvement, with disappearance of the occupation of the middle ear and decrease in the volume of the tissues originally affected. Unspecific enhancement at the jugular foramen (red arrow).