



# Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



## Brief report

### Pacemaker-induced endocarditis by *Gordonia bronchialis*

Saray Mormeneo Bayo<sup>a,\*</sup>, María Pilar Palacián Ruíz<sup>a</sup>, Uxua Asin Samper<sup>b</sup>,  
María Isabel Millán Lou<sup>a</sup>, Ascensión Pascual Catalán<sup>b</sup>, María Cruz Villuendas Usón<sup>a</sup>

<sup>a</sup> Servicio de Microbiología, Hospital Universitario Miguel Servet Zaragoza, Spain

<sup>b</sup> Servicio de Medicina Interna y Enfermedades Infecciosas, Hospital Universitario Miguel Servet Zaragoza, Spain

## ARTICLE INFO

### Article history:

Received 17 July 2020

Accepted 3 November 2020

### Keywords:

*Gordonia*  
Endocarditis  
Pacemaker  
Bacteremia  
Actinomycetales

### Palabras clave:

*Gordonia*  
Endocarditis  
Marcapasos  
Bacteremia  
Actinomycetales

## ABSTRACT

**Purpose:** *Gordonia* species are known to be opportunistic human pathogens causing secondary infections. We present the second case in the world of endocarditis caused by *Gordonia bronchialis* and a review of all the cases of endocarditis caused by *Gordonia* spp.

**Methods:** The identification was performed by matrix-assisted desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) and 16S rRNA gene sequencing were performed to confirm the identification. Antimicrobial susceptibility was performed by MIC test Strip on Mueller-Hinton agar supplemented with 5% defibrinated sheep blood according to Clinical and Laboratory Standards Institute.

**Results:** Pacemaker-induced endocarditis due to *Gordonia bronchialis* infection was determined in an 88-year old woman. The patient was treated with ceftriaxone and ciprofloxacin until completing 6 weeks from the pacemaker explant with a good evolution.

**Conclusion:** The case presented supports the pathogenic role of *Gordonia bronchialis* as an opportunistic pathogen and highlights the high risk of suffering infections caused by environmental bacteria.

© 2020 Sociedad Española de

Enfermedades Infecciosas y Microbiología Clínica. Published by Elsevier España, S.L.U. All rights reserved.

### Endocarditis del marcapasos debido a infección por *Gordonia bronchialis*

## RESUMEN

**Objetivo:** Las especies de *Gordonia* son patógenos humanos oportunistas que causan infecciones secundarias. Presentamos el segundo caso en el mundo de endocarditis causada por *Gordonia bronchialis*, así como una revisión de todos los casos de endocarditis causados por *Gordonia* spp.

**Métodos:** La identificación fue realizada mediante espectrometría de masas MALDI-TOF MS, y se confirmó mediante secuenciación del gen 16S rRNA. La susceptibilidad antimicrobiana se realizó mediante tiras reactivas MIC en agar Müller-Hinton suplementado con un 5% de sangre ovina desfibrinada, conforme al Clinical and Laboratory Standards Institute (CLSI).

**Resultados:** La endocarditis del marcapasos debido a infección por *Gordonia bronchialis* se encontró en una mujer de 88 años. La paciente fue tratada con ceftriaxona y ciprofloxacina hasta completar el periodo de 6 semanas desde el explante del marcapasos, con buena evolución.

**Conclusión:** Este caso respalda el rol patogénico de *Gordonia bronchialis* como patógeno oportunista, subrayando el alto riesgo de padecer infecciones causadas por bacterias ambientales.

© 2020 Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

## Introduction

Infective endocarditis is a serious disease with a high mortality in the short and long terms. A special situation of endocarditis that has grown in parallel with the more frequent use of implantable

\* Corresponding author.

E-mail address: samorbayo@hotmail.com (S. Mormeneo Bayo).

defibrillator, cardiac resynchronization therapy and pacemaker implantation is infection developed on these devices<sup>1</sup>.

Members of the genus *Gordonia* are aerobic, catalase-positive, Gram-positive, nonmotile, nocardioform actinomycetes that are weakly acid-fast. The name for this genus, initially called *Gordona*, was chosen by Tsukamura in 1971 to pay tribute to the American bacteriologist Ruth E. Gordon<sup>2</sup>.

*Gordonia* genus comprises 39 different species that are recognized for symbiotic associations with multiple hosts allowing its use in bioremediation. Some *Gordonia* species (including *Gordonia bronchialis*) isolated from clinical specimens are known to be opportunistic human pathogens causing secondary infections in immunocompetent and immunosuppressive individuals<sup>3</sup>.

Here, we present the second case in the world of endocarditis caused by *Gordonia bronchialis* in a patient with implanted pacemaker. Our main purpose is to show our experience and describe all the cases published to date of endocarditis caused by *Gordonia* spp.

## Material and methods

The identification was performed by matrix-assisted desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) (Maldi Biotyper® Bruker Daltonics) and 16S rRNA gene sequencing were performed to confirm the identification. The DNA was extracted using the PrepMan DNA Kit method (Applied Biosystems,) following the manufacturer's recommendations. Bacterial universal PCR was performed systematically on sample with primers 16S-F (5'-AGA GTT TGA TCC TGG CTC AG-3') and 16S-R (5'-ACG GCT ACC TTG TTA CGA CTT-3'), producing an amplicon of approximately 1500 base pairs. The sequences obtained were compared with those stored in GenBank using BLAST software (<http://www.ncbi.nlm.nih.gov/blast>).

Antimicrobial susceptibility was performed by MIC test Strip (Liofilchem®) on Mueller-Hinton agar supplemented with 5% defibrinated sheep blood according to Clinical and Laboratory Standards Institute (CLSI) criteria for aerobic actinomycetes<sup>4</sup>.

## Case report

An 88-year old woman with chronic atrial fibrillation developed an atrioventricular block 20 years ago, with the implantation of a pacemaker. Generator was replaced in 2005 and 2018. One month after the second exchange, the patient began with erythema and itching on the generator wound and spontaneous drainage of hemopurulent material, exudate from the surgical wound was cultivated, growing a Gram-positive bacilli identified as *Gordonia bronchialis* beginning treatment with ciprofloxacin. The patient undergoes a transthoracic echocardiogram without signs of infective endocarditis.

Given the persistence of signs of local infection and fever, the patient undergone a pace-maker and electrodes (located in right ventricle) exchange surgery. The explanted pacemaker (generator, pacemaker pocket and pacemaker leads) was sent to the Microbiology Department. Next day, the patient has fever so blood cultures and a new transesophageal echocardiogram was performed, showing vegetations in the tricuspid septal veil and pacemaker electrode, so pace-maker explant was performed by cardiac surgeon, requesting cultures of the pacemaker.

Aerobic, anaerobic and fungal culture was performed and there was growth of *Gordonia bronchialis* (except in the generator) after 48 h of incubation at 37 °C in blood and chocolate agar. Given the isolation of *Gordonia* in the first explanted system, it was decided to extend the incubation time of blood cultures from 5 days to 14 days being positive at 11 days and 18 h for *Gordonia bronchialis*.

So it was a pacemaker-associated endocarditis, ceftriaxone 2 g and ciprofloxacin 400 mg/12 h were combined. The second pacemaker explanted culture was negative.

Control blood cultures were performed at 72 h, being negative. Intravenous treatment with ceftriaxone and ciprofloxacin was maintained until completing 6 weeks from the pacemaker explant.

Upon discharge, a new transthoracic echocardiogram was performed in which no endocarditis signs were observed. Since the patient was aged, dependent for basic activities and no atrial-block or bradiarrhythmia was observed in subsequent medical visits so no pace-maker implant took place. The patient remained stable, presenting progressive normalization of acute phase reactants, asymptomatic and without presenting recurrences until the present. No control blood culture was performed after the resolution of clinical presentation.

The identification of the microorganism by MALDI-TOF resulting in *Gordonia bronchialis* with a score of 1.68. Identification was repeated with previous extraction with ethanol-formic recommended by the manufacturer, improving to 2.08 score. The sequence was 99% identical to *Gordonia bronchialis* type strain DMS 43247 using the NCBI 16SrRNA gene database.

The strain was sensitive to all the antibiotics tested (amoxicillin-clavulanate, ceftriaxone, ciprofloxacin, imipenem, tobramycin, amikacin, linezolid and trimethoprim-sulfamethoxazole).

## Discussion

*Gordonia bronchialis* is an emerging opportunistic pathogen widely distributed in the environment. The genus *Gordonia* is able to form biofilm by producing gordonan, and acidic cell aggregation-inducing polysaccharide and consequently is responsible for persistent infections and need for prolonged treatments<sup>5</sup>.

*Gordonia bronchialis* is involved in bacteraemia and sequestered lung<sup>6</sup>, recurrent breast abscess<sup>7</sup>, pleural infection<sup>8</sup>, osteomyelitis<sup>9</sup>, sternal infection<sup>10–14</sup>, pacemaker-induced endocarditis<sup>15</sup>, cutaneous abscess<sup>16</sup>, peritoneal dialysis associated peritonitis<sup>17</sup> and endophthalmitis<sup>18</sup>. To our knowledge, we describe the second case in the world of endocarditis caused by *Gordonia bronchialis* and the fourth case of endocarditis caused by *Gordonia* spp since two other cases produced by *Gordonia polyisoprenivorans*<sup>19</sup> and *Gordonia* sp. most closely resembling *Gordonia sputi*<sup>20</sup> have been described. In these last two cases, endocarditis was caused by bacteremia associated with long-standing central catheter, in our case as well as in the case of Titécát M et al., endocarditis was caused by infection of the pacemaker system. The main characteristics of these 4 cases are described in Table 1.

With the advancement of molecular techniques, the number of recognized species of aerobic actinomycetes is increasing, and differentiation between related species is allowed. In our case, the identification by MALDI-TOF achieved a high score after extraction with ethanol-formic acid. However, correct identification is not always possible as it is not in the database of some automated systems and for its morphological similarity of the colonies with other species. In fact, in the other three published articles<sup>15,19,20</sup> in which endocarditis occurred, the definitive identification required the 16S rRNA.

Regarding antibiotic treatment, there are currently no standardized guidelines, there are numerous combinations of antibiotics described in the literature. The combination of vancomycin and ceftazidime, trimethoprim-sulfamethoxazole and imipenem, piperacilia-tazobactam and daptomycin among others have been described. Later, according to evolution, the treatment has been sequenced to ciprofloxacin, linezolid and amoxicillin. A case has been described in which ceftaroline monotherapy has been used<sup>21</sup>.

**Table 1**  
Reported infective endocarditis due to *Gordonia* spp.

Reference	Sex (F/M) Age (years)	Underlying condition	Source of infection	Echocardiogram	Treatment	Outcome
Lesens O et al. (2000)	F/31	Splenectomy, beta-thalassemia and haemoglobin E disease. Hepatitis C with cirrhosis and secondary hemochromatosis, treated with deferoxamine.	Catheter-related bacteremia	Native mitral valve vegetation.	i.v. amoxicillin + i.v. netilmicin (1 week). Then i.v. amoxicillin alone (3 weeks) followed by ceftriaxone for 2 weeks.	Cure
Verma P et al. (2006)	M/78	Osler-Weber-Rendu and myelodysplastic syndrome with pancytopenia	Catheter-related bacteremia	Native mitral and aortic valve vegetation.	I.v. ampicillin + gentamicin. After, imipenem and amikacin (unspecified duration).	Died
Titécát M et al. (2014)	M/92	High blood pressure, atrial fibrillation, pulmonary embolism	Pacemaker	Vegetation on the ventricular lead in the right auricle.	Piperacillin-tazobactam + daptomycin (6 days), i.v. amoxicillin (6 weeks)	Cure
Our case	F/88	Atrial fibrillation	Pacemaker	Endocarditis in the tricuspid septal veil and pacemaker electrode	Ceftriaxone + ciprofloxacin (6 weeks)	Cure

The case presented supports the pathogenic role of *Gordonia bronchialis* as an opportunistic pathogen and highlights the high risk of suffering infections caused by environmental bacteria and the importance of molecular analyses in the definitive identification of these and other related species.

### Conflict of interests

The authors declare that they have no conflict of interest.

### References

- Carrasco F, Anguita M, Ruiz M, Castillo JC, Delgado M, Mesa D, et al. Clinical features and changes in epidemiology of infective endocarditis on pacemaker devices over a 27-year period (1987–2013). *Europace*. 2016;18:836–41.
- Tsukamura M. Proposal of a new genus, *Gordonia*, for slightly acid-fast organisms occurring in sputa of patients with pulmonary disease and in soil. *J Gen Microbiol*. 1971;68:15–26.
- Sowani H, Kulkarni M, Zinjarde S. An insight into the ecology, diversity and adaptations of *Gordonia* species. *Crit Rev Microbiol*. 2018;44:393–413.
- Clinical and Laboratory Standards Institute. *Susceptibility Testing of Mycobacteria, Nocardiae, and Other Aerobic Actinomycetes* – Second Edition: approved standard (M24-A2). Wayne, PA: The Institute; 2011.
- Kondo T, Yamamoto D, Yokota A, Suzuki A, Nagasawa H, Sakuda S. *Gordonan*, an acidic polysaccharide with cell aggregation-inducing activity in insect BM-N4 cells, produced by *Gordonia* sp. *Biosci Biotechnol Biochem*. 2000;64:2388–94.
- Sng LH, Koh TH, Toney SR, Floyd M, Butler WR, Tan BH. Bacteremia caused by *Gordonia bronchialis* in a patient with sequestered lung. *J Clin Microbiol*. 2004;42:2870–1.
- Werno AM, Anderson TP, Chambers ST, Laird HM, Murdoch DR. Recurrent breast abscess caused by *Gordonia bronchialis* in an immunocompetent patient. *J Clin Microbiol*. 2005;43:3009–10.
- Johnson JA, Onderdonk AB, Cosimi LA, Yawetz S, Lasker BA, Bolcen SJ, et al. *Gordonia bronchialis* bacteremia and pleural infection: case report and review of the literature. *J Clin Microbiol*. 2011;49:1662–6.
- Siddiqui N, Toumeh A, Georgescu C. Tibial osteomyelitis caused by *Gordonia bronchialis* in an immunocompetent patient. *J Clin Microbiol*. 2012;50:3119–.
- Wright SN, Gerry JS, Busowski MT, Klochko AY, McNulty SG, Brown SA, et al. *Gordonia bronchialis* sternal wound infection in 3 patients following open heart surgery: intraoperative transmission from a healthcare worker. *Infect Control Hosp Epidemiol*. 2012;33:1238–41.
- Vasquez MA, Marne C, Villuendas MC, Arazo P. Osteomyelitis esternal subaguda por *Gordonia bronchialis* tras cirugía cardíaca. *Enferm Infecc Microbiol Clin*. 2013;31:559–60.
- Chang JH, Ji M, Hong HL, Choi SH, Kim YS, Chung CH, et al. Sternal osteomyelitis caused by *Gordonia bronchialis* after open-heart surgery. *Infect Chemother*. 2014;46:110–4.
- Rodríguez-Lozano J, Pérez-Llantada E, Agüero J, Rodríguez-Fernández A, De Alegría CR, Martínez-Martínez L, et al. Sternal wound infection caused by *Gordonia bronchialis*: identification by MALDI-TOF MS. *JMM Case Rep*. 2016;3:e005067.
- Ambesh P, Kapoor A, Kazmi DH, Elsheshtawy M, Shetty V, Lin YS, et al. Sternal osteomyelitis by *Gordonia bronchialis* in an immunocompetent patient after open heart surgery. *Ann Card Anaesth*. 2019;22:221–4.
- Titécát M, Loiez C, Courcol RJ, Wallet F. Difficulty with *Gordonia bronchialis* identification by Microflex mass spectrometer in a pacemaker-induced endocarditis. *JMM Case Rep*. 2014;1:1–4.
- Bartolomé-Álvarez J, Sáez-Nieto JA, Escudero-Jiménez A, Barba-Rodríguez N, Galán-Ros J, Carrasco G, et al. Cutaneous abscess due to *Gordonia bronchialis*: case report and literature review. *Rev Esp Quimioter*. 2016;29:170–3.
- Sukackiene D, Rimsevicius L, Kiveryte S, Marcinkeviciene K, Bratchikov M, Zokaityte D, et al. A case of successfully treated relapsing peritoneal dialysis-associated peritonitis caused by *Gordonia bronchialis* in a farmer. *Nephrol Ther*. 2018;14:109–11.
- Choi R, Strnad L, Flaxel CJ, Lauer AK, Suhler EB. *Gordonia bronchialis*-associated endophthalmitis, Oregon, USA. *Emerg Infect Dis*. 2019;25:1017–9.
- Verma P, Brown JM, Nunez VH, Morey RE, Steigerwalt AG, Pellegrini GJ, et al. Native valve endocarditis due to *Gordonia polyisoprenivorans*: case report and review of literature of bloodstream infections caused by *Gordonia* species. *J Clin Microbiol*. 2006;44:1905–8.
- Lesens O, Hansmann Y, Riegel P, Heller R, Benaissa-Djellouli M, Martinot M, et al. Bacteremia and endocarditis caused by a *Gordonia* species in a patient with a central venous catheter. *Emerg Infect Dis*. 2000;6:382–5.
- Akrami K, Coletta J, Mehta S, Fierer J. *Gordonia* sternal wound infection treated with ceftaroline: case report and literature review. *JMM Case Rep*. 2017;4:e005113.