

# Effects of aerobic exercise on cognitive function in adults with major depressive disorder: A systematic review and meta-analysis

Fei-Fei Ren<sup>a</sup>, Charles H. Hillman<sup>b,c</sup>, Wei-Guang Wang<sup>a</sup>, Ruei-Hong Li<sup>d</sup>, Wen-Sheng Zhou<sup>e</sup>, Wen-Ming Liang<sup>f</sup>, Yong Yang<sup>g</sup>, Feng-Tzu Chen<sup>h,\*\*</sup>, Yu-Kai Chang<sup>d,i,j,\*</sup>

<sup>a</sup> Department of Physical Education, Beijing Language and Culture University, Beijing, China

<sup>b</sup> Department of Psychology, Northeastern University, Boston, MA, USA

<sup>c</sup> Department of Physical Therapy, Movement, and Rehabilitation, Northeastern University, Boston, MA, USA

<sup>d</sup> Department of Physical Education and Sport Sciences, National Taiwan Normal University, Taipei, Taiwan

<sup>e</sup> Department of Physical Education, Jiangsu Second Normal University, Jiangsu, China

<sup>f</sup> Life Sciences Center, Vilnius University, Vilnius, Lithuania

<sup>g</sup> Laboratory of Kinesiology and Rehabilitation, School of Physical Education and Sport, Chaohu University, Anhui, China

<sup>h</sup> Department of Kinesiology, National Tsing Hua University, Hsinchu, Taiwan

<sup>i</sup> Social Emotional Education and Development Center, National Taiwan Normal University, Taipei, Taiwan

<sup>j</sup> Institute for Research Excellence in Learning Science, National Taiwan Normal University, Taipei, Taiwan

## ARTICLE INFO

### Keywords:

Physical activity  
Cognition  
Memory  
Executive function  
Depression

## ABSTRACT

**Background:** Major Depressive Disorder (MDD) is a highly prevalent psychiatric disorder that impairs the cognitive function of individuals. Aerobic exercise stands out as a promising non-pharmacological intervention for enhancing cognitive function and promoting brain health.

While positive impacts of aerobic exercise on executive function in adults with depression have been documented, a comprehensive understanding of its benefits on overall cognitive function, including memory, attention, and processing speed, along with key moderating factors in adults with MDD, remains unexplored. The purpose of the systematic review and meta-analysis was to investigate the effects of aerobic exercise on overall cognitive function in adults with MDD, and to explore whether cognitive sub-domains, aerobic exercise characteristics, and study and sample variables modify the effects of aerobic exercise on cognition.

**Methods:** Six English electronic databases (Embase, Cochrane Central, Scopus, APA PsycInfo, PubMed, Web of Science) were searched from inception to 2 April 2023. Randomized trials, including adults aged 18 years or above with a diagnosis of clinical depression, of the effects of aerobic exercise on cognitive function in adults with MDD compared to non-aerobic exercise groups were included. A three-level meta-analysis was conducted utilizing a random-effects model in R. The quality of the studies was evaluated using the Physiotherapy Evidence Database (PEDro) scale. The PROSPERO registration number is CRD42022367350.

**Results:** Twelve randomized trials including 945 adults with MDD were included. Results indicated that aerobic exercise significantly improved overall cognitive function ( $g = 0.21$ ; 95 % confidence intervals [CI] = 0.07, 0.34), and the sub-domains of memory ( $g = 0.25$ ; 95 % CI = 0.06, 0.44) and executive function ( $g = 0.12$ ; 95 % CI = 0.04, 0.20). Significant benefits in cognitive function were found from moderate-to-vigorous (mixed) intensity ( $g = 0.19$ ; 95 % CI = 0.02, 0.37), aerobic exercise conducted 3 times per week ( $g = 0.23$ ; 95 % CI = 0.10, 0.38), in sessions < 45 min ( $g = 0.59$ ; 95 % CI = 0.28, 0.90), and 45–60 min ( $g = 0.16$ ; 95 % CI = 0.07, 0.26), in aerobic exercise intervention  $\leq 12$  weeks ( $g = 0.26$ ; 95 % CI = 0.08, 0.44).

**Limitations:** This review only included peer-reviewed English-language studies, which may lead to a language bias. The results of the Egger's test suggested a potential publication bias.

**Conclusions:** Aerobic exercise is efficacious in improving overall cognitive function and the sub-domains of memory and executive function in adults with major depressive disorder.

\* Corresponding author at: Department of Physical Education and Sport Sciences, National Taiwan Normal University, Taipei, Taiwan.

\*\* Corresponding author at: Department of Kinesiology, National Tsing Hua University, Hsinchu, Taiwan.

E-mail addresses: [alexnewtaipei@gmail.com](mailto:alexnewtaipei@gmail.com) (F.-T. Chen), [yukaichangnew@gmail.com](mailto:yukaichangnew@gmail.com) (Y.-K. Chang).

<https://doi.org/10.1016/j.ijchp.2024.100447>

Received 7 August 2023; Accepted 30 January 2024

Available online 9 February 2024

1697-2600/© 2024 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Introduction

Depression has emerged as a leading cause of worldwide illness burden, disability, and suicide (Birnie et al., 2022; Kawilapat et al., 2022; Li et al., 2023). However, roughly two-thirds of adults diagnosed with depression do not engage in therapy actively (Heissel et al., 2023). Major Depressive Disorder (MDD), as one of the costliest psychiatric disorders (Patel et al., 2023), is a devastating, recurrent, and highly prevalent psychiatric disorder (Lan et al., 2023; Lee et al., 2020) affecting approximately 5 % of adults worldwide (Kowalec et al., 2022). MDD is responsible for an annual economic loss of billions of dollars globally (Greenberg et al., 2021; Sepehrmanesh et al., 2017), and is associated with a lower quality of life (Hammar et al., 2022; Irwin et al., 2023), poor academic performance, marital issues, and a higher rate of unemployment and absenteeism (Iancu et al., 2020; Lépine & Briley, 2011; Sepehrmanesh et al., 2017). MDD symptoms include negative emotions, hopelessness (Chai et al., 2019), insomnia, suicidal ideation, and cognitive dysfunction (Zhao et al., 2020).

Cognitive impairments have been widely reported to be a core deficit of MDD individuals (Kriesche et al., 2022; Pan et al., 2019), and meta-analyses have demonstrated that these individuals have deficits in several cognitive domains, such as processing speed, attention, memory, and executive function (EF) (Kriesche et al., 2022; Rock et al., 2014; Snyder, 2013). There is evidence that cognitive impairment is observed in both the acute and remission phases of depression (Kriesche et al., 2022; Legemaat et al., 2022), and cognitive impairment may deteriorate over time and with recurrent episodes when untreated (Hammar et al., 2022; Maramis et al., 2021). Moreover, cognitive impairment negatively impacts quality of life, social, interpersonal functioning (Hammar et al., 2022), daily activities, education and work (Keefe et al., 2022) in individuals with depression. Antidepressants, despite being the first-line treatment for MDD (Zhdanov et al., 2020), have shown minimal and limited effects on cognitive function (Rosenblat et al., 2016). In addition, antidepressants can cause side effects, such as nausea, headaches, and sleepiness (Du et al., 2020). Thus, non-pharmacological interventions to enhance cognitive function in MDD participants have scientific and practical relevance.

Aerobic exercise, a low-cost non-pharmacological treatment, has been shown to reduce depressive symptoms (standardized mean difference [SMD] = -1.156;  $p < 0.001$ ) in adults (18 years or older) with diagnosed or indicated depression (Heissel et al., 2023) and alleviate cognitive fatigue (Zhang et al., 2023). Further, aerobic exercise has also shown significant benefits to cognitive function in the healthy population (McSween et al., 2019; Nanda et al., 2013), people with Alzheimer's Disease (Zhang et al., 2022), participants with schizophrenia (Shimada et al., 2022), children with preterm birth (Ren, Feng, et al., 2023), and adults aged 50 years and older (Northey et al., 2018). Several randomized trials have suggested that aerobic exercise produces positive effects on attention (Buschert et al., 2018), memory (Buschert et al., 2018; Khatri et al., 2001; Krogh et al., 2012), and executive function (Imboden et al., 2020; Khatri et al., 2001) in participants with MDD. However, other studies have suggested that aerobic exercise does not have a significant effect on cognitive outcomes in participants with MDD (Brush et al., 2020; Krogh et al., 2009). In addition, several meta-analyses have explored the effects of physical exercise on cognitive function (Brondino et al., 2017; Sun et al., 2018) or executive function (Contreras-Osorio et al., 2022; Ren, Alderman, et al., 2023) in depressed adults. Brondino et al. (2017) found that exercise did not enhance processing speed, attention/vigilance, global cognition, verbal learning and memory, reasoning/problem-solving, and working memory in depressed adults (Brondino et al., 2017). Further, Sun et al. (2018) also found that exercise did not significantly improve these same outcomes; however, interventions that included both physical and cognitive activities, and exercise with low-intensity did produce significant improvements in cognition (Sun et al., 2018). Contreras-Osorio et al. (2022) suggested that exercise selectively improved working memory (effect size [ES] =

0.33), but was unrelated to inhibitory control and cognitive flexibility in adults with depression (Contreras-Osorio et al., 2022), suggesting differential effects across the different aspects of executive function. However, Ren et al. (2023) found that aerobic exercise broadly improved executive function (ES = 0.203) in adults with depression (Ren, Alderman, et al., 2023). In general, the findings from meta-analyses on the effects of exercise on cognitive function in depressed adults are mixed, and few meta-analyses have specifically focused on aerobic exercise. While randomized trials of different modes of exercise on cognitive function in depression have grown in recent years, most trials have used aerobic exercise as an intervention. Synthesizing results support the notion that aerobic exercise improves EF in this population. However, the effects of aerobic exercise on overall cognitive function and other cognitive sub-domains by meta-analysis remain unknown. Given the conflicting results observed in randomized trials regarding the effects of aerobic exercise on cognitive function in adults with MDD, it is necessary to conduct a comprehensive examination of the effects of aerobic exercise on overall cognitive function and its sub-domains of memory, attention, and processing speed to fill this gap. More importantly, the detailed prescription (frequency, intensity, session time, duration) for the effect of aerobic exercise on cognitive function in adults with MDD has not been determined. This may hinder the development of optimal aerobic exercise prescriptions. In addition, important moderators between aerobic exercise and cognition have not been examined, such as the characteristics of study design (aerobic exercise plus antidepressants; comparator/control group), and sample characteristics (proportion of female participants; mean age; clinical setting). In fact, these moderators are crucial factors to consider when scientific researchers try to understand how aerobic exercise improves cognitive function in adults with depression.

This review addresses the disparate outcomes in randomized trials examining aerobic exercise impact on cognitive function in MDD. Given the absence of comprehensive evidence from systematic reviews and meta-analyses, our objective is to scrutinize aerobic exercise effects on overall cognitive function. Additionally, we aim to explore potential moderation in cognitive sub-domains, exercise characteristics, and study/sample variables. This meta-analysis serves as a theoretical foundation for clinicians and depressed patients considering aerobic exercise as a non-pharmacological intervention.

## Methods

### Design and eligibility criteria

The meta-analysis followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Page et al., 2021). Its protocol was preregistered in the International Prospective Register of Systematic Reviews (PROSPERO) on 24 October 2022 (registration number CRD42022367350).

The criteria for inclusion were: (1) participants were adults (aged 18 years or older) with a diagnosis of MDD using recognized or established diagnostic criteria (e.g. the International Classification of Diseases, Tenth Revision (ICD-10), the Diagnostic and Statistical Manual of Mental Disorders, (DSM-IV or DSM-5), or the Mini-International neuropsychiatric interview (MINI), and by applying a cut-off on reliable depression scales (such as Beck Depression Inventory, Beck Depression Inventory-II, the Montgomery-Åsberg Depression Rating Scale, and Hamilton Depression Rating Scale); (2) studies employed aerobic exercise as the experimental intervention, or aerobic exercise as main intervention combined with other intervention (such as antidepressant); (3) studies included a comparator/control conditions that performed non-aerobic exercises, standard care, active control, placebo pill, no treatment, or a wait list; (4) studies included an overall cognitive function task, and at least one measure of a cognitive sub-domain such as attention, processing speed, memory, and executive function; and (5) randomized trials published in English-language peer-reviewed

journals.

The exclusion criteria were: (1) studies that employed a single bout of (acute) aerobic exercise; (2) experimental group used mind-body exercises (Yoga, Tai-Chi, and Qigong), resistance exercise or aerobic exercise combined with resistance exercise or multicomponent exercise; (3) both experimental and comparator groups used aerobic exercise; and (4) studies in which data could not be extracted for ES estimation even after contacting the authors.

#### Literature search and selection

Six English electronic databases (Embase, Cochrane Central, Scopus, APA PsycInfo, PubMed, Web of Science) were searched from inception to 12 October 2022, and an updated search was conducted on 2 April 2023. The search term (i.e., keywords) included medical subject headings ("depression" AND "aerobic exercise" AND "cognition" OR "executive function" AND "randomized controlled trial" [Publication Type]) and related free terms (see Supplementary Table 1). After removing duplicates, two authors (FFR and FTC) independently reviewed the titles and abstracts of the remaining literature. Next, the same two authors independently read and identified the full-text of studies that had been preliminarily accepted. In instances when the two authors disagreed on which studies should be included, a third author (YKC) was consulted to reach consensus.

#### Data extraction and coding strategy

Cochrane Collaboration Handbook methods and strategies were followed throughout the data extraction process (Higgins et al., 2019). Two authors (FFR and FTC) extracted data separately by using a pre-developed extraction form in Microsoft Excel. Extracted data and information included: first author, year of publication, depression diagnostic criteria and instruments, characteristics of groups and participants (including sample size of experimental and control groups, mean age, gender, severity of depression, clinical setting, percentage of antidepressant use of exercise group, exercise frequency, exercise intensity, session time of exercise, duration of exercise, cognitive function domains and tasks, mean values and standard deviations of post-test outcome. If raw data were unavailable or inaccessible in the included studies, FFR would contact the authors. When an included study presented several relevant outcome measures on the same cognitive sub-domain, all of them were included.

The cognitive sub-domains were categorized according to their features and previous meta-analysis (Northey et al., 2018; Sanders et al., 2019), these sub-domains included attention, processing speed, memory, and executive function (which together comprise overall cognitive function). The characteristics of aerobic exercise were coded: (1) intensity (moderate only, and moderate-to-vigorous (mixed)); (2) frequency (2 times per week, and 3 times per week); (3) session time (< 45 min, 45–60 min, and > 60 min) (Chen et al., 2020); (4) duration ( $\leq 12$  weeks, and > 12 weeks). Study and sample characteristics were coded: (1) comparator/control group (active control group, placebo pill, and treatment as usual care), (2) exercise plus antidepressant (NO: participants of aerobic exercise group did not use antidepressant, YES: more than 95 % participants of aerobic exercise group used antidepressant, and MIXED: some participants of aerobic exercise group used antidepressant); (3) proportion of female (> 50 %,  $\leq 50$  %, and unclear); (4) mean age (young adults (18–44 years), middle adults (45–64 years), old adults ( $\geq 65$  years), and unclear); (5) clinical setting (inpatients, and outpatients) (Krogh et al., 2017; Lee et al., 2021).

#### Assessment of study quality

Two authors (WSZ and FFR) independently assessed the study quality of all included studies using PEDro scale (Maher et al., 2003). Study quality was ranked as high (excellent: scoring 9–10 points), (good:

scoring 6–8 points), moderate (scoring 4–5 points), and poor (scoring 0–3 points). If the two authors disagreed, the third author (YKC) was engaged to reach a consensus.

#### Statistical analysis

This statistical analysis was conducted using R software (R Core Team, 2013; Viechtbauer, 2010). To manage the dependency of effect sizes within studies, a three-level meta-analysis with a random-effects model was performed. This model considers the following three categories of effect size (ES) variability: sampling variance (level 1); within-study variance (level 2); and between-study variance (level 3) (Cheung, 2014). Standardized mean difference and its variance were calculated based on post-test means, standard deviations, and sample sizes of the intervention and comparator/control groups. With the use of restricted maximum-likelihood estimation (REML) method, pooled ES of aerobic exercise on cognitive function was determined using Hedges'  $g$  and 95 % confidence intervals (CIs). A positive Hedges'  $g$  indicated that aerobic exercise had a favorable effect for cognitive function compared to the comparator/control group. The Hedges'  $g$  values were classified as follows:  $g = 0.20$  (small),  $g = 0.20$  to  $0.49$  (small-to-medium),  $g = 0.5$  to  $0.79$  (medium), and  $g \geq 0.80$  (large) (Cohen, 1992). Testing for statistical significance in level 2 (within-study) and level 3 (between-study) variances were performed using likelihood ratio tests (LRT). 95 % prediction intervals (PIs) were calculated to determine the expected range in which an ES in future identical studies will fall. Statistical significance was defined as a  $p$ -value < 0.05. The  $I^2$  statistic was used to estimate of total between-study heterogeneity (1–49 % indicating low, 50–74 % indicating moderate, and 75–100 % indicating high) (Higgins et al., 2003). Funnel plots and multilevel Egger's regression tests were used to examine the possibility of publication bias (Egger et al., 1997). Sensitivity analyses were conducted to investigate if outliers and influential studies affect pooled ES (Viechtbauer & Cheung, 2010).


The steps comprising the three-level meta-analysis were: (1) all cognitive tests were utilized to calculate the effect of aerobic exercise on overall cognitive function in adults with MDD; and (2) moderator analyses were performed to explore potential moderators of the observed effects. The subgroup analyses included cognitive sub-domains (attention, processing speed, memory, and executive function), exercise intensity, frequency, session time, duration, comparator/control group, exercise plus antidepressant or not, proportion of female, mean age, and clinical setting. The meta-regression analyses included the proportion of female, mean age, session time of exercise (per time), duration of exercise, volume of exercise per week, total intervention time, and sample size.

#### Results

Fig. 1 depicts the literature search and article selection procedure. The literature search yielded 18,031 records. Following the removal of duplicates ( $k = 8,157$ ), 9,874 studies met the criteria for title and abstract screening. Subsequently, 9,838 studies were excluded. The next step included reading the full text of the remaining 36 studies. Following this step, 12 studies met the inclusion criteria for this analysis. The first author of one article (Brush et al., 2020) provided more detailed data via direct email request.

#### Characteristics of included studies

Tables 1 and 2 provide the study characteristics of the 12 included randomized trials published between 2001 and 2023 (Brush et al., 2020; Buschert et al., 2018; Foley et al., 2008; Hoffman et al., 2008; Imboden et al., 2020; Khatri et al., 2001; Krogh et al., 2014; Krogh et al., 2009; Krogh et al., 2012; Neviani et al., 2017; Oertel-Knöchel et al., 2014; Olson et al., 2017). In total, 979 participants (678 females) with MDD (mean age from 20.23 to 75.20 years) are included in this review, and



**Fig. 1.** PRISMA flow chart of the literature search and selection procedure.

Note. n = number; PRISMA = Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

the number of female participants ranged from 11 to 153. However, 945 participants completed posttest and are thus included in the meta-analysis, with 437 in aerobic exercise group and 508 in comparator/control group. Depressive severity of participants ranged from mild-to-moderate (five studies), mild-to-severe (two studies), moderate-to-severe (one study), and unspecified (four studies). Six studies had 8.50 % to 64.60 % participants, four studies had 96.67 % to 100 % participants using antidepressant medications in the exercise group, and two studies did not use antidepressant medications in the exercise group. There were 9 studies that examined outpatients. Within the 12 included studies, diagnostic criteria for MDD were based on the DSM-IV in six studies, ICD-10 in three studies, and MINI in three studies. The scores of baseline depression were measured using the Beck Depression Inventory (BDI) or Beck Depression Inventory-II (BDI-II) in 10 studies, the Hamilton Depression Rating Scale (HAM-D or HDRS or HRSD) in 8 studies, and the Montgomery-Åsberg Depression Rating Scale (MADRAS) in one study. Regarding the comparator/control groups, ten studies used active comparators (e.g., stretching, occupational or art therapy, relaxation, and non-aerobic progressive physical activity such as flexibility, resistance training, and balance training), four studies used treatment as usual care (antidepressants, including one study that used antidepressants while serving as a waiting-list control), and one study used a placebo medication group. The frequency of

aerobic exercise ranged from 2 to 3 times per week, and each session time lasted from 30 to 90 min. Five studies utilized moderate only intensity, while another seven studies used moderate-to-vigorous (mixed) intensity exercise. Duration of the aerobic exercise intervention varied from 3 to 24 weeks. As for the cognitive sub-domains, six studies assessed processing speed, eight studies assessed memory, six studies assessed attention, and nine studies assessed executive function.

#### Methodological quality assessment

The results of the PEDro scale evaluation of 12 randomized studies revealed that two studies received 4–5 points, indicating moderate quality, while ten scored 6–8 points, indicating high quality. The included studies averaged 6.2 (see Supplementary Table 2).

#### Overall cognitive function

The results of this meta-analysis from the 12 included studies with 101 effect sizes indicated that the effect of aerobic exercise on overall cognitive function of adults with MDD was significant ( $g = 0.21$ ; 95 % CI = 0.07, 0.34;  $p = 0.003$ ) compared with the comparator/control group (see Fig. 2). The 95 % prediction intervals (PIs) suggested that a future ES will fall between -0.21 and 0.62 (Hedges'  $g$ ). The likelihood ratio tests

**Table 1**Overview of participants' characteristics of included studies ( $k = 12$ ).

| Author (Year)                | Sample size (N)                                                     | Age (mean or range) | Gender (N)                  | Depressive severity | Clinical setting | Antidepressant use in exercise group (%) | Diagnostic criteria/instruments  |
|------------------------------|---------------------------------------------------------------------|---------------------|-----------------------------|---------------------|------------------|------------------------------------------|----------------------------------|
| Brush et al. (2020)          | E: 35<br>C: 31<br>66 MDD                                            | 20.23               | M: 17<br>F: 49              | Mild-to-severe      | Outpatients      | 11.43 %                                  | MINI/<br>BDI-II                  |
| Buschert et al. (2018)       | E: 15<br>C: 15<br>30 MDD                                            | 47.37               | M: 11<br>F: 19              | Unspecified         | Inpatients       | 96.67 %                                  | ICD-10/<br>BDI-II;<br>HAMD       |
| Foley et al. (2008)          | E: 10<br><br>C: 13<br>23 MDD                                        | 18-55               | M:<br>Unclear<br>F: Unclear | Unspecified         | Outpatients      | 60 %                                     | DSM-IV/<br><br>BDI-II;<br>MADRAS |
| Hoffman et al. (2008)        | E: 104<br><br>C <sup>1</sup> : 49<br>C <sup>2</sup> : 49<br>202 MDD | 51.7                | M: 49<br><br>F: 153         | Mild-to-moderate    | Outpatients      | 0.00 %                                   | DSM-IV/<br><br>HAMD;<br>BDI-II   |
| Imboden et al. (2020)        | E: 22<br><br>C: 20<br>42 MDD                                        | 39.9                | M: 22<br><br>F: 20          | Moderate-to-severe  | Inpatients       | 100 %                                    | ICD-10/<br><br>HDRS;<br>BDI      |
| Khatri et al. (2001)         | E: 42<br>C: 42<br>84 MDD                                            | 56.73               | M: 20<br>F: 64              | Mild-to-severe      | Outpatients      | 0.00 %                                   | DSM-IV/<br>HAMD;<br>BDI-II       |
| Krogh et al. (2009)          | E: 55<br><br>C <sup>1</sup> : 55<br>C <sup>2</sup> : 55<br>165 MDD  | 38.9                | M: 43<br><br>F: 122         | Mild-to-moderate    | Outpatients      | 64.60 %                                  | ICD-10/<br><br>HRSD;<br>BDI      |
| Krogh et al. (2012)          | E: 56<br><br>C: 59<br>115 MDD                                       | 41.55               | M: 38<br><br>F: 77          | Mild-to-moderate    | Outpatients      | 8.50 %                                   | MINI/<br><br>HAMD;<br>BDI        |
| Krogh et al. (2014)          | E: 41<br><br>C: 38<br>79 MDD                                        | 41.3                | M: 26<br><br>F: 53          | Mild-to-moderate    | Outpatients      | 12.10 %                                  | MINI/<br><br>HAMD                |
| Neviani et al. (2017)        | E: 42<br><br>C <sup>1</sup> : 37<br>C <sup>2</sup> : 42<br>121 MDD  | 75.2                | M: 35<br><br>F: 86          | Mild-to-moderate    | Outpatients      | 100 %                                    | DSM-IV/<br><br>HAMD              |
| Oertel-Knochel et al. (2014) | E: 8<br>C <sup>1</sup> : 6<br>C <sup>2</sup> : 8<br>22 MDD          | 40                  | M: 11<br>F: 11              | Unspecified         | Inpatients       | 100 %                                    | DSM-IV/<br>BDI-II                |
| Olson et al. (2017)          | E: 15<br>C: 15<br>30 MDD                                            | 21.1                | M: 6<br>F: 24               | Unspecified         | Outpatients      | 14 %                                     | DSM-IV/<br>BDI-II                |

Note.  $k$  = number of included studies; N = number; % = percentage; E = experimental group; C = control group; C<sup>1</sup> = first control group; C<sup>2</sup> = second control group; MDD = Major depressive disorder; M = male; F = female; MINI = The Mini-International neuropsychiatric interview; DSM-5 = The Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition; DSM-IV = The Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition; ICD-10 = International Classification of Diseases, Tenth Revision; BDI = Beck Depression Inventory; BDI-II = Beck Depression Inventory-II; MADRAS = the Montgomery-Åsberg Depression Rating Scale; HAMD/HDRS/HRSD = Hamilton Depression Rating Scale; Diagnostic criteria/instruments = Diagnostic criteria and depression scales.

indicated that the influence of within-study variance was not statistically significant ( $LRT = 0$ ;  $p > 0.05$ ), while the effect of between-study variance was significant ( $LRT = 16.03$ ;  $p < 0.001$ ), explaining 44.74 % of the variance in level 3. Statistical heterogeneity in the pooled ES was low ( $I^2 = 44.74$  %).

### Moderator analyses

#### Cognitive sub-domains

Table 3 summarized the results of all subgroup analyses. The test for moderators indicated that there were no statistically significant subgroup differences in ES for cognitive sub-domains ( $F = 1.24$ ,  $p = 0.298$ ). The results indicated that aerobic exercise only had statistically significant effects on memory ( $g = 0.25$ ; 95 % CI = 0.06, 0.44;  $p = 0.012$ ; 95 % PIs = -0.20, 0.69) and executive function ( $g = 0.12$ ; 95 % CI = 0.04,

0.20;  $p = 0.005$ ; 95 % PIs = -0.02, 0.26) of adults with MDD compared with a comparator/control group. However, there were no significant effects on attention ( $g = 0.01$ ; 95 % CI = -0.12, 0.13;  $p = 0.928$ ; 95 % PIs = -0.12, 0.13) and processing speed ( $g = 0.08$ ; 95 % CI = -0.11, 0.26;  $p = 0.387$ ; 95 % PIs = -0.25, 0.40) following aerobic exercise.

#### Aerobic exercise characteristics

The results of the test of moderation for aerobic exercise characteristics revealed that there were no statistically significant subgroup differences in ES for exercise intensity ( $F = 0.11$ ,  $p = 0.735$ ), exercise frequency ( $F = 1.74$ ,  $p = 0.190$ ), and exercise duration ( $F = 0.83$ ,  $p = 0.364$ ), but there was a statistically significant subgroup differences in ES for session time ( $F = 5.36$ ,  $p = 0.006$ ). Studies with session time < 45 min yielded a larger effect ( $g = 0.59$ ; 95 % CI = 0.28, 0.90;  $p < 0.001$ ) than session time between 45–60 min ( $g = 0.16$ ; 95 % CI = 0.07, 0.26;  $p$

**Table 2**Overview of groups, characteristics of aerobic exercise and cognitive domains and tasks of included studies ( $k = 12$ ).

| Author (Year)                | Groups                                                                                                                   | Characteristics of aerobic exercise                                              | Cognitive domains and tasks                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|------------------------------|--------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Brush et al. (2020)          | E: Aerobic exercise<br>C: Stretching                                                                                     | F: 3 times/wk<br>I: 40–65 % HRR<br>T: 45 min<br>D: 8 wks                         | <b>PS:</b> Flanker Task-Congruent-Accuracy (FT-Con-Acc); Flanker Task-Congruent-Reaction Time (FT-Con-RT);<br><b>EF:</b> Flanker Task-Incongruent-Accuracy (FT-Incon-Acc); Flanker Task-Incongruent-Reaction Time (FT-Incon-RT)                                                                                                                                                                                                                                                                                                                                                          |
| Buschert et al. (2018)       | E: Endurance training<br>C: Occupational or art therapy                                                                  | F: 2–3 times/wk<br>I: 85 % HRmax (as the upper limit)<br>T: 30 min<br>D: 3–4 wks | <b>Attention:</b> TAP-Simple Reaction Times (TAP-S-RT); TAP-Audio-Visual Conditions (TAP-AVC); Digit span forward (DSF);<br><b>Memory:</b> Verbaler Lern-und Merkfähigkeitstest Learning (VLMT Learning); Verbaler Lern-und Merkfähigkeitstest Recall (VLMT Recall);<br><b>EF:</b> Digits span backward (DSB); Standard progressive matrices (SPM); Computergestütztes Kartensortierverfahren (CKV)                                                                                                                                                                                      |
| Foley et al. (2008)          | E: Aerobic exercise<br>C: Stretching                                                                                     | F: 3 times/wk<br>I: Moderate<br>T: 30–40 min<br>D: 12 wks                        | <b>Memory:</b> Un-Cued Recall (UCR); Cued Recall (CR)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Hoffman et al. (2008)        | E: Aerobic exercise<br>C <sup>1</sup> : Antidepressant<br>C <sup>2</sup> : Placebo pill                                  | F: 3 times/wk<br>I: 70–85 % HRR<br>T: 45 min<br>D: 16 wks                        | <b>Memory:</b> Logical Memory Subtest from the WMS (LMS-WMS); Verbal Paired Associates Subtest from the WMS-easy (VPAS-WMS-easy); Verbal Paired Associates Subtest from the WMS (VPAS-WMS-hard); <b>Attention:</b> Digit Span Forward (DSF); <b>EF:</b> Digits Span Backward (DSB); Animal Naming Test (ANT); Controlled Oral Word Association Test (COWAT); Stroop Color/Word Test-Interference (SCWT-Int); Ruff 2 & 7 Test total (Ruff-total); Digit Symbol Test-Wechsler Adult Intelligence Scale-Revised (DST-WAIS-R); Trail Making Test B-A (TMT B-A)                               |
| Imboden et al. (2020)        | E: Aerobic exercise<br>C: Stretching                                                                                     | F: 3 times/wk<br>I: 60–75 % HRmax<br>T: 45 min<br>D: 6 wks                       | <b>Attention:</b> Testbatterie zur Aufmerksamkeitsprüfung-Alertness, no warning signal (TAP-A-NWS)/with warning signal (TAP-A WWS); <b>EF:</b> Testbatterie zur Aufmerksamkeitsprüfung-Working Memory-Difficulty Level 3 (TAP-WM); Testbatterie zur Aufmerksamkeitsprüfung-Cognitive Flexibility-Verbal Change (TAP-CF); Testbatterie zur Aufmerksamkeitsprüfung-Go/No-Go-Reaction Time (TAP-GNG-RT)                                                                                                                                                                                     |
| Khatiri et al. (2001)        | E: Aerobic exercise<br>C: Antidepressant                                                                                 | F: 3 times/wk<br>I: 70–85 % HRR<br>T: 45 min<br>D: 16 wks                        | <b>Memory:</b> WMS-Logical Memory Immediate Recall (WMS-LMIR); WMS-Logical Memory Delayed Recall (WMS-LMDR); WMS-Logical Memory Visual Reproduction Immediate Recall (WMS-LMVRIR); WMS-Logical Memory Visual Reproduction Delayed Recall (WMS-LMVRDR); <b>Attention:</b> Digit Span Forward (DSF); <b>PS:</b> Trail Making Test A (TMT-A); Stroop Word Test (SWT); Stroop Color Test (SCT); <b>EF:</b> Digit Symbol Test-Wechsler Adult Intelligence Scale-Revised (DST-WAIS-R); Stroop Color/Word Test-Interference (SCWT-Int); Digits Span Backward (DSB); Trail Making Test B (TMT-B) |
| Krogh et al. (2009)          | E: Aerobic exercise<br>C <sup>1</sup> : Resistance training<br>C <sup>2</sup> : Relaxation                               | F: 2 times/wk<br>I: 70–89 % HRmax<br>T: 90 min<br>D: 16 wks                      | <b>Attention:</b> Digit Span (DS); Subtracting Serial Sevens (SSS);<br><b>PS:</b> Trail Making Test A (TMT-A); Design Fluency Test (DFT);<br><b>EF:</b> Digit Symbol Test-Wechsler Adult Intelligence Scale-Revised (DST-WAIS-R); Verbal Fluency Test-Animal (VFT-A); Verbal Fluency Test-S word (VFT-S); Trail Making Test B (TMT-B);<br><b>Memory:</b> Buschke Selective Reminding Test (BSRT); Rey Complex Figure Test (RCFT)                                                                                                                                                         |
| Krogh et al. (2012)          | E: Aerobic exercise<br>C: Stretching                                                                                     | F: 3 times/wk<br>I: 65–80 % HRmax<br>T: 45 min<br>D: 12 wks                      | <b>Memory:</b> Buschke Selective Reminding Test (BSRT); Rey Complex Figure Test (RCFT);<br><b>Attention:</b> Digit span forward (DSF); Subtracting Serial Sevens (SSS);<br><b>PS:</b> Stroop Color/Word Test-congruent-Reaction time (SCWT-Con-RT); Trail Making Test A (TMT-A);<br><b>EF:</b> Stroop Color/Word Test-Incongruent-Reaction time (SCWT-Incon-RT); Digits Span Backward (DSB); Trail Making Test B (TMT-B); Digit Symbol Test-Wechsler Adult Intelligence Scale-Revised (DST-WAIS-R); Verbal Fluency Test-Animal (VFT-A); Verbal Fluency Test-S word (VFT-S)               |
| Krogh et al. (2014)          | E: Aerobic exercise<br>C: Stretching                                                                                     | F: 3 times/wk<br>I: 80 % HRmax<br>T: 45 min<br>D: 12 wks                         | <b>Memory:</b> Buschke Selective Reminding Test (BSRT); Rey Complex Figure Test (RCFT)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Neviani et al. (2017)        | E: Progressive aerobic exercise<br>C <sup>1</sup> : Non-progressive physical activity<br>C <sup>2</sup> : Antidepressant | F: 3 times/wk<br>I: 60–85 % HRmax<br>T: 60 min<br>D: 24 wks                      | <b>Overall:</b> Montreal Cognitive Assessment (MoCA)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Oertel-Knochel et al. (2014) | E: Aerobic exercise<br>C <sup>1</sup> : Relaxation<br>C <sup>2</sup> : Waiting control                                   | F: 3 times/wk<br>I: 60–70 % HRmax<br>T: 45 min<br>D: 4 wks                       | <b>PS:</b> Trail Making Test A (TMT-A);<br><b>EF:</b> Wechsler Memory Scale-Third Edition-Spatial Span/Letter-Number-Span (WMS-III-SS/LNS);<br><b>Memory:</b> Hopkins Verbal Learning Test-Revised (HVLTR); Brief Visuospatial Memory Test-Revised (BVMTR)                                                                                                                                                                                                                                                                                                                               |
| Olson et al. (2017)          | E: Aerobic exercise<br>C: Stretching                                                                                     | F: 3 times/wk<br>I: 40–65 % HRR<br>T: 45 min<br>D: 8 wks                         | <b>PS:</b> Flanker Task-Congruent-Accuracy (FT-Con-Acc); Flanker Task-Congruent-Reaction Time (FT-Con-RT); <b>EF:</b> Flanker Task-Incongruent-Accuracy (FT-Incon-Acc); Flanker Task-Incongruent-Reaction Time (FT-Incon-RT)                                                                                                                                                                                                                                                                                                                                                             |


Note.  $k$  = number of included studies; E = experimental group; C = control group; C<sup>1</sup> = first control group; C<sup>2</sup> = second control group; F = frequency; I = intensity; T = time; D = duration; /wk = per week; % = percentage; HRR = Heart rate reserve; HRmax = maximum heart rate; min = minutes; wks = weeks; PS = processing speed; EF = executive function

= 0.001), and session time > 60 min ( $g = -0.04$ ; 95 % CI = -0.26, 0.19;  $p = 0.734$ ) on cognitive function.

#### Study design and sample characteristics

The results of the test of moderation for study design and sample

characteristics indicated that there were no statistically significant subgroup differences in ES for comparator/control group ( $F = 0.33$ ,  $p = 0.720$ ), proportion of females in the study sample ( $F = 1.17$ ,  $p = 0.314$ ), and mean age ( $F = 0.56$ ,  $p = 0.644$ ); however, there were statistically significant subgroup differences in ES for aerobic exercise plus antidepressant use ( $F = 7.55$ ,  $p = 0.001$ ), and clinical setting ( $F = 8.90$ ,  $p =$



(caption on next page)

**Fig. 2.** Forest plot of aerobic exercise on cognitive function.

*FT-Con-Acc* = Flanker Task-Congruent-Accuracy; *FT-Incon-Acc* = Flanker Task-Incongruent-Accuracy; *DS* = Digit Span; *AE* = aerobic exercise; *vs.* = versus.; *VFT-A* = Verbal Fluency Test-Animal; *VFT-S* = Verbal Fluency Test-S word; *TMT-A* = Trail Making Test A; *SSS* = Subtracting Serial Sevens; *TAP-CF* = Testbatterie zur Aufmerksamkeitsprüfung-Cognitive Flexibility; *TAP-A-NWS* = Testbatterie zur Aufmerksamkeitsprüfung-Alertness, no warning signal; *DFT* = Design Fluency Test; *TAP-A-WWS* = Testbatterie zur Aufmerksamkeitsprüfung-Alertness, with warning signal; *BSRT* = Buschke Selective Reminding Test; *TMT-B* = Trail Making Test B; *DSF* = Digit Span Forward; *RCFT* = Rey Complex Figure Test; *COWAT* = Controlled Oral Word Association Test; *DSB* = Digits Span Backward; *VPAS-WMS-easy* = Verbal Paired Associates Subtest from the WMS-easy; *SCT* = Stroop Color Test; *DST-WAIS-R* = Digit Symbol Test-Wechsler Adult Intelligence Scale-Revised; *ANT* = Animal Naming Test; *SCWT-Int* = Stroop Color/Word Test-Interference; *VPAS-WMS-hard* = Verbal Paired Associates Subtest from the WMS; *FT-Con-RT* = Flanker Task-Congruent-Reaction Time; *FT-Incon-RT* = Flanker Task-Incongruent-Reaction Time; *WMS-LMDR* = WMS-Logical Memory Delayed Recall; *WMS-LMVRIR* = WMS-Logical Memory Visual Reproduction Immediate Recall; *LMS-WMS* = Logical Memory Subtest from the WMS; *WMS-LMVRDR* = WMS-Logical Memory Visual Reproduction Delayed Recall; *SCWT-Incon-RT* = Stroop Color/Word Test-Incongruent-Reaction time; *SWT* = Stroop Word Test; *WMS-LMIR* = WMS-Logical Memory Immediate Recall; *SCWT-Con-RT* = Stroop Color/Word Test-congruent-Reaction time; *TMT B-A* = Trail Making Test B-A; *MoCA* = Montreal Cognitive Assessment; *VLMT Recall* = Verbaler Lern-und Merkfähigkeitstest Recall; *TAP-S-RT* = Testbatterie zur Aufmerksamkeitsprüfung-Simple Reaction Times; *WMS-III-SS/LNS* = Wechsler Memory Scale-Third Edition-Spatial Span/Letter-Number-Span; *UCR* = Un-Cued Recall; *TAP-GNG-RT* = Testbatterie zur Aufmerksamkeitsprüfung-Go/No-Go-Reaction Time; *HVLT-R* = Hopkins Verbal Learning Test-Revised; *TAP-WM* = Testbatterie zur Aufmerksamkeitsprüfung-Working Memory-Difficulty Level 3; *CR* = Cued Recall; *TAP-AVC* = Testbatterie zur Aufmerksamkeitsprüfung-Audio-Visual Conditions; *VLMT Learning* = Verbaler Lern-und Merkfähigkeitstest Learning; *CKV* = Computergestütztes Kartensortierverfahren; *BVMT-R* = Brief Visuospatial Memory Test-Revised; *SPM* = Standard progressive matrices; *RE Model* = random-effects model.

0.004). The findings of aerobic exercise plus antidepressant use indicated that the ES was larger for YES (which represents participants of aerobic exercise group used antidepressant) ( $g = 0.44$ ; 95 % CI = 0.27, 0.61;  $p < 0.001$ ) than ESs for NO (which represents participants of aerobic exercise group did not use antidepressant) ( $g = 0.14$ ; 95 % CI = 0.01, 0.27;  $p = 0.044$ ) and MIXED (which represents some participants of aerobic exercise group used antidepressant) ( $g = 0.05$ ; 95 % CI = -0.06, 0.16;  $p = 0.401$ ) when compared with a comparator/control group. For clinical setting, the results indicated that aerobic exercise produced a larger effect on cognitive function for inpatients ( $g = 0.47$ ; 95 % CI = 0.25, 0.69;  $p < 0.001$ ) than outpatients ( $g = 0.11$ ; 95 % CI = 0.01, 0.21;  $p = 0.042$ ).

Table 4 summarizes the results of all meta-regressions analyses. The results suggested that the influence of aerobic exercise on cognitive function was significantly moderated by the sample size of included studies ( $\beta = -0.003$ ; 95 % CI = -0.006, -0.001;  $p = 0.039$ ), but not significantly moderated by proportion of female ( $\beta = -0.012$ ; 95 % CI = -0.026, 0.001;  $p = 0.069$ ), mean age ( $\beta = 0.005$ ; 95 % CI = -0.004, 0.015;  $p = 0.273$ ), session time of exercise per time ( $\beta = -0.007$ ; 95 % CI = -0.014, 0.001;  $p = 0.096$ ), duration of exercise ( $\beta = -0.014$ ; 95 % CI = -0.038, 0.009;  $p = 0.226$ ), volume of exercise per week ( $\beta = -0.004$ ; 95 % CI = -0.009, 0.001;  $p = 0.112$ ), and total intervention time ( $\beta = -0.001$ ; 95 % CI = -0.001, 0.001;  $p = 0.299$ ).

### Sensitivity analyses

In one randomized trial (Oertel-Knöchel et al., 2014), the residual for one ES (Oertel-Knochel et al. used the brief visuospatial memory test-revised) was more than three standard deviations from the mean. Removing this outlier reduced the overall ES of aerobic exercise on cognitive function by 0.02 ( $g = 0.19$ ; 95 % CI = 0.07, 0.32;  $p = 0.003$ ). In total, thirteen effect sizes had a Cook's distance more than three standard deviations from the mean. These ES's included Brush et al. (2020) FT-Con-Acc; Buschert et al. (2018) SPM and CKV; Imboden et al. (2020) TAP-A-NWS, TAP-A-WWS, TAP-WM, TAP-CF, and TAP-GNG-RT; Oertel-Knochel et al. (2014) HVLT-R, and BVMT-R; Neviani et al. (2017) MoCA; Foley et al. (2008) CR; Krogh et al. (2014) BSRT (Brush et al., 2020; Buschert et al., 2018; Foley et al., 2008; Imboden et al., 2020; Krogh et al., 2014; Neviani et al., 2017; Oertel-Knöchel et al., 2014), see Table 2 for the full names of the task abbreviations. After removing those outliers, the overall ES of aerobic exercise on cognitive function decreased by 0.05 and remained significant ( $g = 0.16$ ; 95 % CI = 0.05, 0.28;  $p = 0.005$ ).

### Publication bias

The funnel plot (see Fig. 3) showed asymmetry across ES, which was

confirmed by the multilevel extension of Egger's test ( $F = 13.36$ ;  $p = 0.0004$ ). The findings suggested a possibility of publication bias in the included studies.

### Discussion

This meta-analytic review investigated evidence strictly from randomized controlled trials on the effects of aerobic exercise on cognitive function in adults with MDD. Twelve randomized trials were included, and the key results revealed that aerobic exercise significantly improved overall cognitive function, and its sub-domains of memory and executive function in adults with MDD when compared with a comparator/control group. In addition, the effects of aerobic exercise on overall cognitive function were moderated by several factors including session time, aerobic exercise plus antidepressant use, clinical setting, and sample size. These results may contribute to the development of non-pharmacological treatments to enhance cognitive function in adults with MDD.

### Overall cognitive function

Our findings suggested a small to moderate effect on overall cognitive function in adults with MDD following aerobic exercise. However, this result was not consistent with prior meta-analyses (Brondino et al., 2017; Sun et al., 2018), which find nonsignificant effects of physical exercise on cognitive function in depressed adults. Such disparity is not unexpected as the current meta-analysis deviated from previous ones in several meaningful ways. For example, our review focused on aerobic exercise, while the two previous reviews included aerobic, non-aerobic, and mind-body exercises, underscoring potential disparate outcomes when contrasting aerobic exercise with alternative modes of physical activity. As such, our review is the first to confirm that aerobic exercise is effective for improving cognitive function in adults with MDD. Researchers, clinicians, and health professionals can contemplate aerobic exercise as a viable alternative or adjunctive therapy to augment cognitive function in individuals with MDD. Additionally, our finding that aerobic exercise in conjunction with anti-depressant therapy makes it especially apparent that aerobic exercise may not only be complementary to standard therapies but raises the possibility that it may also be synergistic. Future research will need to unpack that question. Interesting, our finding is also consonant with previous meta-analyses that found aerobic exercise had significant improvements on cognitive function in other mental health disorders such as schizophrenia (Shimada et al., 2022) and dementia (Balbim et al., 2022), as well as in healthy individuals including adults older than 50 years (Northey et al., 2018).

**Table 3**  
Categorical moderator analyses of aerobic exercise on cognition.

| Subgroup analyses               | k  | g (95 % CI)              | p-value           | Test of Moderation                 |
|---------------------------------|----|--------------------------|-------------------|------------------------------------|
| <b>Cognitive sub-domains</b>    | 11 |                          |                   | $F(3, 95) = 1.24$ ,<br>$p = 0.298$ |
| Memory                          | 8  | <b>0.25 (0.06, 0.44)</b> | <b>0.012</b>      |                                    |
| Executive Function              | 9  | <b>0.12 (0.04, 0.20)</b> | <b>0.005</b>      |                                    |
| Attention                       | 6  | 0.01 (-0.12, 0.13)       | 0.928             |                                    |
| Processing speed                | 6  | 0.08 (-0.11, 0.26)       | 0.387             |                                    |
| <b>Exercise intensity</b>       | 12 |                          |                   | $F(1, 99) = 0.11$ ,<br>$p = 0.735$ |
| Moderate only                   | 5  | 0.24 (-0.01, 0.49)       | 0.057             |                                    |
| Moderate-to-vigorous (mixed)    | 7  | <b>0.19 (0.02, 0.37)</b> | <b>0.032</b>      |                                    |
| <b>Exercise frequency</b>       | 12 |                          |                   | $F(1, 99) = 1.74$ ,<br>$p = 0.190$ |
| 2 times per week                | 1  | -0.04 (-0.43, 0.35)      | 0.844             |                                    |
| 3 times per week                | 11 | <b>0.23 (0.10, 0.38)</b> | <b>0.001</b>      |                                    |
| <b>Session time</b>             | 12 |                          |                   | $F(2, 98) = 5.36$ ,<br>$p = 0.006$ |
| < 45 min                        | 2  | <b>0.59 (0.28, 0.90)</b> | <b>&lt; 0.001</b> |                                    |
| 45-60 min                       | 9  | <b>0.16 (0.07, 0.26)</b> | <b>0.001</b>      |                                    |
| > 60 min                        | 1  | -0.04 (-0.26, 0.19)      | 0.734             |                                    |
| <b>Exercise duration</b>        | 12 |                          |                   | $F(1, 99) = 0.83$ ,<br>$p = 0.364$ |
| ≤ 12 weeks                      | 8  | <b>0.26 (0.08, 0.44)</b> | <b>0.005</b>      |                                    |
| > 12 weeks                      | 4  | 0.13 (-0.09, 0.35)       | 0.234             |                                    |
| <b>AE plus antidepressant</b>   | 12 |                          |                   | $F(2, 98) = 7.55$ ,<br>$p = 0.001$ |
| No                              | 2  | <b>0.14 (0.01, 0.27)</b> | <b>0.044</b>      |                                    |
| Yes                             | 4  | <b>0.44 (0.27, 0.61)</b> | <b>&lt; 0.001</b> |                                    |
| Mixed                           | 6  | 0.05 (-0.06, 0.16)       | 0.401             |                                    |
| <b>Comparator/control group</b> | 12 |                          |                   | $F(2, 98) = 0.33$ ,<br>$p = 0.720$ |
| Active                          | 10 | <b>0.18 (0.02, 0.35)</b> | <b>0.030</b>      |                                    |
| Placebo pill                    | 1  | 0.24 (-0.03, 0.52)       | 0.078             |                                    |
| Treatment as usual care         | 4  | <b>0.28 (0.04, 0.53)</b> | <b>0.026</b>      |                                    |
| <b>Proportion of female</b>     | 12 |                          |                   | $F(2, 98) = 1.17$ ,<br>$p = 0.314$ |
| > 50 %                          | 9  | <b>0.16 (0.02, 0.30)</b> | <b>0.031</b>      |                                    |
| ≤ 50 %                          | 2  | <b>0.41 (0.06, 0.76)</b> | <b>0.023</b>      |                                    |
| Unclear                         | 1  | 0.56 (-0.33, 1.46)       | 0.217             |                                    |
| <b>Mean age</b>                 | 12 |                          |                   | $F(3, 97) = 0.56$ ,<br>$p = 0.644$ |
| Young adults (18-44 yrs)        | 6  | 0.13 (-0.06, 0.33)       | 0.173             |                                    |
| Middle adults (45-64 yrs)       | 4  | <b>0.25 (0.02, 0.48)</b> | <b>0.030</b>      |                                    |
| Old adults (≥ 65 yrs)           | 1  | 0.37 (-0.15, 0.89)       | 0.161             |                                    |
| Unclear                         | 1  | 0.56 (-0.35, 1.48)       | 0.224             |                                    |
| <b>Clinical setting</b>         | 12 |                          |                   | $F(1, 99) = 8.90$ ,<br>$p = 0.004$ |
| Inpatients                      | 3  | <b>0.47 (0.25, 0.69)</b> | <b>&lt; 0.001</b> |                                    |
| Outpatients                     | 9  | <b>0.11 (0.01, 0.21)</b> | <b>0.042</b>      |                                    |

Note. k = number of included studies; g = Hedges' g; CI = confidence interval, AE = aerobic exercise; yrs = years

**Table 4**  
Continuous moderator analyses of aerobic exercise on cognition.

| Meta-regression             | k  | $\beta$ (95 % CI)             | p-value      | Test of Moderation |
|-----------------------------|----|-------------------------------|--------------|--------------------|
| Proportion of female        | 11 | -0.012 (-0.026, 0.001)        | 0.069        | $F(1, 97) = 3.39$  |
| Mean age                    | 11 | 0.005 (-0.004, 0.015)         | 0.273        | $F(1, 97) = 1.22$  |
| Session time of exercise/T  | 12 | -0.007 (-0.014, 0.001)        | 0.096        | $F(1, 99) = 2.83$  |
| Duration of exercise        | 12 | -0.014 (-0.038, 0.009)        | 0.226        | $F(1, 99) = 1.49$  |
| Volume of exercise per week | 12 | -0.004 (-0.009, 0.001)        | 0.112        | $F(1, 99) = 2.57$  |
| Total intervention time     | 12 | -0.001 (-0.001, 0.001)        | 0.299        | $F(1, 99) = 1.09$  |
| Sample size                 | 12 | <b>-0.003(-0.006, -0.001)</b> | <b>0.039</b> | $F(1, 99) = 4.36$  |

Note. k = number of included studies;  $\beta$  = regression coefficient; CI = confidence interval; /T = per time.

#### Cognitive sub-domains

Our meta-analysis yielded a small-to-moderate ES for memory, which is in line with a previous meta-analysis (Brondino et al., 2017) that found physical exercise had a significant effect on visual learning and memory in adults with depression. However, our finding contradicts another prior meta-analysis (Sun et al., 2018), which found physical exercise had no significant benefit on verbal/visual learning and memory in individuals with depression. These conflicting results may be due to differences in inclusion criteria (i.e., severity of depression/depression symptoms), as well as the type of exercise intervention, as the previous meta-analysis included aerobic, resistance, and mind-body exercise, and our meta-analysis specifically focused on aerobic exercise. Further, the previous meta-analysis analyzed verbal learning, visual learning, and memory separately. However, we performed an overall analysis of memory, and found that aerobic exercise had a small-to-moderate ES in adults with MDD.

Regarding executive function, the present findings confirmed the results from a recent meta-analysis (Ren, Alderman, et al., 2023), which observed that aerobic exercise significantly enhanced executive function in depressed adults. Notably, the ES in the present meta-analysis was smaller than that reported by (Ren, Alderman, et al., 2023). Inclusion criteria and the number of included studies may have contributed to the discrepancy in results. The previous meta-analysis (Ren, Alderman, et al., 2023) included both English and Chinese language published studies ( $k = 10$ ); however, the current study only included English language published studies ( $k = 9$ ). Importantly, these results support aerobic exercise as a promising intervention that has positive benefits for executive function in adults with MDD.

Unfortunately, no significant effects were observed for attention or processing speed in this review. Prior meta-analyses (Brondino et al., 2017; Sun et al., 2018) indicated that physical exercise had no significant effect on attention and processing speed in people with depression, similar to our results. In addition, our results are in agreement with a previous meta-analysis, which found that aerobic exercise had non-significant effects on attention and processing speed in participants with schizophrenia (Xu et al., 2022). Furthermore, a non-significant ES on processing speed in participants with ischemic cerebrovascular disorder after aerobic exercise has been observed (Shu et al., 2020). However, it should be noted that Smith et al. (2010) has found positive benefits for attention and processing speed in individuals without depression after aerobic exercise (Smith et al., 2010). Since only six randomized trials have investigated the effects of aerobic exercise on attention and processing speed in adults with MDD. Therefore, more studies are needed in the future to investigate the effects of aerobic exercise on these cognitive sub-domains in adults with MDD.




Fig. 3. Funnel plot of potential publication bias.

#### Aerobic exercise characteristics

Exercise intervention characteristics, including intensity, frequency, session time, and duration, play significant roles in the cognitive benefits of aerobic exercise (Shimada et al., 2022). Our findings suggest that the optimal dose for the effects of aerobic exercise on cognitive function in adults with MDD are as follows: moderate-to-vigorous (mixed), frequency of 3 times per week, each session time < 45 min, and intervention duration of  $\leq 12$  weeks. However, these findings are from a small number of studies, and future work is needed to better understand how to optimize the exercise 'dose' to best improve MDD symptoms.

Regarding intensity, our results showed that moderate-to-vigorous (mixed) aerobic exercise induced a significant small-to-moderate ES on cognitive function in adults with MDD. However, moderate intensity aerobic exercise produced a favorable but non-significant ES on cognitive function. These findings are meaningful, since both moderate intensity and moderate-to-vigorous intensity exercise were analyzed as a whole in previous meta-analyses (Brondino et al., 2017; Ren, Alderman, et al., 2023; Sun et al., 2018). Accordingly, we separated them into two groups for analysis, and these results may provide a fresh perspective for formulating the aerobic exercise prescription. In addition, further investigation into moderate intensity aerobic exercise using well-designed randomized controlled trials is crucial to understanding the relationship of aerobic exercise as a treatment for adults with MDD.

As for frequency, session time, and duration, our meta-analysis found that aerobic exercise performed 3 times per week, for session time < 45 min and between 45–60 min, and for a duration of  $\leq 12$  weeks had significant benefits on cognitive function; however, aerobic exercise performed 2 times per week, for session time > 60 minute, and for a duration of > 12 weeks did not. Our results of frequency and session time align with a recent meta-analysis (Ren, Alderman, et al., 2023), which found that exercise training (resistance, mind-body and aerobic) conducted 3 times per week, with a session time  $\leq 60$  min resulted in significant effects on executive function in depressed adults. However, our results for duration contradict the results of the previous meta-analysis (Ren, Alderman, et al., 2023), which indicated that exercise training practiced with a duration of  $\geq 13$  weeks significantly promoted executive function in depressed adults, but 3–12 weeks did not. What cannot ignore the non-significant result of 2 times per week ( $k = 1$ ), session time > 60 minute ( $k = 1$ ), and durations > 12 weeks ( $k = 4$ )

are clearly due to a lack of statistical power. Therefore, further research is needed to determine if aerobic exercise at less frequent, longer session time and duration can also lead to significant improvements in cognitive function in adults with MDD.

#### Study design and sample characteristics

The present meta-analysis explored two study design characteristics (aerobic exercise plus antidepressant use, and comparator/control group), and three sample characteristics (proportion of females, mean age, and clinical setting) as moderators. The test of moderation indicated that the effect of aerobic exercise on cognitive function was moderated by antidepressant use as well as clinical setting in adults with MDD. Regarding antidepressant use, a small beneficial ES on cognitive function was found when the aerobic exercise group did not use antidepressant medication (NO), although the ES just reached the threshold for statistical significance, this suggests that aerobic exercise intervention may have a positive impact on cognitive function in adults with MDD. In addition, a small-to-moderate benefit on cognitive function was observed when the aerobic exercise group used antidepressant medication (YES). This pattern of results suggests that aerobic exercise combined with antidepressant medication is a more effective intervention for improving cognitive function in adults with MDD. The combination of both interventions may produce a synergistic effect, yielding a better treatment effect. The specific mechanisms behind the observed synergistic effect of antidepressants and aerobic exercise remain unclear; nonetheless, existing evidence supports the idea that both interventions contribute to the promotion of neuroplasticity (Andrade & Rao, 2010; Mellow et al., 2020). The integration of these two interventions might be the reason underlying a greater cognitive effect via the facilitation of neuroplasticity. However, a significant ES did not extend to the aerobic exercise group when only some of the participants used antidepressants (MIXED). The lack of statistical significance in the ES may be due to the use of different comparison/control groups, study design, exercise dosage, and sample size. Considering that only 8.5 % to 64.6 % of participants used aerobic exercise combined with antidepressant across 6 studies, the result requires further investigation, as our results are likely underpowered. For the comparator/control group, the findings suggest that aerobic exercise produced significant effects on cognitive function when compared to an active group or a treatment as

usual group, but did not produce significant effects on cognitive function when compared with placebo medication. Our results are partially consistent with the findings of (Ren, Alderman, et al., 2023), which found that exercise had significant effects on executive function when compared with usual medical treatment and placebo medication, but had no significant effects on executive function when compared to an active control. However, our results suggest aerobic exercise may have more beneficial effects on cognition than active comparator/control groups (stretching, occupational or art therapy, non-progressive physical activity, relaxation, resistance training), and treatment as usual groups (antidepressant).

Results from included trials with more than 50 % and less than or equal to 50 % female participants showed that aerobic exercise significantly enhanced cognitive function, implying that aerobic exercise is an effective intervention to improve cognition regardless of sex. For age, the results indicate that aerobic exercise only produces significant benefits on cognition in middle-aged depressed adults (45–64 years), but not in younger depressed (18–44 years old) and older depressed adults ( $\geq 65$  years old). Our findings were partly comparable to a previous review (Ren, Alderman, et al., 2023). However, it is worth noting that there is only one study that examined older adults with MDD. As such, more studies are required for older participants to properly power this question. Lastly, although both groups demonstrated a significant effect, we found preliminary evidence for greater improvements in cognitive function among inpatient relative to outpatient participants following aerobic exercise.

Aerobic exercise may enhance cognitive function in adults with MDD via different underlying mechanisms. First, aerobic exercise has been shown to increase cerebral blood flow (Kleinloog et al., 2019; Tomoto et al., 2021), which plays a critical role in maintaining proper brain function and cognition (Abdelsaid et al., 2016; Cheng et al., 2022). Second, brain-derived neurotrophic factor (BDNF), a protein that plays an important role in the development and survival of nerve cells (Nowak et al., 2015), is closely associated with cognitive functions (White & Castellano, 2008). Decreased BDNF levels in the brain are associated with impaired cognitive function (Canivet et al., 2017). Depressed participants have evidenced lower serum BDNF levels when compared with healthy controls, and have been found to negatively correlate with depression severity (Chiou & Huang, 2017). Aerobic exercise has been shown to increase the production of BDNF (Kim et al., 2022; Salehi et al., 2016), which support the growth and survival of neurons and may enhance cognitive function. In addition, a recent meta-analysis (da Cunha et al., 2023) suggested that exercise interventions (including aerobic exercise) led to a significantly increase in circulating BDNF in adults with MDD when compared with a control group. Lastly, the hippocampus is a complicated region inside the brain that serves as a pivotal component in the process of memory formation and retrieval (Xiu et al., 2023), and it appears to be highly plastic to intervention (Wilckens et al., 2021). Hippocampal atrophy has been associated with cognitive impairments (De Crescenzo et al., 2016), and decreased hippocampal volume in individuals with depression has been documented (Campbell et al., 2004; Sheline et al., 1996). Previous studies have shown that exercise interventions significantly increase hippocampal volume in typically aging humans (Erickson et al., 2011; Ten Brinke et al., 2015; Wilckens et al., 2021). Therefore, increased hippocampal volume after aerobic exercise may enhance cognition in individuals with MDD. Overall, the mechanisms by which aerobic exercise improves cognitive function in adults with MDD are complex. More studies are needed to examine the potential mechanisms that underlie aerobic exercise effects on cognitive function in depressed individuals.

## Limitations

There are several limitations to the present systematic review and meta-analysis. First, the current review only included those studies published in peer-reviewed English-language journals, which may lead

to a language bias. Second, the results of the Egger's test suggested a potential publication bias, despite the fact that we used a rigorous selection process and criteria to determine which studies we included; therefore, it is still necessary to interpret the findings of the current meta-analysis with caution, especially given the small number of studies that exist in this field of investigation, which further led to multiple instances of underpowered results. Lastly, only two of the 12 included trials underwent follow-up evaluation on cognitive function in adults with MDD. As a consequence, we did not conduct follow-up analysis, and further randomized trials are needed to determine how long the positive effects of aerobic exercise on cognition may last following intervention.

## Conclusions

The present systematic review and meta-analysis concludes that aerobic exercise is an effective approach to improve cognitive function, as well as the sub-domains of memory and executive function, in adults with MDD. Aerobic exercise may be a promising non-pharmacological treatment for cognitive function that supplements standard, pharmacological treatments in this population. More studies are needed to examine the extent and dosage of exercise intervention, as well as the potential mechanisms that underlie the benefits of aerobic exercise on cognitive function in adults with MDD.

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.ijchp.2024.100447](https://doi.org/10.1016/j.ijchp.2024.100447).

## References

- Abdelsaid, M., Williams, R., Hardigan, T., & Ergul, A. (2016). Linagliptin attenuates diabetes-induced cerebral pathological neovascularization in a blood glucose-independent manner: Potential role of ET-1. *Life Sciences*, 159, 83–89. <https://doi.org/10.1016/j.lfs.2015.11.026>
- Andrade, C., & Rao, N. S. (2010). How antidepressant drugs act: A primer on neuroplasticity as the eventual mediator of antidepressant efficacy. *Indian Journal of Psychiatry*, 52(4), 378–386. <https://doi.org/10.4103/0019-5545.74318>
- Balbin, G. M., Falck, R. S., Barha, C. K., Starkey, S. Y., Bullock, A., Davis, J. C., & Liu-Ambrose, T. (2022). Effects of exercise training on the cognitive function of older adults with different types of dementia: A systematic review and meta-analysis. *British Journal of Sports Medicine*, 56(16), 933–940. <https://doi.org/10.1136/bjsports-2021-104955>
- Birnie, M. T., Eapen, A. V., Kershaw, Y. M., Lodge, D., Collingridge, G. L., Conway-Campbell, B. L., & Lightman, S. L. (2022). Time of day influences stress hormone response to ketamine. *Journal of Neuroendocrinology*, 34(10), e13194. <https://doi.org/10.1111/jne.13194>
- Brondino, N., Rocchetti, M., Fusar-Poli, L., Codrons, E., Correale, L., Vandoni, M., Barbui, C., & Politi, P. (2017). A systematic review of cognitive effects of exercise in depression. *Acta Psychiatrica Scandinavica*, 135(4), 285–295. <https://doi.org/10.1111/acps.12690>
- Brush, C. J., Hajcak, G., Bocchine, A. J., Ude, A. A., Muniz, K. M., Foti, D., & Alderman, B. L. (2020). A randomized trial of aerobic exercise for major depression: Examining neural indicators of reward and cognitive control as predictors and treatment targets. *Psychological Medicine*, 52(5), 893–903. <https://doi.org/10.1017/S0033291720002573>
- Buschert, V., Prochazka, D., Bartl, H., Diemer, J., Malchow, B., Zwanzger, P., & Brunner, A. (2018). Effects of physical activity on cognitive performance: A controlled clinical study in depressive patients. *European Archives of Psychiatry and Clinical Neuroscience*, 269(5), 555–563. <https://doi.org/10.1007/s00406-018-0916-0>
- Campbell, S., Marriott, M., Nahmias, C., & MacQueen, G. M. (2004). Lower hippocampal volume in patients suffering from depression: A meta-analysis. *American Journal of Psychiatry*, 161(4), 598–607. <https://doi.org/10.1176/appi.ajp.161.4.598>
- Canivet, A., Albinet, C. T., Rodríguez-Ballesteros, M., Chicherio, C., Fagot, D., André, N., & Audiffren, M. (2017). Interaction between BDNF polymorphism and physical activity on inhibitory performance in the elderly without cognitive impairment.

- Frontiers in Human Neuroscience, 11(541). <https://doi.org/10.3389/fnhum.2017.00541>
- Chai, H., Liu, B., Zhan, H., Li, X., He, Z., Ye, J., Guo, Q., Chen, J., Zhang, J., & Li, S. (2019). Antidepressant effects of rhodomyrtone in mice with chronic unpredictable mild stress-induced depression. *International Journal of Neuropsychopharmacology*, 22(2), 157–164. <https://doi.org/10.1093/Fijnp/Fpyy091>
- Chen, F. T., Etnier, J. L., Chan, K. H., Chiu, P. K., Hung, T. M., & Chang, Y. K. (2020). Effects of exercise training interventions on executive function in older adults: A systematic review and meta-analysis. *Sports Medicine*, 50(8), 1451–1467. <https://doi.org/10.1007/s40279-020-01292-x>
- Cheng, X., Chen, H., Sie, E. J., Marsili, F., & Boas, D. A. (2022). Development of a Monte Carlo-wave model to simulate time domain diffuse correlation spectroscopy measurements from first principles. *Journal of Biomedical Optics*, (8), 27. <https://doi.org/10.1117/1.Jbo.27.8.083009>
- Cheung, M. W. (2014). Modeling dependent effect sizes with three-level meta-analyses: A structural equation modeling approach. *Psychological Methods*, 19(2), 211–229.
- Chiou, Y. J., & Huang, T. L. (2017). Serum brain-derived neurotrophic factors in Taiwanese patients with drug-naïve first-episode major depressive disorder: Effects of antidepressants. *International Journal of Neuropsychopharmacology*, 20(3), 213–218. <https://doi.org/10.1093/ijnp/pyw096>
- Cohen, J. (1992). A power primer. *Psychological Bulletin*, 112(1), 155–159. <https://doi.org/10.1037//0033-2909.112.1.155>
- Contreras-Osorio, F., Ramirez-Campillo, R., Cerda-Vega, E., Campos-Jara, R., Martínez-Salazar, C., Reigal, R. E., Hernández-Mendo, A., Carneiro, L., & Campos-Jara, C. (2022). Effects of physical exercise on executive function in adults with depression: A systematic review and meta-analysis. *International Journal of Environmental Research and Public Health*, 19(22), 15270. <https://doi.org/10.3390/ijerph192215270>
- da Cunha, L. L., Feter, N., Alt, R., & Rombaldi, A. J. (2023). Effects of exercise training on inflammatory, neurotrophic and immunological markers and neurotransmitters in people with depression: A systematic review and meta-analysis. *Journal of Affective Disorders*, 326, 73–82. <https://doi.org/10.1016/j.jad.2023.01.086>
- De Crescenzo, F., Foti, F., Ciabattini, M., Del Giovane, C., Watanabe, N., Sañé Schepisi, M., Quested, D. J., Cipriani, A., Barbui, C., & Amato, L. (2016). Comparative efficacy and acceptability of pharmacological treatments for insomnia in adults: A systematic review and network meta-analysis. *Cochrane Database of Systematic Reviews*, 2016(9). <https://doi.org/10.1002/14651858.CD012364>
- Du, Y., Ruan, J., Zhang, L., & Fu, F. (2020). Jieyu Anshen Granule, a Chinese herbal formulation, exerts effects on poststroke depression in rats. *Evidence-Based Complementary and Alternative Medicine*, 7469068. <https://doi.org/10.1155/2020/7469068>
- Egger, M., Davey Smith, G., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *BMJ*, 315(7109), 629–634. <https://doi.org/10.1136/bmj.315.7109.629>
- Erickson, K. I., Voss, M. W., Prakash, R. S., Basak, C., Szabo, A., Chaddock, L., Kim, J. S., Heo, S., Alves, H., White, S. M., Wojcicki, T. R., Mailey, E., Vieira, V. J., Martin, S. A., Pence, B. D., Woods, J. A., McAuley, E., & Kramer, A. F. (2011). Exercise training increases size of hippocampus and improves memory. *Proceedings of the National Academy of Sciences*, 108(7), 3017–3022. <https://doi.org/10.1073/pnas.1015950108>
- Foley, L. S., Prapaavessis, H., Osuch, E. A., De Pace, J. A., Murphy, B. A., & Podolinsky, N. J. (2008). An examination of potential mechanisms for exercise as a treatment for depression: A pilot study. *Mental Health and Physical Activity*, 1(2), 69–73. <https://doi.org/10.1016/j.mhpa.2008.07.001>
- Greenberg, P. E., Fournier, A. A., Sisitsky, T., Simes, M., Berman, R., Koenigsberg, S. H., & Kessler, R. C. (2021). The economic burden of adults with major depressive disorder in the United States (2010 and 2018). *Pharmacoeconomics*, 39(6), 653–665.
- Hammar, Å., Ronold, E. H., & Rekkedal, G. Å. (2022). Cognitive impairment and neurocognitive profiles in major depression—a clinical perspective. *Frontiers in Psychiatry*, 13, Article 764374. <https://doi.org/10.3389/fpsy.2022.764374>
- Heissel, A., Heinen, D., Brokmeier, L. L., Skarabis, N., Kangas, M., Vancampfort, D., Stubbs, B., Firth, J., Ward, P. B., Rosenbaum, S., Hallgren, M., & Schuch, F. (2023). Exercise as medicine for depressive symptoms? A systematic review and meta-analysis with meta-regression. *British Journal of Sports Medicine*. <https://doi.org/10.1136/bjsports-2022-106282>
- Higgins, J. P., Thompson, S. G., Deeks, J. J., & Altman, D. G. (2003). Measuring inconsistency in meta-analyses. *BMJ*, 327(7414), 557–560. <https://doi.org/10.1136/bmj.327.7414.557>
- Higgins, J. P. T., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M. J., & Welch, V. A. (2019). *Cochrane handbook for systematic reviews of interventions*. John Wiley & Sons.
- Hoffman, B. M., Blumenthal, J. A., Babyak, M. A., Smith, P. J., Rogers, S. D., Doraiswamy, P. M., & Sherwood, A. (2008). Exercise fails to improve neurocognition in depressed middle-aged and older adults. *Medicine and Science in Sports and Exercise*, 40(7), 1344–1352. <https://doi.org/10.1249/MSS.0b013e31816b877c>
- Iancu, S. C., Wong, Y. M., Rhebergen, D., van Balkom, A. J. L. M., & Batelaan, N. M. (2020). Long-term disability in major depressive disorder: A 6-year follow-up study. *Psychological Medicine*, 50(10), 1644–1652. <https://doi.org/10.1017/S0033291719001612>
- Imboden, C., Gerber, M., Beck, J., Holsboer-Trachsler, E., Pühse, U., & Hatzinger, M. (2020). Aerobic exercise or stretching as add-on to inpatient treatment of depression: Similar antidepressant effects on depressive symptoms and larger effects on working memory for aerobic exercise alone. *Journal of Affective Disorders*, 276, 866–876. <https://doi.org/10.1016/j.jad.2020.07.052>
- Irwin, C. L., Coelho, P. S., Kluwe-Schiavon, B., Silva-Fernandes, A., Gonçalves, Ó, F., Leite, J., & Carvalho, S. (2023). Non-pharmacological treatment-related changes of molecular biomarkers in major depressive disorder: A systematic review and meta-analysis. *International Journal of Clinical and Health Psychology*, 23(3), Article 100367. <https://doi.org/10.1016/j.ijchp.2023.100367>
- Kawilapat, S., Maneeton, B., Maneeton, N., Prasitwattanaseree, S., Kongsuk, T., Arunpongpaissal, S., Leejongpermpoon, J., Sukhawaha, S., & Traisathit, P. (2022). Comparison of unweighted and item response theory-based weighted sum scoring for the nine-questions depression-rating scale in the Northern Thai Dialect. *BMC Medical Research Methodology*, 22(1), 268. <https://doi.org/10.1186/s12874-022-01744-0>
- Keefe, R. S. E., Cañadas, E., Farlow, D., & Etkin, A. (2022). Digital intervention for cognitive deficits in major depression: A randomized controlled trial to assess efficacy and safety in adults. *American Journal of Psychiatry*, 179(7), 482–489. <https://doi.org/10.1176/appi.ajp.21020125>
- Khatri, P., Blumenthal, J. A., Babyak, M. A., Craighead, W. E., Herman, S., Baldewicz, T., Madden, D. J., Doraiswamy, M., Waugh, R., & Krishnan, K. R. (2001). Effects of exercise training on cognitive functioning among depressed older men and women. *Journal of Aging and Physical Activity*, 9, 43–57. <https://doi.org/10.1123/japa.9.1.43>
- Kim, H. J., Lee, D., & Lee, Y. (2022). The effect of aerobic exercise on brain-derived neurotrophic factor (BDNF) in individuals with mild cognitive impairment: A systematic review and meta-analysis of a randomized controlled trials. *Physical Therapy Rehabilitation Science*, 11(3), 304–310. <https://doi.org/10.1016/j.neubiorev.2021.05.025>
- Kleinlog, F. P. D., Mensink, R. P., Ivanov, D., Adam, J. J., Uludağ, K., & Joris, P. J. (2019). Aerobic exercise training improves cerebral blood flow and executive function: A randomized, controlled cross-over trial in sedentary older men. *Frontiers in Aging Neuroscience*, 11(333). <https://doi.org/10.3389/fnagi.2019.00333>
- Kowalec, K., Salter, A., Fitzgerald, K. C., Patel, M., Han, J., Lu, Y., Bolton, J. M., Hitchon, C., Bernstein, C. N., & Patten, S. (2022). Depressive symptom trajectories and polygenic risk scores in individuals with an immune-mediated inflammatory disease. *General Hospital Psychiatry*, 77, 21–28. <https://doi.org/10.1016/j.genhosppsych.2022.04.005>
- Kriesche, D., Woll, C. F. J., Tschentscher, N., Engel, R. R., & Karch, S. (2022). Neurocognitive deficits in depression: A systematic review of cognitive impairment in the acute and remitted state. *European Archives of Psychiatry and Clinical Neuroscience*, 273(5), 1105–1128. <https://doi.org/10.1007/s00406-022-01479-5>
- Krogh, J., Hjorthøj, C., Speyer, H., Gluud, C., & Nordentoft, M. (2017). Exercise for patients with major depression: A systematic review with meta-analysis and trial sequential analysis. *BMJ Open*, 7(9), Article e014820. <https://doi.org/10.1136/bmjopen-2016-014820>
- Krogh, J., Rostrup, E., Thomsen, C., Elfving, B., Videbech, P., & Nordentoft, M. (2014). The effect of exercise on hippocampal volume and neurotrophins in patients with major depression—a randomized clinical trial. *Journal of Affective Disorders*, 165, 24–30. <https://doi.org/10.1016/j.jad.2014.04.041>
- Krogh, J., Saltin, B., Gluud, C., & Nordentoft, M. (2009). The DEMO trial: A randomized, parallel-group, observer-blinded clinical trial of strength versus aerobic versus relaxation training for patients with mild to moderate depression. *Journal of Clinical Psychiatry*, 70(6), 790–800. <https://doi.org/10.4088/jcp.08m04241>
- Krogh, J., Videbech, P., Thomsen, C., Gluud, C., & Nordentoft, M. (2012). Aerobic exercise versus stretching exercise in patients with major depression—a randomised clinical trial. *PLoS One*, 7(10), e48316. <https://doi.org/10.1371/journal.pone.0048316>
- Lan, X. F., Wu, F. C., Wang, C. Y., Wu, K., Fang, Z. Y., Lao, G. H., Zhang, B., Ning, Y. P., & Zhou, Y. L. (2023). Sex differences in the association of plasma cytokines and neurocognition in first-episode major depressive disorder. *Journal of Affective Disorders*, 322, 258–266. <https://doi.org/10.1016/j.jad.2022.11.016>
- Lee, J., Gierc, M., Vila-Rodriguez, F., Puterman, E., & Faulkner, G. (2021). Efficacy of exercise combined with standard treatment for depression compared to standard treatment alone: A systematic review and meta-analysis of randomized controlled trials. *Journal of Affective Disorders*, 295, 1494–1511. <https://doi.org/10.1016/j.jad.2021.09.043>
- Lee, Y. J., Kim, H. R., Lee, C. Y., Hyun, S. A., Ko, M. Y., Lee, B. S., Hwang, D. Y., & Ka, M. (2020). 2-Phenylethylamine (PEA) ameliorates corticosterone-induced depression-like phenotype via the BDNF/TrkB/CREB signaling pathway. *International Journal of Molecular Sciences*, 21(23), 9103. <https://doi.org/10.3390/ijms21239103>
- Legemaat, A. M., Semkovska, M., Brouwer, M., Geurtsen, G. J., Burger, H., Denys, D., & Bockting, C. L. (2022). Effectiveness of cognitive remediation in depression: A meta-analysis. *Psychological Medicine*, 52(16), 4146–4161. <https://doi.org/10.1017/S0033291721001100>
- Lépine, J. P., & Briley, M. (2011). The increasing burden of depression. *Neuropsychiatric Disease and Treatment*, 7, 3–7. <https://doi.org/10.2147/NDT.S19617> Suppl. 1.
- Li, X., Huang, Y., Liu, M., Zhang, M., Liu, Y., Teng, T., Liu, X., Yu, Y., Jiang, Y., Ouyang, X., Xu, M., Lv, F., Long, Y., & Zhou, X. (2023). Childhood trauma is linked to abnormal static-dynamic brain topology in adolescents with major depressive disorder. *International Journal of Clinical and Health Psychology*, 23(4), Article 100401. <https://doi.org/10.1016/j.ijchp.2023.100401>
- Maier, C. G., Sherrington, C., Herbert, R. D., Moseley, A. M., & Elkins, M. (2003). Reliability of the PEDro scale for rating quality of randomized controlled trials. *Physical Therapy*, 83(8), 713–721.
- Maramis, M. M., Mahajudin, M. S., & Khotib, J. (2021). Impaired cognitive flexibility and working memory precedes depression: A rat model to study depression. *Neuropsychobiology*, 80(3), 225–233. <https://doi.org/10.1159/000508682>
- McSweeney, M. P., Coombes, J. S., MacKay, C. P., Rodriguez, A. D., Erickson, K. I., Copland, D. A., & McMahon, K. L. (2019). The immediate effects of acute aerobic

- exercise on cognition in healthy older adults: A systematic review. *Sports Medicine*, 49(1), 67–82. <https://doi.org/10.1007/s40279-018-01039-9>
- Mellow, M. L., Goldsworthy, M. R., Coussens, S., & Smith, A. E. (2020). Acute aerobic exercise and neuroplasticity of the motor cortex: A systematic review. *Journal of Science and Medicine in Sport*, 23(4), 408–414. <https://doi.org/10.1016/j.jsams.2019.10.015>
- Nanda, B., Balde, J., & Manjunatha, S. (2013). The acute effects of a single bout of moderate-intensity aerobic exercise on cognitive functions in healthy adult males. *Journal of Clinical and Diagnostic Research*, 7(9), 1883–1885. <https://doi.org/10.7860/JCDR/2013/5855.3341>
- Neviani, F., Belvederi Murri, M., Mussi, C., Triolo, F., Toni, G., Simoncini, E., Tripi, F., Menchetti, M., Ferrari, S., Ceresini, G., Cremonini, A., Bertolotti, M., Neri, G., Squatrito, S., Amore, M., Zanetidou, S., & Neri, M. (2017). Physical exercise for late life depression: effects on cognition and disability. *International Psychogeriatrics*, 29(7), 1105–1112. <https://doi.org/10.1017/s1041610217000576>
- Northey, J. M., Cherbuin, N., Pumpa, K. L., Smee, D. J., & Rattray, B. (2018). Exercise interventions for cognitive function in adults older than 50: A systematic review with meta-analysis. *British Journal of Sports Medicine*, 52(3), 154–160. <https://doi.org/10.1136/bjsports-2016-096587>
- Nowak, A., Majsterek, I., Przybyłowska-Sygut, K., Pytel, D., Szymanek, K., Szaflik, J., & Szaflik, J. P. (2015). Analysis of the expression and polymorphism of APOE, HSP, BDNF, and GRIN2B genes associated with the neurodegeneration process in the pathogenesis of primary open angle glaucoma. *BioMed Research International*, Article 258281. <https://doi.org/10.1155/2015/258281>
- Oertel-Knöchel, V., Mehler, P., Thiel, C., Steinbrecher, K., Malchow, B., Tesky, V., Ademmer, K., Prvulovic, D., Banzer, W., Zopf, Y., Schmitt, A., & Hänsel, F. (2014). Effects of aerobic exercise on cognitive performance and individual psychopathology in depressive and schizophrenia patients. *European Archives of Psychiatry and Clinical Neuroscience*, 264(7), 589–604. <https://doi.org/10.1007/s00406-014-0485-9>
- Olson, R. L., Brush, C. J., Ehmann, P. J., & Alderman, B. L. (2017). A randomized trial of aerobic exercise on cognitive control in major depression. *Clinical Neuropsychology*, 128(6), 903–913. <https://doi.org/10.1016/j.clinph.2017.01.023>
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J., Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S., & Moher, D. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *International Journal of Surgery*, 88, Article 105906. <https://doi.org/10.1016/j.ijsu.2021.105906>
- Pan, Z., Park, C., Brietzke, E., Zuckerman, H., Rong, C., Mansur, R. B., Fus, D., Subramanipillai, M., Lee, Y., & McIntyre, R. S. (2019). Cognitive impairment in major depressive disorder. *CNS Spectrums*, 24(1), 22–29. <https://doi.org/10.1017/S1092852918001207>
- Patel, S., Keating, B. A., & Dale, R. C. (2023). Anti-inflammatory properties of commonly used psychiatric drugs. *Frontiers in Neuroscience*, 16, Article 1039379. <https://doi.org/10.3389/fnins.2022.1039379>
- R Core Team. (2013). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. <http://www.r-project.org/index.html>
- Ren, F. F., Alderman, B. L., Wang, W. G., Chen, F. T., Zhou, W. S., Zong, W. J., Liang, W. M., & Chang, Y. K. (2023). Effects of exercise training on executive functioning in adults with depression: A systematic review and meta-analysis of randomized controlled trials. *Sports Medicine*, 1–24. <https://doi.org/10.1007/s40279-023-01869-2>
- Ren, F. F., Feng, S. H., Li, R. H., Chu, C. H., Chang, Y. K., & Chen, F. T. (2023). The effects of acute aerobic and resistance exercise on the planning aspect of executive functions in children with preterm birth. *International Journal of Sport and Exercise Psychology*, 1–14. <https://doi.org/10.1080/1612197X.2023.2268852>
- Rock, P. L., Roiser, J. P., Riedel, W. J., & Blackwell, A. D. (2014). Cognitive impairment in depression: A systematic review and meta-analysis. *Psychological Medicine*, 44(10), 2029–2040.
- Rosenblat, J. D., Kakar, R., & McIntyre, R. S. (2016). The cognitive effects of antidepressants in major depressive disorder: A systematic review and meta-analysis of randomized clinical trials. *International Journal of Neuropsychopharmacology*, 19(2). <https://doi.org/10.1093/ijnp/pyv082>
- Salehi, I., Hosseini, S. M., Haghighi, M., Jahangard, L., Bajoghli, H., Gerber, M., Pühse, U., Holsboer-Trachsler, E., & Brand, S. (2016). Electroconvulsive therapy (ECT) and aerobic exercise training (AET) increased plasma BDNF and ameliorated depressive symptoms in patients suffering from major depressive disorder. *Journal of Psychiatric Research*, 76, 1–8. <https://doi.org/10.1016/j.jpsychires.2016.01.012>
- Sanders, L. M. J., Hortobágyi, T., la Bastide-van Gemert, S., van der Zee, E. A., & van Heuvelen, M. J. G. (2019). Dose-response relationship between exercise and cognitive function in older adults with and without cognitive impairment: A systematic review and meta-analysis. *PLoS One*, 14(1), Article e0210036. <https://doi.org/10.1371/journal.pone.0210036>
- Sepehrmanesh, Z., Fahimi, H., Akasheh, G., Davoudi, M., Gilasi, H., & Ghaderi, A. (2017). The effects of combined sertraline and aspirin therapy on depression severity among patients with major depressive disorder: A randomized clinical trial. *Electronic Physician*, 9(11), 5770–5777. <https://doi.org/10.19082/5770>
- Sheline, Y. I., Wang, P. W., Gado, M. H., Csernansky, J. G., & Vannier, M. W. (1996). Hippocampal atrophy in recurrent major depression. *Proceedings of the National Academy of Sciences*, 93(9), 3908–3913. <https://doi.org/10.1073/pnas.93.9.3908>
- Shimada, T., Ito, S., Makabe, A., Yamanushi, A., Takenaka, A., Kawano, K., & Kobayashi, M. (2022). Aerobic exercise and cognitive functioning in schizophrenia: An updated systematic review and meta-analysis. *Psychiatry Research*, 314, Article 114656. <https://doi.org/10.1016/j.psychres.2022.114656>
- Shu, Y., He, Q., Xie, Y., Zhang, W., Zhai, S., & Wu, T. (2020). Cognitive gains of aerobic exercise in patients with ischemic cerebrovascular disorder: A systematic review and meta-analysis. *Frontiers in Cell and Developmental Biology*, 8, Article 582380. <https://doi.org/10.3389/fcell.2020.582380>
- Smith, P. J., Blumenthal, J. A., Hoffman, B. M., Cooper, H., Strauman, T. A., Welsh-Bohmer, K., Brownlyke, J. N., & Sherwood, A. (2010). Aerobic exercise and neurocognitive performance: A meta-analytic review of randomized controlled trials. *Psychosomatic Medicine*, 72(3), 239–252. <https://doi.org/10.1097/PSY.0b013e3181d14633>
- Snyder, H. R. (2013). Major depressive disorder is associated with broad impairments on neuropsychological measures of executive function: A meta-analysis and review. *Psychological Bulletin*, 139(1), 81–132. <https://doi.org/10.1037/a0028727>
- Sun, M., Lancot, K., Herrmann, N., & Gallagher, D. (2018). Exercise for cognitive symptoms in depression: A systematic review of interventional studies. *Canadian Journal of Psychiatry*, 63(2), 115–128. <https://doi.org/10.1177/0706743717738493>
- Ten Brinke, L. F., Bolandzadeh, N., Nagamatsu, L. S., Hsu, C. L., Davis, J. C., Miran-Khan, K., & Liu-Ambrose, T. (2015). Aerobic exercise increases hippocampal volume in older women with probable mild cognitive impairment: A 6-month randomised controlled trial. *British Journal of Sports Medicine*, 49(4), 248–254. <https://doi.org/10.1136/bjsports-2013-093184>
- Tomoto, T., Liu, J., Tseng, B. Y., Pasha, E. P., Cardim, D., Tarumi, T., Hynan, L. S., Munro Cullum, C., & Zhang, R. (2021). One-year aerobic exercise reduced carotid arterial stiffness and increased cerebral blood flow in amnesic mild cognitive impairment. *Journal of Alzheimer's Disease*, 80(2), 841–853. <https://doi.org/10.3233/jad-201456>
- Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, 36(3), 1–48. <https://doi.org/10.18637/jss.v036.i03>
- Viechtbauer, W., & Cheung, M. W. (2010). Outlier and influence diagnostics for meta-analysis. *Research Synthesis Methods*, 1(2), 112–125. <https://doi.org/10.1002/jrsm.11>
- White, L. J., & Castellano, V. (2008). Exercise and brain health—implications for multiple sclerosis: Part 1—neuronal growth factors. *Sports Medicine*, 38(2), 91–100. <https://doi.org/10.2165/00007256-200838020-00001>
- Wilckens, K. A., Stillman, C. M., Waiwood, A. M., Kang, C., Leckie, R. L., Peven, J. C., Foust, J. E., Fraundorf, S. H., & Erickson, K. I. (2021). Exercise interventions preserve hippocampal volume: A meta-analysis. *Hippocampus*, 31(3), 335–347. <https://doi.org/10.1002/hipo.23292>
- Xiu, M., Fan, Y., Liu, Q., Chen, S., Wu, F., & Zhang, X. (2023). Glucose metabolism, hippocampal subfields and cognition in first-episode and never-treated schizophrenia. *International Journal of Clinical and Health Psychology*, 23(4), Article 100402. <https://doi.org/10.1016/j.ijchp.2023.100402>
- Xu, Y., Cai, Z., Fang, C., Zheng, J., Shan, J., & Yang, Y. (2022). Impact of aerobic exercise on cognitive function in patients with schizophrenia during daily care: A meta-analysis. *Psychiatry Research*, 312, Article 114560. <https://doi.org/10.1016/j.psychres.2022.114560>
- Zhang, S., Zhen, K., Su, Q., Chen, Y., Lv, Y., & Yu, L. (2022). The effect of aerobic exercise on cognitive function in people with Alzheimer's disease: A systematic review and meta-analysis of randomized controlled trials. *International Journal of Environmental Research and Public Health*, 19(23), 15700. <https://doi.org/10.3390/ijerph192315700>
- Zhang, Y., Wang, T., Chen, M., Lou, H., Ye, J., Shi, J., & Wen, X. (2023). Effects of moderate-intensity aerobic exercise on cognitive fatigue relief: A randomised self-controlled study. *International Journal of Sport and Exercise Psychology*, 1–19. <https://doi.org/10.1080/1612197X.2023.2229358>
- Zhao, S., Chi, A., Yan, J., & Yao, C. (2020). Feature of heart rate variability and metabolic mechanism in female college students with depression. *BioMed Research International*, 2020, Article 5246350. <https://doi.org/10.1155/2020/5246350>
- Zhdanov, A., Atluri, S., Wong, W., Vaghei, Y., Daskalakis, Z. J., Blumberger, D. M., Frey, B. N., Giacobbe, P., Lam, R. W., Milev, R., Mueller, D. J., Turecki, G., Parikh, S. V., Rotzinger, S., Soares, C. N., Brenner, C. A., Vila-Rodriguez, F., McAndrews, M. P., Kleffner, K., & Farzan, F. (2020). Use of machine learning for predicting escitalopram treatment outcome from electroencephalography recordings in adult patients with depression. *JAMA Network Open*, 3(1), Article e1918377. <https://doi.org/10.1001/jamanetworkopen.2019.18377>