

The literature on the subject suggests an association between the neurotoxicity generated by beta-lactam antibiotics and their ability to interact with gamma-aminobutyric acid (GABA) receptors.⁶ However recent studies in rats suggest the involvement of other factors, such as increased release of excitatory aminoacids,⁷ which has yet to be demonstrated.

Considering the above, we conclude that in patients who are receiving treatment with carbapenem and present neurological alterations or altered mental state, especially in cases of patients with CKD or elderly patients, we should consider an adverse effect of the drug to be a potential cause after ruling out other possibilities. According to the pattern observed in these cases, symptoms can persist up to 2 weeks after suspension of the drug.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

1. Morales RI. Ertapenem: una nueva clase de carbapenem. *Rev Chil Infect.* 2003;20:270–6.
2. Musson D, Majumdar A, Birk K, Sherry H, Wickersham P, Li S, et al. Pharmacokinetics of intramuscularly administered

- ertapenem. *Antimicrob Agents Chemother.* 2003;47:1732–5.
3. Wen MJ, Sung CC, Chau T, Lin SH. Acute prolonged neurotoxicity associated with recommended doses of ertapenem in 2 patients with advanced renal failure. *Clin Nephrol.* 2013;80:474–8.
4. Dupuaine S, Kitchell E, Tate T, Tannen RC, Wickremasinghe IM. Central nervous system associated with ertapenem use. *Ann Pharmacother.* 2011;45:e6.
5. Naranjo CA. A clinical pharmacologic perspective on the detection and assessment of adverse drug reaction. *Drug Inf J.* 1986;20:387–93.
6. Miller AD, Ball AM, Bookstaver PB, Dornblaser EK, Bennett CL. Epileptogenic potential of carbapenem agents: mechanism of action, seizure rates, and clinical considerations. *Pharmacotherapy.* 2011;31:408–23.
7. De Sarro A, Ammendola D, Zappala M, Grasso S, de Sarro GB. Relationship between structure and convulsant properties of some beta-lactam antibiotics following intracerebroventricular microinjection in rats. *Antimicrob Agents Chemother.* 1995;39:232–7.

R. Padilla Peinado*, J. Esteban Fernández,
S. Rodríguez Álvarez, T. Villa Albuguer

Servicio de Medicina Interna, Hospital Universitario de Getafe, Getafe, Madrid, Spain

*Corresponding author.

E-mail addresses: Rebeca.padilla.p@gmail.com,
rbk_b12@hotmail.com (R. Padilla Peinado).

Logopenic progressive aphasia associated with idiopathic Parkinson's disease☆☆☆



Afasia progresiva logopénica asociada a enfermedad de Parkinson idiopática

Dear Editor,

Primary progressive aphasia is a clinical syndrome characterised by a language deficit of neurodegenerative origin with no other cognitive manifestations, at least during the initial stages.¹ Three clinical variants have been described to date (nonfluent, semantic, and logopenic), each associated with its distinct topography and anatomical pathology.³ Of the 3 variants, logopenic aphasia is mainly associated with Alzheimer disease, and it is considered

an atypical form of AD onset.¹ However, the association between logopenic aphasia and Alzheimer disease is still a matter of debate in the literature. Associations with other diseases have been found in a high percentage of cases in studies using molecular imaging, cerebrospinal fluid biomarkers, or anatomical pathology findings.^{3,4} Furthermore, cognitive impairment and dementia associated with Parkinson's disease are regarded as frequent. They are characterised by executive and/or memory deficits, while language typically remains preserved.⁵ We present the case of a patient with idiopathic Parkinson's disease who developed symptoms of progressive logopenic aphasia. Her symptoms finally progressed to generalised dementia with biomarkers of Alzheimer disease, thereby supporting the idea that this type of aphasia is a marker of Alzheimer disease.

Our patient is a 65-year-old woman with high blood pressure and dyslipidaemia, who in 2009 was diagnosed with idiopathic Parkinson's disease after a one-year period in which she displayed slowness and tremor. Neurological examination revealed rigidity, bradykinesia, and resting tremor predominantly on the right side. She responded favourably to levodopa and subsequently developed motor fluctuations. She also presented REM sleep behaviour disorder. Since mid 2012, the patient began reporting increasing difficulty finding words without any associated memory impairment or behaviour disorder. She experienced no hallucinations and her neurological examination did not

☆ Please cite this article as: Matías-Guiu JA, García-Ramos R, Cabrera-Martin MN, Moreno-Ramos T. Afasia progresiva logopénica asociada a enfermedad de Parkinson idiopática. *Neurología.* 2015;30:521–524.

☆☆ This study was presented at the 65th Annual Meeting of the Spanish Society of Neurology (Barcelona, 19–23 November 2013).

reveal oculomotor alterations, pyramidal signs, or cerebellar signs. Neuropsychological evaluation performed in early 2013 showed qualitative impairment of language fluency consisting of frequent pauses to find words, with no aphasic transformations and well-articulated speech. We observed anomic aphasia both in visual-verbal naming and verbal-verbal naming, and conduction aphasia, with preserved verbal comprehension, grammatical structures, and semantic usage. She also presented a mild divided attention deficit (TMT-Part B). Verbal memory tested as borderline, while the rest of the evaluation yielded normal results (Table 1). The PET-TC study showed left temporoparietal hypometabolism (Fig. 1) and magnetic resonance scan of

that region revealed asymmetry. Analysis of cerebrospinal fluid showed a decreased concentration of A- β 42 proteins (199 pg/mL, normal value > 500 pg/mL), increased T-protein (443 pg/mL, normal value < 400 pg/mL), and increased phosphotau levels (62 pg/mL, normal value < 61 pg/mL). Over the following months, the patient's clinical symptoms progressed and her speech symptoms grew more marked. Memory deficit, disorientation, and functional decline also began to manifest.

The main signs of aphasia and findings from the FDG-PET study met diagnostic criteria for the logopenic variant of primary progressive aphasia,² which subsequently progressed to overall cognitive impairment. In this case, we

Table 1 Results from the neuropsychological evaluation.

	Raw score	Scaled score ^b
MMSE	26/30	
Addenbrooke's Cognitive Examination	63/100	
Orientation	35/35	18
Boston Naming Test	31/60	5
Verbal-verbal naming. Answer the question ^a	11/18	Deficient
Verbal-verbal naming. Complete the sentence ^a	18/18	Normal
Token Test	30.5	8
Word comprehension ^a	36/36	Normal
Word comprehension (body parts) ^a	18/18	Normal
Comprehension of instructions ^a	14/16	Normal
Repetition of syllables ^a	8/8	Normal
Repetition of syllable pairs ^a	8/8	Normal
Repetition of logotomes ^a	8/8	Normal
Word repetition ^a	10/10	Normal
Repetition of sentences ^a	44/60	Deficient
Pyramids and palm trees test	47/52 (90.4%)	Normal
Orobuccal apraxia ^a	20/20	Normal
Direct and inverse verbal span	5/3	10/9
Trail Making Test A/B	106/356	5/4
Stroop A/B/C	63/31/26	7/5/9
Symbol Digit Modalities Test	19	8
Symbolic gesture, on command and imitation (one hand) ^a	10/10	Normal
Mimicry of object use, on command and imitation ^a	10/10	Normal
Imitation of postures: unilateral, bilateral ^a	10/10; 7/8	Normal
Basic constructive praxis: copying ^a	14/18	Normal
Rey-Osterrieth complex figure: copying	30	9
Rey-Osterrieth complex figure: 3 min	6.5	7
Rey-Osterrieth complex figure: 30 min	6.5	7
Free and Cued Selective Reminding Test. Total free recall	10	6
Free and Cued Selective Reminding Test. Total recall	28	6
Free and Cued Selective Reminding Test. Delayed free recall	4	6
Free and Cued Selective Reminding Test. Delayed total recall	8	6
Semantic fluency task (animals in 1 min)	8	3
Lexical fluency task (words beginning with 'P' in 1 min)	7	7
Tower of London test. Correct movements	2	8
Visual object and space perception battery. Object decision	17	11
Visual object and space perception battery. Progressive silhouettes	12	10
Visual object and space perception battery. Position discrimination	18	6
Visual object and space perception battery. Number location	10	18

^a Subtest from the revised Barcelona Test.

^b According to patient's age and years of education (8 years), using normative data from Neuronorma project. A scaled score less than or equal to 5 is considered deficient.¹⁰




Figure 1 FDG-PET scan showing left temporoparietal hypometabolism compatible with the logopenic variant of progressive aphasia.

might suggest the association of 2 neurodegenerative diseases in the same patient (Parkinson's disease and Alzheimer disease), or else that logopenic aphasia had manifested as the initial form of dementia associated with Parkinson's disease. Parkinson's disease and REM sleep behaviour disorder are indicative of dementia with Lewy bodies and alpha-synuclein aggregates, while biomarkers in cerebrospinal fluid and the cognitive profile point to Alzheimer disease. An association between these 2 diseases is considered frequent in dementia associated with Parkinson's disease, and researchers have indicated that copresence of Alzheimer disease could influence the clinical presentation of the former.⁶ Recent anatomical pathology studies suggest that we recognise 2 main subgroups of dementia associated with Parkinson's disease: one subgroup characterised by neocortical alpha-synuclein deposition and another also featuring A β deposition. However, deposition of neocortical Tau associated with A β would be less frequent. These data seem to support the hypothesis that amyloid disease would also contribute to the pathogenesis of cognitive impairment in Parkinson's disease; in contrast, an association between Alzheimer disease (Tau and A β deposition) and Parkinson's disease would be less frequent.⁷ Clinical distinction between subgroups, and especially determining which patients have concomitant amyloid pathology, could be relevant from a therapeutic point of view.

To our knowledge, only one other case of progressive aphasia in a patient with idiopathic Parkinson's disease has been reported in the literature.⁸ That patient was a woman who also presented hallucinations. Her FDG-PET study displayed more generalised impairment, and the PET scan with Pittsburgh compound B yielded negative results. These findings indicated that symptoms could be the result of cortical involvement in Lewy body disease.

In conclusion, presence of progressive aphasia in patients with idiopathic Parkinson's disease suggests that this type of aphasia may also represent the onset of cognitive impairment, as do other clinical manifestations of posterior cortical deficit.⁹ In our case, detection of biomarkers of Alzheimer disease indicates that the presence of logopenic aphasia could also be a marker of amyloid disease in these patients.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

1. Matías-Guiu JA, García-Ramos R. Primary progressive aphasia: from syndrome to disease. *Neurología*. 2013;28:366–74.
2. Gorno-Tempini ML, Hillis AE, Weintraub S, Kertesz A, Mendez M, Cappa SF, et al. Classification of primary progressive aphasia and its variants. *Neurology*. 2011;76:1006–14.
3. Grossman M. Primary progressive aphasia: clinicopathological correlations. *Nat Rev Neurol*. 2010;6:88–97.
4. Teichmann M, Kas A, Boutet C, Ferrieux S, Nogues M, Samri D, et al. Deciphering logopenic primary progressive aphasia: a clinical, imaging, and biomarker investigation. *Brain*. 2013;136:3478–88.
5. Emre M, Aarsland D, Brown R, Burn DJ, Duyckaerts C, Mizuno Y, et al. Clinical diagnostic criteria for dementia associated with Parkinson's disease. *Mov Dis*. 2008;22:1689–707.
6. Irwin DJ, White MT, Toledo JB, Xie SX, Robinson JL, Van Deerlin V, et al. Neuropathologic substrates of Parkinson's disease dementia. *Ann Neurol*. 2012;72:587–98.
7. Kotzbaauer PT, Cairns NJ, Campbell MC, Willis AW, Racette BA, Tabbar SD, et al. Pathologic accumulation of α -synuclein and A β in Parkinson disease patients with dementia. *Arch Neurol*. 2012;69:1326–31.
8. Sakai K, Ono K, Harada H, Shima K, Notoya M, Yamada M. Parkinson's disease showing progressive conduction aphasia. *Neurol Sci*. 2012;33:399–402.
9. Villa-Bonomo C, Pagonabarraga J, Martínez-Horta S, Fernández de Bobadilla R, García-Sánchez C, Campolongo A, et al. Short-lasting episodes of prosopagnosia in Parkinson's disease. *Parkinsonism Relat Disord*. 2013;19:375–7.
10. Peña-Casanova J, Blesa R, Aguilar M, Gramunt-Fombuena N, Gómez-Ansón B, Oliva R, et al. Spanish Multicenter Normative Studies (NEURONORMA Project): methods and sample characteristics. *Arch Clin Neuropsychol*. 2009;24:307–19.

J.A. Matías-Guiu^{a,*}, R. García-Ramos^a,
M.N. Cabrera-Martín^b, T. Moreno-Ramos^a

^a Servicio de Neurología, Hospital Clínico San Carlos,
Universidad Complutense, Madrid, Spain

^b Servicio de Medicina Nuclear, Hospital Clínico San Carlos,
Universidad Complutense, Madrid, Spain

* Corresponding author.

E-mail address: jordimatiassguiu@hotmail.com
(J.A. Matías-Guiu).

Eversion endarterectomy and re-implantation in carotid stenosis and distal kink[☆]



Endarterectomía por eversión y reimplante en una estenosis carotídea y un bucle distal

Kinking and coiling of the arteries are rare morphological alterations of the internal carotid artery (ICA) described as the most frequent sole cause of cerebrovascular insufficiency (CI), or as a combined cause together with carotid atherosclerosis. Although incidence of kinking and coiling of the ICA has been estimated to range from 10% to 16% in the general population, only 4% to 16% of these features cause symptomatic CI.^{1,2} Their natural course seems to be benign, and these alterations are asymptomatic in most cases. Morphological diagnosis is easy to determine with the help of current imaging techniques, but it is essential to determine, by means of a neurological examination and transcranial Doppler ultrasound, whether the anomaly is causing cerebral ischaemia.

This letter presents the case of a patient with carotid stenosis and distal kink.

Our patient is an 80-year-old man, a former smoker of one packet a day during 60 years with a medical history of arterial hypertension treated with 2 antihypertensive drugs (losartan and hydrochlorothiazide); he was also fitted with a cochlear implant. The patient visited the emergency department due to an episode of left-sided hemiparesia, plegic left arm, dysarthria, and amaurosis fugax of the right eye; symptoms had been present for several hours without remitting fully. Physical examination revealed left supranuclear facial paresis, distal brachial paresis with a tendency to pronation of the left forearm, and crural paresis with slow claudication after 10 seconds. Right flexor plantar reflex was observed.

A Doppler ultrasound of the supra-aortic trunks revealed a heterogeneous and irregular plaque in the right carotid bifurcation, preocclusive stenosis (70%-99%) of the right internal carotid (RIC), and homogeneous plaque with moderate stenosis in the left internal carotid (LIC). Transcranial Doppler ultrasound showed the right middle cerebral artery (MCA) to be less curved than the left one. Head CT showed necrosis on the right superior parietal postrolandic area, and signs of diffuse brain atrophy (Fig. 1). A subsequent com-

puted axial tomography angiography of the brain and neck revealed a fibrolipid atheromatous plaque on the proximal third of the RIC causing critical stenosis (90% approximately) and coiling of the distal third of the RIC. Intracranial images showed a total occlusion of the M3 segment of the MCA. The LIC showed an atheromatous plaque causing moderate stenosis (50%-70%).

The patient started treatment with antiplatelet drugs (ASA 300 mg) and statins (atorvastatin 80 mg). Twenty-four hours after the patient was admitted, his motor symptoms on the left side improved with only the tendency towards left forearm pronation persisting. Six days later, after the presurgical evaluation and brain reserve test had been carried out, the surgery was performed with the patient under general anaesthesia (Fig. 2). Surgery involved complete resection of the right ICA, eversion endarterectomy (TEA) and reimplantation of the ICA (by lateroterminal anastomosis) on the common carotid to remove the kink. The post-operative period was uneventful, with no neurological focal signs or other complications.



Figure 1 Cranial CT scan showing necrotic lesion on the right superior parietal area.

[☆] Please cite this article as: López de La Franca Beltrán B, Pérez Ramírez P, Esteban Gracia C, Llagostera Pujol S. Endarterectomía por eversión y reimplante en una estenosis carotídea y un bucle distal. *Neurología*. 2015;30:524–525.