

Scientific letter

Prognostic Value of Consolidation in Lung Tomography in Patients With Acute Respiratory Distress Syndrome



Dear Editor,

Morphological alterations evaluated by computed tomography (CT) are part of the diagnosis of acute respiratory distress syndrome (ARDS).¹ These morphological alterations include: ground glass opacification (heterogeneous density with recognizable vessels, GGO) and airspace consolidation (homogeneous density without recognizable vessels or bronchi, CO).² CO on CT has been associated with a more severe clinical course and greater progression to death.³ Furthermore, CO refers to lung parenchyma with alveolar occupation, collapse, or a combination of both. Therefore, CO would be related to oxygenation and the degree of lung static compliance (Crs). For this reason, this study aims to evaluate the relationship between CO quantification present in the CT and respiratory variables assessed during the first day of mechanical ventilation (MV).

This prospective study was conducted in the intensive care unit of the Hospital Regional Río Gallegos, Argentina. The institutional ethics board approved the study and it was subsequently registered (NCT02799940). Individuals who met criteria for ARDS according to the Berlin definition¹ and who required MV for more than 48 h were included. Patients with chronic lung disease, high risk of death within three months for reasons other than ARDS, decision to limit therapeutic effort, and unable to be transferred to radiology in the first 72 h from the diagnosis of ARDS. Patients were transferred to Radiology under MV in the first 72 h after the ARDS diagnosis. A CT scan of the chest was performed during an end-inspiratory pause. CT was evaluated and quantified according to a previous study.⁴ Briefly, the CT was assessed at three levels: apex (upper part of the aortic arch), hilum (first section below the carina), and base (2 cm above the highest diaphragm). Each level (in each of the lungs) was divided into three sections: ventral, medial and dorsal. In each of the 18 locations, the score was performed through the visualization of three morphological findings: normal lung, GGO, and CO. In each location, the occupation was quantified with a score from one to three, so the sum for each subsection should equal three. Subsequently, total GGO and total CO were considered as the sum of all GGO and CO scores from the 18 locations. Scoring was performed by two authors (RSC and CF) where consensus resolved differences.

Data were recorded on the first day of ARDS: date of ICU admission, age, gender, APACHE-II score,⁵ ARDS risk factors, PaO₂/FiO₂ ratio, lung injury score (LIS),⁶ SOFA,⁷ Vt/PBW, PEEP, RR, plateau pressure (Pplat), driving pressure (ΔP : Pplat – PEEP), oxygenation index (OI: (mean airway pressure $\times$ FiO₂ $\times$ 100)/PaO₂), respiratory minute volume (RMV: product between Vt and RR). Ventilator-free days (VFD), ICU and hospital mortality were also recorded. The end-points of the study were: relationship between total CO and Crs

(primary) and relationship between total CO and PaO₂/FiO₂; Pplat; ΔP ; RMV; OI; LIS; and VFD (secondaries). Categorical variables were presented as numbers and percentages, while continuous variables were presented as mean (SD) or median and interquartile range (IQR), as appropriate. Pearson or Spearman correlation tests were also used as appropriate. A value of $p < 0.05$ was considered statistically significant. The sample size was calculated according to a previous study.⁴

We included 29 patients. Time between the ARDS diagnosis and the CT's performance: 1 day (IQR: 0–1). Baseline characteristics and respiratory parameters are detailed in Table 1. In CT, according to quantification, 11.6% were total GGO (with the greater medial location in the ventral–dorsal gradient), 26.4% corresponded to total CO (with the greater dorsal location), while 61.9% was normal lung. There was a negative correlation between total CO and Crs: -0.38 ($p = 0.04$). In relation to the secondary results, only PaO₂/FiO₂ (r : -0.4775 , p : 0.008), ΔP (r : 0.43, p : 0.01) and VFD (r : -0.42 , p : 0.02) presented a statistically significant correlation.

The main findings of this study are (1) in patients with ARDS, CO present at baseline CT was the most frequent finding and (2) the greater the consolidation (total CO), greater alterations in the respiratory variables: Crs, ΔP , VFD and less oxygenation.

In lung CT, CO refers to absence of alveolar aeration, product of occupation, collapse or a combination of both. A plausible expression in lung mechanics of this morphological alteration should be the decrease in Crs. This pilot study supports this hypothesis, showing that the higher the CO, the lower the Crs. The importance of this result lies in the fact that the quantification of the consolidation in CT can be done, through a simple method for intensivists, without the use of any additional software. In addition, assuming the absence of mathematical coupling between variables (Crs and ΔP), we found a correlation between total CO and ΔP , which is the normalized volume according to the degree of lung compliance. In turn, this relationship, could be of importance in ventilatory management. A recent study found that the benefit of using a low Vt in ARDS varies depending on elastance (inverse of Crs), suggesting that the lung protective ventilation strategy should focus primarily on ΔP to improve elastance rather than from Vt.⁸ For this reason, the magnitude of CO in relation to mechanical parameters (Crs and ΔP), would allow us to assess the risk of injury related to the Vt according to the Crs. The magnitude of CO, in this way, would have prognostic implications. Furthermore, due to alveolar compromise, the increase in CO should be related to less oxygenation. However, the previous evidence is not conclusive about CO and oxygenation.^{9–11} The findings of this study support this association, since a negative correlation was found between total CO and PaO₂/FiO₂. Data from this study also show correlation between VFD and total CO. VFD is a measurement used in studies that address lung function in critically ill patients.¹² Its measurement focuses on the duration of MV days, and thus, indirectly, in patients with lung

Table 1
Baseline characteristics and respiratory parameters.

Number of patients	29	
Male sex, n (%)	23 (79.31)	
Age, Md (IQR)	48 (30–64)	
APACHE II, mean (SD)	18.5 (7.6)	
SOFA, mean (SD)	8.5 (3.1)	
LIS, mean (SD)	2.25 (0.5)	
VFD, Md (IQR)	12 (0–19)	
ICU mortality, n (%)	9 (31)	
Hospital mortality (%)	11 (37.93)	
Risk factor, n (%)	Pulmonary	Extrapulmonary
	CAP: 10 (34.48)	Sepsis: 7 (24.14)
	Pulmonary contusion: 5 (17.2)	Pancreatitis: 2 (6.90)
	Inhalation injury: 1 (3.4)	
	Aspiration pneumonia: 4 (13.79)	
Vt/PBW, mean (SD)	7.5 (0.9)	
RR, mean (SD)	18 (3.7)	
Pplat, mean (SD)	20 (3.8)	
PEEP, Md (IQR)	7 (5–8)	
Crs, mean (SD)	38 (9.7)	
ΔP, mean (SD)	13 (2.9)	
RMV, Md (IQR)	8100 (7500–9600)	
OI, Md (IQR)	7.73 (6.38–11.39)	
PaO ₂ /FiO ₂ , mean (SD)	144 (49.5)	

APACHE II: acute physiological and chronic health evaluation; CAP: community-acquired pneumonia; Crs: static compliance (ml/cmH₂O); ΔP: driving pressure; ICU: intensive care unit; IQR: interquartile range; LIS: lung injury score; Md: median; OI: oxygenation index; PaO₂/FiO₂: partial pressure of arterial oxygen/fraction of inspired oxygen ratio; PEEP: positive end-expiratory pressure (cmH₂O); Ppl: plateau pressure (cmH₂O); RMV: respiratory minute volume (ml/min); RR: respiratory rate; SD: standard deviation; SOFA: Sepsis related Organ Failure Assessment; VFD: ventilator-free days; Vt/PBW: predicted tidal volume (ml/kg of predicted body weight).

involvement, it would be associated with mortality.¹³ Our study has certain limitations. We used three axial slices as an expression of the entire lung and this method has been questioned, arguing that the more heterogeneous the lung is, the less representative the assessment may be with only three slices.² However, several authors have used this methodology,^{14,15} considering that it is sufficient to assess the lesions present in ARDS. In conclusion, this study evaluated the pulmonary expression of consolidation evidenced in the CT of ARDS patients. It was corroborated that the higher the degree of CO, the lower both oxygenation and Crs. Thus, the CO magnitude, assessed through a simple method, could have prognostic implications. This physiological study presented here will be useful in planning a larger study evaluating clinical endpoints, which may reaffirm the findings of this study.

Authors’ contributions

Literature search: Roberto Santa Cruz, Fernando Pagano, and Gustavo Domeniconi.
Data collection: Roberto Santa Cruz and Carolina Favot.
Study design: Roberto Santa Cruz.
Analysis of data: Roberto Santa Cruz, Carolina Favot, and Marcelo Choi.
Manuscript preparation: Roberto Santa Cruz and Gustavo Domeniconi.
Review of manuscript: All authors.

Funding

The study had no source of funding.

Conflicts of interest

The authors declare that they have no conflict of interest.

Acknowledgements

The authors would like to thank the Academic Writing Center (Universidad de Magallanes), who helped us with the writing of the

manuscript. We also want to thank Engineer Roberto de Miguel, who helped us with the Stata analysis.

References

1. Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, Fan E, et al. Acute respiratory distress syndrome: the Berlin definition. *JAMA*. 2012;307:2526–33.
2. Gattinoni L, Caironi P, Pelosi P, Goodman LR. What has computed tomography taught us about the acute respiratory distress syndrome? *Am J Respir Crit Care Med*. 2001;164:1701–11.
3. Marchiori E, Zanetti G, Fontes CA, Santos ML, Valiante PM, Mano CM, et al. Influenza A (H1N1) virus-associated pneumonia: high-resolution computed tomography–pathologic correlation. *Eur J Radiol*. 2011;80:e500–4.
4. Goodman LR, Fumagalli R, Tagliabue P, Tagliabue M, Ferrario M, Gattinoni L, et al. Adult respiratory distress syndrome due to pulmonary and extrapulmonary causes: CT, clinical, and functional correlations. *Radiology*. 1999;213:545–52.
5. Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. *Crit Care Med*. 1985;13:818–29.
6. Murray JF, Matthay MA, Luce JM, Flick MR. An expanded definition of the adult respiratory distress syndrome. *Am Rev Respir Dis*. 1988;138:720–3.
7. Vincent JL, de Mendonça A, Cantraine F, Moreno R, Takala J, Suter PM, et al. Use of the SOFA score to assess the incidence of organ dysfunction/failure in intensive care units: results of a multicenter, prospective study. Working group on “sepsis-related problems” of the European Society of Intensive Care Medicine. *Crit Care Med*. 1998;26:1793–800.
8. Goligher EC, Costa ELV, Yarnell CJ, Brochard LJ, Stewart TE, Tomlinson G, et al. Effect of lowering Vt on mortality in acute respiratory distress syndrome varies with respiratory system elastance. *Am J Respir Crit Care Med*. 2021;203:1378–85.
9. Brett SJ, Hansell DM, Evans TW. Clinical correlates in acute lung injury: response to inhaled nitric oxide. *Chest*. 1998;114:1397–404.
10. Raghu MG, Wig JD, Kochhar R, Gupta D, Gupta R, Yadav TD, et al. Lung complications in acute pancreatitis. *JOP*. 2007;8:177–85.
11. Gattinoni L, Pesenti A, Bombino M, Baglioni S, Rivolta M, Rossi F, et al. Relationships between lung computed tomographic density, gas exchange, and PEEP in acute respiratory failure. *Anesthesiology*. 1988;69:824–32.
12. Yehya N, Harhay MO, Curley MAQ, Schoenfeld DA, Reeder RW. Reappraisal of ventilator-free days in critical care research. *Am J Respir Crit Care Med*. 2019;200:828–36.
13. Schoenfeld DA, Bernard GR. Statistical evaluation of ventilator-free days as an efficacy measure in clinical trials of treatments for acute respiratory distress syndrome. *Crit Care Med*. 2002;30:1772–7.
14. Ichikado K, Suga M, Muranaka H, Guyima Y, Miyakawa H, Tsubamoto M, et al. Prediction of prognosis for acute respiratory distress syndrome with thin-section CT: validation in 44 cases. *Radiology*. 2006;238:321–9.
15. Grieser C, Goldmann A, Steffen IG, Kastrup M, Pérez Fernandez CM, Engert U, et al. Computed tomography findings from patients with ARDS due to Influenza A (H1N1) virus-associated pneumonia. *Eur J Radiol*. 2012;81:389–94.

Santa Cruz Roberto^{a,b,c,f,*}, Domeniconi Gustavo^d,
Favot Carolina^e, Pagano Fernando^f, Choi Marcelo^{g,h}

^a Servicio de Terapia Intensiva, Hospital Regional Rio Gallegos, Santa Cruz, Argentina

^b Laboratory of Physiology, School of Medicine, University of Magallanes, Punta Arenas, Chile

^c Fundación H.A. Barceló, Instituto Universitario de Ciencias de la Salud, Ciudad Autónoma de Buenos Aires, Argentina

^d Servicio de Terapia Intensiva, Sanatorio de la Trinidad de San Isidro, Buenos Aires, Argentina

^e Servicio de Radiología, Sanatorio de la Trinidad de San Isidro, Buenos Aires, Argentina

^f Servicio de Terapia Intensiva, Hospital Ramos Mejía, Ciudad Autónoma de Buenos Aires, Argentina

^g Universidad de Buenos Aires, Facultad de Farmacia y Bioquímica, Departamento de Ciencias Biológicas, Cátedra de Anatomía e Histología, Buenos Aires, Argentina

^h Universidad de Buenos Aires, CONICET, Instituto Alberto C. Taquini de Investigaciones en Medicina Traslacional (IATIMET), Buenos Aires, Argentina

* Corresponding author.

E-mail address: roberto.santacruz@umag.cl (S.C. Roberto).