



ORIGINAL ARTICLE

Spatial distribution of canine zoonotic enteroparasites in Bahía Blanca, Argentina

Luciano F. La Sala^{a,b}, Anastasia Leiboff^a, Julián M. Burgos^c, Sixto R. Costamagna^{a,*}

^a Cátedra de Parasitología Clínica, Depto. de Biología, Bioquímica y Farmacia, Universidad Nacional del Sur, Bahía Blanca, Argentina

^b Consejo Nacional de Investigaciones Científicas y Técnicas (CCT Bahía Blanca), Bahía Blanca, Buenos Aires, Argentina

^c Marine Research Institute, Reykjavík, Iceland

Received 22 August 2014; accepted 29 December 2014

Available online 19 February 2015

KEYWORDS

Dogs;
Zoonotic
enteroparasites;
Spatial distribution;
Bahía Blanca;
Quality of life

Abstract The objectives of this research were: (1) to determine the occurrence of zoonotic enteroparasites in dog feces from Bahía Blanca, Argentina; (2) to characterize the spatial distribution of the parasites found in association with the quality of life index (QLI) in neighborhoods of Bahía Blanca; and (3) to determine if the presence of a particular parasite genus in a stool sample was facilitated or impeded by the presence of other parasite genera. Samples of dog stools ($n = 475$) were collected between December 2012 and December 2013 in areas with varying QLI. The association between QLI values and the presence of parasites was analyzed using logistic regression. Overall enteroparasite occurrence was 36.6%. Parasitic forms found included nematode larvae, cysts of *Blastocystis* spp., *Giardia* spp., and oocysts of *Cryptosporidium* spp., and eggs of *Ancylostoma caninum*, *Toxocara canis*, cestodes and *Trichuris* spp. For certain enteroparasites, we detected significant associations between their occurrence and QLI. Feces collected in areas with medium and low QLI were 2.46 and 5.43 times more likely, respectively, to contain *A. caninum* than stools from the high-QLI area. Samples from areas with low QLI were 2.36 times more likely to contain *Trichuris* spp. than those from the high QLI area. Regarding protozoa, feces from areas with low QLI were 2.4 times more likely to be positive than those from areas with high QLI. We demonstrated that canine zoonotic parasites have a wide distribution in the study area, and that occurrence is higher in neighborhoods with lower QLI.

© 2014 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail addresses: rcosta@uns.edu.ar, rcostama2001@yahoo.com.ar (S.R. Costamagna).

PALABRAS CLAVE

Perros;
Enteroparásitos
zoonóticos;
Distribución espacial;
Bahía Blanca;
Calidad de vida

Distribución espacial de enteroparásitos zoonóticos de caninos en Bahía Blanca, Argentina

Resumen Los objetivos del presente trabajo fueron los siguientes: a) determinar la ocurrencia de enteroparásitos zoonóticos en heces de perros de Bahía Blanca, Argentina; b) caracterizar la distribución espacial de los parásitos hallados en función del índice de calidad de vida (ICV) en barrios de Bahía Blanca, y c) determinar si la presencia de un género parasitario en particular en las heces fue facilitada o impedida por la presencia de otro/s. Se recolectaron muestras de heces de perro ($n = 475$) durante el período diciembre 2012-diciembre 2013 en áreas con diferente ICV. La asociación entre el ICV y la presencia de parásitos fue analizada mediante regresión logística. La ocurrencia global de enteroparásitos fue del 36,6%. Las formas parasitarias detectadas correspondieron a larvas de nematodos, a quistes de *Blastocystis* spp., *Giardia* spp. y *Cryptosporidium* spp., y a huevos de *Ancylostoma caninum*, *Toxocara canis*, cestodos y *Trichuris* spp. Se hallaron asociaciones significativas entre ocurrencia de algunos enteroparásitos e ICV. Las heces recolectadas en áreas con ICV medio y bajo mostraron una probabilidad de contener *A. caninum* 2,46 y 5,43 veces mayor, respectivamente, que las recolectadas en la zona de ICV alto. Las muestras de zonas con ICV bajo tuvieron una probabilidad 2,36 veces mayor de contener *Trichuris* spp. que aquellas de áreas con ICV alto. En lo referido a protozoos, las heces de áreas con ICV bajo tuvieron una probabilidad 2,4 veces mayor de presentar quistes que las heces de áreas con ICV alto. Se demostró que los parásitos zoonóticos caninos tienen una amplia distribución en el área estudiada y que la ocurrencia es más alta en los barrios con menor ICV. © 2014 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Close relationships between people and animals date from very early in human history. The domestication of the dog (*Canis lupus familiaris*) some 12,000 years ago has enabled this species to become the most abundant carnivore in the world today, with an estimated population of 500 million. Cultural transformations over the ensuing centuries have seen a remarkable increase in the interaction between humans and domestic animals, and mainly the domestic dog¹⁹.

The benefits of contact with domestic animals, such as psychological well-being, are now well recognized. Family pets provide acceptance, joy and companionship for family members, and particularly for young children^{32,34,45}. However, this close relationship between dogs and humans also brings potential problems such as parasitic zoonoses (PZs), which represent a threat to public health. Humans most commonly become infected with these zoonotic parasites through consumption of infected food or water or via direct fecal-oral contamination. The results of these infections may vary from asymptomatic carriage to long-term morbidity and even death. Although data are still scarce, it is clear that these PZs present a significant burden for public health, particularly in poor and marginalized communities^{30,52} and especially for young children, immunosuppressed and/or malnourished people and pregnant women^{8,33,37,40,54}. Moreover, PZs can lead to significant economic losses, both directly through their adverse effects on human and animal health, and indirectly through the control measures required in the food production chain⁵².

In recent years, there has been a fairly large amount of research regarding the presence of enteroparasites in

domestic dog populations worldwide^{5,7,22,43}. Comparatively, fewer studies exist in Argentina^{3,12,38,47,56}; however, only a minority addressed the spatial distribution of infection^{26,46}.

Several studies reported elevated prevalence of intestinal parasites, mostly among children dwelling in the peripheral and poorest areas of Bahía Blanca^{13,15,17}, in vegetable gardens grown in the area (Visciarelli pers. comm.), and in dog feces collected from the streets⁶.

Local and updated information is essential to understand the epidemiology of gastrointestinal parasitic infections in dogs and to design preventive strategies at local, national, or regional levels. With this background, the objectives of this study were: (1) to determine the occurrence of zoonotic enteroparasites in dog feces from the city of Bahía Blanca, Argentina; (2) to characterize the spatial distribution of the parasites found in association with the quality of life index (QLI) in neighborhoods of Bahía Blanca; and (3) to determine if the presence of a particular parasite genus in a sample was facilitated or impeded by the presence of other parasite genera.

Materials and methods**Study area sampling design**

Bahía Blanca (38°44' S, 62°16' W) is the third largest city in Buenos Aires province, Argentina, with a population of a little over 300,000 people and a growth rate of 6% reported for the period 2001–2010. Bahía Blanca presents profound social inequalities leading to heterogeneous spatial structuring of well-being levels⁴¹. Temperature extremes are notably

frequent, with an average daily mean of 15.1 °C (ranges 8.9–21.9 °C). Annual precipitation averages 613.7 mm.

A prospective study was conducted between December 2012 and December 2013 in different neighborhoods of the city of Bahía Blanca, Argentina. Considering the second objective of this work (to analyze parasite occurrence as a function of quality of life), our sampling scheme was done by following Prieto⁴¹, who defined four levels for the quality of life in the city (very high, high, low, and very low) by considering variables such as housing quality, educational level, and indicators of public and environmental health. Considering a dog population of 68,000 dogs¹⁴ in the city, an infection prevalence of 50% (the most conservative value) for each of the parasites studied, a relative maximum error of 4.5%, and a 95% confidence level, the minimum sample size calculated was 475 stool samples for the entire study area. Then, fresh dog stool samples were collected in neighborhoods comprising only areas with very high ($n=244$), high ($n=131$) and low ($n=100$) QLI. Because of the safety risks implied, samples were not collected in the area with very low QLI. If two samples were separated by less than 30 m, only one was collected. This approach was used to minimize the chances of double sampling feces from the same dog, which could falsely inflate occurrence values.

A complete record of small animal veterinary attention centers (henceforth referred to as VACs) in the city was provided by the District's professional association of veterinarians.

Sample processing

Stool specimens were collected from the sidewalks and were placed in individual sterile plastic jars with screw caps, transported to the laboratory, and fixed in 10% formalin until assayed. Samples were sieved through a single gauze followed by concentration using the Ritchie method. Finally, they were observed under a light microscope. The presence of *Cryptosporidium* spp. was assessed using Kinyoun acid-fast stain. All samples were observed during 5 min. The eggs, cysts, and oocysts found were identified according to morphological and staining characteristics under light microscopy.

Statistical analysis

Since the results were originated from stool samples in the environment, and not from individual dogs, we used the term "occurrence" (expressed as percentage), which we considered more appropriate than "prevalence". Therefore, overall occurrence represents the proportion of fecal samples that were positive for at least one parasitic form (i.e. eggs, cysts, oocysts, adults), whereas specific occurrence was defined as the proportion of samples that were positive for each of the recorded parasite forms. For overall occurrence, feces were classified as positive if at least one of the subsamples had a parasitic form, and negative if otherwise. The same approach was used to define specific occurrence.

Results for *Giardia* spp., *Blastocystis* spp., and *Cryptosporidium* spp. were grouped into one single category as protozoa. Then, protozoa occurrence was defined as the

percentage of samples that were positive for at least one of the protozoan species observed.

The association between the independent variable QLI (categorical: very high, high, low) and the outcome variables (overall occurrence, specific occurrence, protozoa occurrence; all binomial), as well as their interaction, was assessed using generalized linear models (GLM; logistic regression with *logit link* function). In this study, QLI was the only independent variable included in the model because (1) an unknown proportion of owners take their dogs to VACs outside their corresponding neighborhood (mostly owners from the area with a very high QLI value), and (2) QLI could be used as proxy for other unmeasured and epidemiologically relevant variables (e.g. family income, responsible pet ownership).

The model's goodness of fit was assessed by comparing the full model with the corresponding intercept-only model using the Akaike information criterion (AIC)¹ as described in Burnham and Anderson¹⁰. A model was considered to provide substantial support of the data if AIC increased by more than 2 units after removing the independent variable QLI (intercept-only model).

The strength of association between exposure and outcome was assessed using the occurrence ratios (OcR) as an approximation to the prevalence ratios frequently reported in cross-sectional studies. For example, if parasite occurrence is higher in the exposed group compared with the unexposed one (e.g. low QLI vs. very high QLI), the ratio will be greater than one; if occurrence is the same, the ratio will be close to one, and if occurrence is lower in the exposed group, the ratio will be less than one. The level of significance for statistical analyses was defined as $p < 0.05$.

To assess if the presence of any particular parasite was facilitated or impeded by the presence of other parasite genera, the frequency distribution of the number of parasitic taxa per sample was tested for a Poisson distribution using a goodness-of-fit test that included samples from all neighborhoods.

All statistical analyses were done using the R and Quantum GIS softwares^{42,44}.

Results

Parasite occurrence and spatial distribution

The occurrence (expressed as percentage) of enteroparasites in dog feces from Bahía Blanca, Buenos Aires, Argentina is shown in Fig. 1.

Overall parasite occurrence was 36.6%. The highest occurrence was recorded for nematode larvae (22.3%), followed by *Ancylostoma caninum* eggs (21.1%), protozoan cysts (19.8%), trichurid eggs (18.1%), *Toxocara canis* eggs (2.3%), and cestode eggs (0.6%). When protozoa were analyzed separately (all three areas combined), occurrence was 1.4% (95% CI: 0.1–5.4) for *Giardia* spp., 2.9% (95% CI: 0.9–7.4) for *Blastocystis* spp., and 14.7% (95% CI: 0.1–22.3) for *Cryptosporidium* spp.

No significant differences were detected between QLI and overall occurrence (Table 1). However, for some parasites, occurrence varied significantly among neighborhoods with different QLI values (Table 2).




Figure 1 Maps showing the occurrence (expressed as percentage) of enteroparasites in dog feces from Bahía Blanca, Buenos Aires, Argentina. Images show: neighborhoods and their quality of life index (A), overall occurrence of enteroparasites (B), and specific occurrences of *A. caninum* (C), *Trichuris* spp. (D), nematode larvae (E), protozoa (F), *Toxocara* spp. (G), and cestodes (H).

VACs were considerably more abundant in neighborhoods with very high and high QLI, compared with areas with low QLI. The number of VACs was strongly correlated ($R=0.89$) with the QLI of different areas as defined by Prieto (2008) (very high = 36; high = 34; low = 3). These results justified the exclusion of the VACs variable from the logistic regression model.

Parasite competition

The frequency distribution of the number of species per stool sample (all neighborhoods combined) was described by a Poisson distribution (mean = 0.69; variance = 1.2; $\chi^2 = 152.3$; $p < 0.0001$), indicating that the presence of a parasite genus in individual samples was neither facilitated nor

Table 1 Overall occurrence of parasites (expressed as percentage) followed by 95% CI and the number of parasitic genera are reported for areas with different QLI in Bahía Blanca, Argentina.

Neighborhood	Occurrence (95% CI)	P^a	No. of parasite genera in positive stools (%) ^b					
			1	2	3	4	5	6
Very high QLI	38.5 (32.6–44.8)	–	66.3	12.8	15.1	5.8	0.0	0.0
High QLI	29.8 (22.6–38.1)	0.092	33.3	33.3	28.2	5.1	0.0	0.0
Low QLI	49.0 (39.4–58.7)	0.074	38.8	2.0	30.6	24.5	2.0	2.0

^a Level of significance for the comparison between the baseline level (very high QLI) and reference categories (high and low QLI).

^b Calculated as the percentage of samples among the positive ones by neighborhood.

Table 2 Logistic regression models describing the association between areas with different quality of life and occurrence (expressed as percentage) of enteroparasites in dog feces from Bahía Blanca, Buenos Aires, Argentina.

Parasite and QLI	Oc (95% CI) ^a	OcR (95% CI) ^b	p-Value	ΔAIC ^c
<i>Ancylostoma caninum</i>				
Very high QLI ^d	9.0 (6.0–13.3)			
High QLI	22.1 (15.8–30.0)	2.46 (1.47–4.10)	<0.001	60.0
Low QLI	49.0 (39.4–58.6)	5.43 (3.48–8.49)	<0.0001	
<i>Trichuris</i> spp.				
Very high QLI ^d	12.7 (9.1–17.5)			
High QLI	19.1 (13.2–26.7)	1.50 (0.93–2.43)	>0.05	10.0
Low QLI	30.0 (21.9–39.6)	2.36 (1.51–3.68)	<0.001	
Nematode larvae				
Very high QLI ^d	25.4 (20.3–31.2)			
High QLI	11.4 (7.0–18.1)	0.45 (0.27–0.76)	<0.01	10.0
Low QLI	29.0 (21.0–38.6)	1.14 (0.78–1.66)	>0.05	
Protozoans				
Very high QLI ^d	5.7 (3.4–9.5)			
High QLI	6.9 (3.5–12.7)	1.2 (0.53–2.69)	>0.05	0.0
Low QLI	14.0 (8.4–22.3)	2.44 (1.21–4.93)	<0.05	
<i>Toxocara</i> spp.				
Very high QLI ^d	2.9 (1.3–6.1)			
High QLI	1.5 (0.3–6.0)	0.53 (0.11–2.53)	>0.05	0.0
Low QLI	2.0 (0.3–7.7)	0.70 (0.15–3.30)	>0.05	
Cestodes				
Very high QLI ^d	0.8 (0.1–3.2)			
High QLI	0.0 (0.0–3.5)	0.0	>0.05	0.0
Low QLI	1.0 (0.1–6.2)	1.2 (0.1–13.3)	>0.05	

^a Occurrence of parasites presented as percentage with 95% confidence interval in parenthesis.

^b Occurrence ratios (OcR) and their 95% CI.

^c Level of empirical support of models reported as ΔAIC, which represents the AIC value increment if the single term is dropped. Values of ΔAIC >2 are considered as evidence of substantial support of the data.

^d Significance (p values) are presented for the levels of main effects (neighborhood) compared with the reference category (very high QLI).

impeded by the presence of other parasite genera. The analyses showed that when *A. caninum* co-occurred with another parasite, 80% of those samples also had *Trichuris* spp. No co-occurrence associations were detected in samples with more than two parasitic genera.

The number of parasite genera in positive stools varied between one and six, and the neighborhood with low QLI had the widest range (Table 1).

Discussion

This work describes, both qualitatively and quantitatively, the occurrence of canine zoonotic enteroparasites in Bahía Blanca, Argentina, and, for the first time, the association between parasite occurrence and QLI in the studied area.

The overall occurrence found (36.6%) was near the low-end range of prevalence (37.9–77.4%) reported by other authors in urban and rural areas of Argentina^{35,47,49,50}, but fell within the range reported in other countries; e.g., 17.6% in Prague²¹, 28.7% in Australia⁹, 35.5% in Venezuela⁴³, 34.8% in the United States³¹, 18–42% in Japan⁵, 53% in Hungary²⁵,

68% in Nigeria⁴, 71.3% in Spain³⁶, and 85% in Mexico²². Here, we identified eight species of parasites, which fall within the range of 6–12 species documented in dogs worldwide²⁶.

In the present work, we characterized the spatial distribution of canine zoonotic enteroparasites in Bahía Blanca according to different QLI values. Although we found no significant association between parasite overall occurrence and QLI, the specific occurrence level for *Ancylostoma caninum*, *Trichuris* spp., nematode larvae, and protozoa was spatially structured following a heterogeneous pattern with higher values in neighborhoods with lower QLI (Table 2).

A. caninum showed the largest and most significant differences in occurrence between areas with different QLI. When considering all the neighborhoods, occurrence was not particularly high (21.1%), being 3.5 times lower than the 73% value reported for an urban area in Zambia¹¹ and very similar to the 21.6% value reported in southern Buenos Aires province²⁶.

In Bahía Blanca there are no reported cases of cutaneous *larva migrans*, which could be attributable to environmental characteristics (e.g. low average ambient temperature) and cultural factors such as shoe wearing. In our study,

the highest occurrence of *A. caninum* eggs in fecal samples was in Villa Nocito, which is considered one of the longest neglected areas in the city. This neighborhood has the lowest QLI among those included in the study, and has no VACs. Although no official data are available, the stray dog population in this area is suspected to be larger, and pet owners are more prone to letting their dogs roam free. This scenario, coupled with the presence of *A. caninum* in stools from this area, would be expected to predispose the human population, and mainly children, to cutaneous *larva migrans*.

Trichuriasis is a disease associated with infection by nematode *Trichuris* spp. with an estimated 1 billion infected people worldwide in 2002²⁰. This disease poses a huge burden on some areas of the world such as sub-Saharan Africa, where estimations suggest that 24% of people are infected²⁹. In the present study, occurrence of *Trichuris* spp. was higher as the QLI decreased, and overall occurrence (18.1%) was almost twice as high as that reported in other research works conducted in urban areas of Argentina (10.1%)²⁶. Similarly, occurrence in Villa Nocito (30.0%) was almost two-fold higher than the maximum level (17.9%) reported by Fontanarrosa et al.²⁶.

Moreover, nematode larvae found here were not identified, which precludes drawing any definite conclusions about their zoonotic importance.

In our research, overall occurrence of *Toxocara* spp. was low (2.3%) and was not higher in the area with lowest QLI, where only 2% of samples were positive. Previous research in people from Villa Nocito reported the presence of antibodies against *Toxocara* spp. in 18% of the studied subjects¹⁸, whereas another work found that 83% of water samples from a natural stream that crosses the city were contaminated with *Toxocara* spp. eggs¹⁶. Occurrence of *Toxocara* spp. in the present study was remarkably low compared with other research works conducted in Argentina, which ranged between 3.4% and 35.1% in soil samples from a subtropical city² and a dry steppe habitat in Patagonia²⁴, respectively. This finding is in agreement with a hypothesis that people living in urban areas are at lower risk for infection than those dwelling in rural regions^{27,28}. We suggest that different awareness levels regarding health practices and responsible pet ownership in rural regions and urban areas might account for the differences observed.

Regarding zoonotic protozoa, the excretion of their cysts/oocysts in the host's feces is intermittent. Therefore, the reported occurrence levels should be considered with some caution⁴⁸. Occurrence of *Giardia* spp. in the area with low QLI (14.0%) was 2.4 times higher than in the area with very high QLI (5.7%). This difference is in line with other research works¹⁵ conducted in Bahía Blanca, where prevalence of *Giardia* spp. was 9% in children stools (under 4 yr. old) dwelling in areas with very high QLI, and 33.3% in older children (4–10 yr. old) from areas with low QLI. Furthermore, the level of overall occurrence (1.4%) found here was very similar to the 1.3% reported by other authors in Argentina⁴⁹.

On the contrary, *Cryptosporidium* spp. occurred 73.5 times more frequently (14.7% vs. 0.2%) in the present study compared with related research in southern Buenos Aires province²⁶. The occurrence of *Cryptosporidium* spp. in dog feces contaminating public places raises concern about the

potential transmission of this protozoan to immunosuppressed people^{45,55}.

Blastocystis spp. was found in 2.9% (4/139) of the samples from areas with very high and high QLI combined. Recent research conducted in people from Bahía Blanca reported *Blastocystis* spp. fecal prevalence between 19.0 and 30.6%²³. However, a paucity of knowledge regarding the zoonotic potential of this pathogen precludes any conclusions about the role of dog defecation as a relevant source of infection in humans. Despite this fact, recent research proposes domestic dogs as a potential source of *Blastocystis* infection to humans³⁹. However, considering the low prevalence of this intestinal protozoan in dogs and a lack of evidence supporting a dog-specific or predominant genetic subtype, dogs are unlikely to act as natural hosts of *Blastocystis*, only becoming infected opportunistically via coprophagia of other host feces or by drinking contaminated water⁵³.

In our study, the absolute number of VACs in the neighborhoods correlated positively with the QLI. However, as mentioned before, an unknown proportion of owners take their dogs to VACs outside their corresponding neighborhood. This fact could bias any conclusions regarding the role of unequal access to VACs as a predisposing factor for enteric parasites in the studied population. Additionally, the variety of helminth species found here warrants a critical revision of deworming protocols currently used in dogs from the study area. It is worth noting that the neighborhood with lowest QLI studied here showed the largest number of parasite genera per stool sample (Table 1). This could be attributable to the markedly lower number of VACs and, consequently, of deworming practices in that area⁵¹.

The distribution of parasite genera per stool sample cannot be assumed to be significantly different from a random distribution, suggesting that in the samples assayed, the presence of a parasite genus was neither facilitated nor impeded by the presence of other parasite genera. One possible explanation for the high frequency (80%) of co-occurrence between *A. caninum* and *Trichuris* spp. is the high occurrence of these genera in the dog population from Bahía Blanca.

In conclusion, several factors associated with decreased quality of life (e.g. income, housing conditions, etc.) would seem to play an important role in the occurrence of zoonotic enteroparasites in the dog population of Bahía Blanca. Our findings will facilitate the implementation of preventive public health measures toward the control of these parasites.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that no patient data appear in this article.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Acknowledgements

We are grateful to Secretaría de Ciencia y Tecnología de la Universidad Nacional del Sur, to Fundación. Alberto J. Roemmers for financial support, and to the Municipal Government of Bahía Blanca for declaring this project of national interest (Ordinance No. 14177).

References

- Akaike H. A new look at the statistical model identification. *IEEE Trans Automat Control*. 1974;19:716–22.
- Alonso JM, Stein M, Chamorro M, Bojanich MV. Contamination of soils with eggs of *Toxocara* in a subtropical city in Argentina. *J Helminthol*. 2001;75:165–8.
- Andresiuk MV, Denegri GM, Sardella NH, Hollmann P. Encuesta coproparasitológica canina realizado en plazas públicas de la ciudad de Mar del Plata, Buenos Aires, Argentina. *Parasitol Latinoam*. 2003;58:17–22.
- Anene BM, Nnaji TO, Chime AB. Intestinal parasitic infections of dogs in the Nsukka area of Enugu State, Nigeria. *Prev Vet Med*. 1996;27:89–94.
- Asano K, Suzuki K, Matsumoto T, Sakai T, Asano R. Prevalence of dogs with intestinal parasites in Tochigi, Japan in 1979, 1991 and 2002. *Vet Parasitol*. 2004;120:243–8.
- Baillie E, Argañín L, Costamagna SR. Contaminación de la vía pública con parásitos de importancia zoonótica: en un sector de Bahía Blanca, Buenos Aires, Argentina. *Rev AMBB*. 2007;17:29–33.
- Basualdo J, Pezzani B, De Luca M, Córdoba A, Apezteguía M. Screening of the municipal watersystem of La Plata, Argentina, for human intestinal parasites. *Int J Hyg Environ Health*. 2000;203:177–82.
- Bojar I, Szymańska J. Environmental exposure of pregnant women to infection with *Toxoplasma gondii* – state of the art. *Ann Agric Environ Med*. 2010;17:209–14.
- Bugg RJ, Robertson ID, Elliot AD, Thompson RCA. Gastrointestinal parasites of urban dogs in Perth, Western Australia. *Vet J*. 1999;157:295–301.
- Burnham KP, Anderson DR. Information and likelihood theory: a basis for model selection and inference. In: Burnham KP, Anderson DR, editors. *Model selection and multimodel inference: a practical information-theoretic approach*. 2nd ed. New York: Springer-Verlag Inc.; 2002. p. 49–97.
- Bwalya EC, Nalubamba KS, Hankanga C, Namangala B. Prevalence of canine gastrointestinal helminths in urban Lusaka and rural Katete Districts of Zambia. *Prev Vet Med*. 2011;100:252–5.
- Córdoba A, Ciarmela ML, Pezzani B, Gamboa I, de Luca M, Minvielle M, Basualdo J. Presencia de parásitos intestinales en paseos públicos urbanos en La Plata Argentina. *Parasitol Latinoam*. 2002;57:25–9.
- Costamagna SR. Enfermedades parasitarias transmitidas por alimentos (EPTAs). In: Ganduglia R, editor. *Calidad y seguridad agroalimentaria*. 1st ed. Bahía Blanca, Argentina: Editorial EdIUNS; 2013. p. 39–50.
- Costamagna SR, Baillie E, Racca L, Marro A, Bin L, Pirlas M, Schiaffino L, Echenique C. Estudio seroepidemiológico prospectivo, sobre toxocarosis humana en Bahía Blanca, Argentina. *Rev Arg Parasitol*. 2012;1:25–39.
- Costamagna SR, García S, Gutiérrez M, Visciarelli E, Torno Cafasso O, Prat MI. Investigación de entero y ectoparasitosis en el área periférica de Bahía Blanca, Argentina. *Rev AMBB*. 1991;2:21–7.
- Costamagna SR, García S, Visciarelli E, Casas N. Epidemiología de las parasitosis en la ciudad de Bahía Blanca, Argentina. *Parasitol Latinoam*. 2002;57:103–10.
- Costamagna SR, López J, Torno Cafasso O, Visciarelli E, Sagua M, Dasso P. Investigación de enteroparasitosis en el área periférica de Bahía Blanca, Argentina. Parte I. *Rev AMBB*. 1998;7:13–6.
- Costamagna SR, Viscerelli E, Lucchi L, Basualdo J. Parásitos en aguas del arroyo Napostá, aguas de recreación y de consumo en la ciudad de Bahía Blanca (Provincia de Buenos Aires, Argentina). *Parasitol Latinoam*. 2005;60:122–6.
- Davis SJM, Valla FR. Evidence for domestication of the dog 12,000 years ago in the Natufian of Israel. *Nature*. 1978;276:608–10.
- de Silva NR, Booker S, Hotez PJ, Montresor A, Engels D, Savioli L. Soil-transmitted helminth infections: updating the global picture. *Trends Parasitol*. 2003;19:547–51.
- Dubná S, Langrová I, Nápravník J, Jankovská I, Vadlejch J, Pekár S, Fechtner J. The prevalence of intestinal parasites in dogs from Prague, rural areas, and shelters of the Czech Republic. *Vet Parasitol*. 2007;145:120–8.
- Eguía-Aguilar P, Cruz-Reyes A, Martínez-Maya JJ. Ecological analysis and description of the intestinal helminths present in dogs in Mexico city. *Vet Parasitol*. 2005;127:139–46.
- Felice V, Vilariño M, Hernández D, Scarafía M, Costamagna SR. Parásitos intestinales en niños y adultos que concurren a Laboratorio Privado y Hospital Público en la ciudad de Bahía Blanca, Argentina: Comparación de prevalencias. *Rev Méd Rosario*. 2011;77:S81.
- Fillaux J, Santillan G, Magnaval JF, Jensen O, Larrieu E, Sobrino-Becaria CD. Epidemiology of toxocariasis in a steppe environment: the Patagonia study. *Am J Trop Med Hyg*. 2007;76:1144–7.
- Fok E, Szatmári V, Busák K, Rozgonyi F. Prevalence of intestinal parasites in dogs in some urban and rural areas of Hungary. *Vet Q*. 2001;2:96–8.
- Fontanarrosa MF, Vezzani D, Basabe J, Eiras DF. An epidemiological study of gastrointestinal parasites of dogs from Southern Greater Buenos Aires (Argentina): age, gender, breed, mixed infections, and seasonal and spatial patterns. *Vet Parasitol*. 2006;136:283–95.
- Glickman LT, Magnaval JF, Domanski LM, Shofer FS, Lauria SS, Gottstein B, Brochier B. Visceral larva migrans in French adults: a new disease syndrome. *Am J Epidemiol*. 1987;125:1019–34.
- Herrmann N, Glickman LT, Schantz PM, Weston MG, Domanski LM. Seroprevalence of zoonotic Toxocariasis in the United States: 1971–1973. *Am J Epidemiol*. 1985;122:890–6.
- Hotez PJ, Alibek K. Central Asia's hidden burden of neglected tropical diseases. *PLoS Negl Trop Dis*. 2011;5:1–4.
- Hotez PJ, Kamath A. Neglected tropical diseases in sub-Saharan Africa: review of their prevalence, distribution, and disease burden. *PLoS Negl Trop Dis*. 2009;3:1–10.
- Kirkpatrick CE. Epizootiology of endoparasitic infections in pet dogs and cats presented to a veterinary teaching hospital. *Vet Parasitol*. 1988;30:113–24.
- Kravetz JD, Federman DG. Cat-associated zoonoses. *Arch Int Med*. 2002;162:1945–52.
- Lewthwaite P, Gill G, Hart CA, Beeching NJ. Gastrointestinal parasites in the immunocompromised. *Curr Opin Infect Dis*. 2005;18:427–35.
- Love M, Overall KL. How anticipating relationships between dogs and children can help prevent disasters. *J Am Vet Med Assoc*. 2001;219:446–53.
- Martín UO, Demonte MA. Urban contamination with zoonotic parasites in the central region of Argentina. *Medicina (Buenos Aires)*. 2008;68:363–6.

36. Martínez-Moreno FJ, Hernández S, López-Cobos E, Becerra C, Acosta I, Martínez-Moreno A. Estimation of canine intestinal parasites in Córdoba (Spain) and their risk to public health. *Vet Parasitol.* 2007;143:7–13.
37. Meamar AR, Rezaian M, Mohraz M, Zahabian F, Hadighi R, Kia EB. A comparative analysis of intestinal parasitic infections between HIV+/AIDS patients and non-HIV infected individuals. *Iran J Parasitol.* 2007;2:1–6.
38. Minvielle MC, Pezzani BC, Basualdo Farjat JA. Frequency of finding helminths eggs in canine stool samples collected in public places from La Plata city, Argentina. *Bol Chil Parasitol.* 1993;48:63–5.
39. Nagel R, Cuttall L, Stensvold CR, Mills PC, Bielefeldt-Ohmann H, Traub RJ. *Blastocystis* subtypes in symptomatic and asymptomatic family members and pets and response to therapy. *Intern Med J.* 2012;42:1187–95.
40. Pavlinová J, Kinčeková J, Ostró A, Saksun L, Vasilková Z, Königová A. Parasitic infections and pregnancy complications. *Helminthologia.* 2011;48:8–12.
41. Prieto MB. Fragmentación socio-territorial y calidad de vida urbana en Bahía Blanca. *Geograficando.* 2008;4:193–214.
42. QGIS Development Team. QGIS Geographic Information System. Open Source Geospatial Foundation Project; 2013 [accessed 25 July 2014] <http://qgis.osgeo.org>
43. Ramírez-Barrios RA, Barboza-Mena G. Prevalence of intestinal parasites in dogs under veterinary care in Maracaibo, Venezuela. *Vet Parasitol.* 2004;121:11–20.
44. R Core Team. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2013 [accessed 25 July 2014] <http://www.R-project.org>
45. Robinson RA, Pugh RN. Dogs, zoonoses and immunosuppression. *J R Soc Health.* 2002;122:95–8.
46. Rubel D, Zunino G, Santillán G, Wisnivesky C. Epidemiology of *Toxocara canis* in the dog population from two areas of different socioeconomic status, Greater Buenos Aires, Argentina. *Vet Parasitol.* 2003;115:275–86.
47. Sánchez Thevenet P, Jensen O, Mellado I, Torrecillas C, Raso S, Flores ME, Minvielle MC, Basualdo JA. Presence and persistence of intestinal parasites in canine fecal material collected from the environment in the Province of Chubut, Argentine Patagonia. *Vet Parasitol.* 2003;117:263–9.
48. Solarczyk P, Majewska AC. A survey of the prevalence and genotypes of *Giardia duodenalis* infecting household and sheltered dogs. *Parasitol Res.* 2010;106:1015–9.
49. Soriano SV, Pierangeli NB, Rocca I, Bergagna HF, Lazzarini LE, Celescinco A, Saiz MS, Kossman A, Contreras PA, Arias C, Basualdo JA. A wide diversity of zoonotic intestinal parasites infects urban and rural dogs in Neuquén, Patagonia, Argentina. *Vet Parasitol.* 2010;167:81–5.
50. Taranto NJ, Passamonte L, Marinconz R, de Marzi MC, Cajal SP, Malchiodi EL. Zoonotic parasitosis transmitted by dogs in the Chaco Salteño, Argentina. *Medicina (Buenos Aires).* 2000;60:217–20.
51. Torgerson PR, Macpherson CN. The socioeconomic burden of parasitic zoonoses: global trends. *Vet Parasitol.* 2011;182:79–95.
52. Torgerson PR, Karaeva RR, Corkeri N, Abdyjaparov TA, Kuttubaev OT, Shaikenov BS. Human cystic echinococcosis in Kyrgyzstan: an epidemiological study. *Acta Trop.* 2003;85:51–61.
53. Wang W, Cuttall L, Bielefeldt-Ohmann H, Inpankaew T, Owen H, Traub RJ. Diversity of *Blastocystis* subtypes in dogs in different geographical settings. *Parasites Vectors.* 2013;6:215–9.
54. Wiwanitkit V. Intestinal parasite infestation in HIV infected patients. *Curr HIV Res.* 2006;4:87–96.
55. Xiao L, Cama VA, Cabrera L, Ortega Y, Pearson J, Gilman RH. Possible transmission of *Cryptosporidium canis* among children and a dog in a household. *J Clin Microbiol.* 2007;45:2014–6.
56. Zunino MG, De Francesco MV, Kuruc JA, Schweigmann N, Wisnivesky-Colli MC, Jensen O. Contaminación por helmintos en espacios públicos de la provincia de Chubut, Argentina. *Bol Chil Parasitol.* 2000;55:78–83.