



BRIEF REPORT

Atypical hand, foot, and mouth disease caused by *Coxsackievirus A6* in Argentina in 2015



Daniel M. Cisterna^{a,*}, Cristina L. Lema^a, Leila M. Martinez^a, Elizabeth Verón^b, Lorena P. Contarino^b, Denise Acosta^c, María Cecilia Freire^a

^a Servicio de Neurovirosis, INEI-ANLIS Dr. Carlos G. Malbran, Ciudad Autónoma de Buenos Aires, Argentina

^b Hospital Rural, Gan Gan, Chubut, Argentina

^c Hospital Zonal "Dr. Andres Isola", Puerto Madryn, Chubut, Argentina

Received 1 August 2017; accepted 28 May 2018

Available online 27 September 2018

KEYWORDS

Atypical hand, foot,
and mouth disease;
Coxsackievirus A6;
Enterovirus

PALABRAS CLAVE

Enfermedad de
mano-pie-boca
atípica;
Virus Coxsackie A6;
Enterovirus

Abstract We present two groups of cases of atypical hand, foot, and mouth disease (HFMD) caused by *Coxsackievirus A6* (CV-A6) detected in Argentina in 2015. The first group involved 14 patients from Chubut province and the second group affected 12 patients from San Luis province. Molecular analysis of the complete VP1 protein gene revealed the circulation of E2 sublineage, the most predominant worldwide. To our knowledge, this is the first report of CV-A6 infections associated with atypical HFMD in Argentina and South America.

© 2018 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Enfermedad de mano-pie-boca atípica causada por virus Coxsackie A6 en Argentina, 2015

Resumen Se describen dos grupos de casos de enfermedad de mano-pie-boca (HFMD) atípica causada por el virus Coxsackie A6 (*Coxsackievirus A6*, CV-A6) detectados en Argentina en el año 2015. El primero de los grupos involucró a 14 pacientes de Chubut y el segundo a 12 pacientes de San Luis. El análisis molecular del gen de la proteína VP1 completa reveló la circulación del sublinaje E2, el predominante a nivel global. Hasta donde sabemos, este es el primer reporte de infecciones CV-A6 asociadas con HFMD atípica en Argentina y Sudamérica.

© 2018 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: dcisterna@anlis.gov.ar (D.M. Cisterna).

Classical hand, foot, and mouth disease (HFMD) is an acute febrile infection characterized by vesicular exanthema on the palms or soles and oral mucosa, typically occurring in young children and infants. More than 90% of HFMD cases are caused by *Enterovirus A71* (EV-A71) and *Coxsackievirus A16* (CV-A16), both members of *Enterovirus A* species (EV-A), *Picornaviridae* family. However, sporadic

cases of the atypical form of HFMD can also be caused by other EV-A viruses¹⁹. We present two groups of atypical HFMD cases caused by *Coxsackievirus A6* (CV-A6) detected in Argentina in 2015.

The first group of 14 cases was reported in Chacabuco (a rural population of 89 inhabitants in Chubut province) between September 2nd 2015 and October 1st 2015. All




Figure 1 Phylogenetic tree of the complete VP1 gene of Argentinean *Coxsackievirus A6* strains from atypical hand, foot and mouth disease, 2015. For lineage classification, other selected strains from GenBank were included (accession numbers are given in the phylogenetic tree). The maximum likelihood method was used to construct the tree using MEGA version 6.0 (<https://www.megasoftware.net>). The phylogenetic tree was determined for 1000 replicates with random seeds. Only strong bootstrap values (>70%) were shown; (black dot) indicates Argentinean strains (GenBank accession No. KX575862-65).

Table 1 Detection and typing of human enterovirus serotypes from Argentina

	Acute Flaccid Paralysis Program ⁶	Aseptic Meningitis Cases ¹	Environmental surveillance ⁷
Period	2000–2015	1998–2014	2005–2015
Samples types	Stool	CSF	Water samples from Matanza-Riachuelo river
No. of samples	2480	4964	112
No. HEV-positive	188 ^a	1884 ^b	102 ^b
No. HEV-typed	152	326	60
No. HEV-A Species	23/152	2/326	2/60
HEV serotypes ^c	E-11	E-30	E-19
	E-13	E-4	CV-B3
	CV-B5	E-6	E-11
	CV-B4, E-14, E-18	CV-B1	PV-2, PV-3
	CV-A16	CV-B5	CV-B5

E: *Echovirus*, CV: *Coxsackievirus*, PV: *Poliovirus*.

^a Cell culture.

^b Nested RT-PCR.

^c The five most frequently identified serotypes.

patients were treated at the Gan Gan Rural Hospital, Chubut. The Department of Epidemiology of the province of Chubut reported the cases. Patients presented with fever with 48 h of evolution and the characteristic vesiculobulbous and erosive eruption, which primarily affected palms and soles. To a lesser extent, the rash affected the folds of the large joints, cheeks, perianal region and in some cases the whole body. Several patients presented onychomadesis on the fingernails. The second group cases occurred in General San Martín, San Luis province, a town with a population of 697 inhabitants. Twelve cases of atypical HFMD were reported between November 11th 2015 and November 22nd 2015 through the National System of Epidemiological Surveillance of measles and rubella. Initially, the patients were screened for these viral agents, being all the patients negative.

A total of eight stool samples from the 14 patients in Chubut and six from the 12 patients in San Luis were obtained. Viral RNA was extracted from stool suspensions using the QIAamp Viral RNA Mini Kit (Qiagen, Hilden, Germany). After that, a generic PCR for enterovirus detection was performed as previously described⁴. For the positive enterovirus cases, we used primer pairs 222/224 and 88/89 for partial VP1 sequencing for the identification of the enterovirus type. We subjected the sequences to BLAST search (<https://www.ncbi.nlm.nih.gov/BLAST>) and defined the virus type using the criteria of $\geq 75\%$ nucleotide and $\geq 85\%$ amino acid similarity in the VP1 region¹⁵. A complete VP1 gene of representative strains of both groups was obtained using nested primers specific for CV-A6 as previously described¹⁸. Moreover, the isolation of human enteroviruses in the human rhabdomyosarcoma (Rd) cell line was attempted for all specimens¹⁷.

Enteroviruses were detected in five (62.5%) of eight patients from Chubut and in four (66.7%) of six patients from San Luis. CV-A6 was identified in all enterovirus-positive samples. No virus was isolated in any of the samples tested. Partial sequencing of VP1 (position 2560–2865, based on strain Gdula, GenBank accession No. AY421764) of the

nine strains detected showed a high nucleotide similarity between them (99.1–99.7%), which suggests that both groups of cases are related. Phylogenetic analysis of the complete VP1 gene shows that CV-A6 strains are grouped into five lineages (A–E), with two sublineages E1 and E2². In our study, it was possible to obtain complete VP1 sequences of four strains (two from each cluster), which revealed that they belong to the E2 lineage (Fig. 1). This information is consistent with the latest notifications that showed that strains of the sublineage E2 are predominant worldwide^{2,10}.

HFMD outbreaks caused by CV-A6 have been reported in Singapore (2009), Taiwan (2009–2010), Japan (2011), Thailand (2012) and China (2013). In North America, CV-A6-associated HFMD outbreaks were reported in the USA and Cuba from 2011 to 2013². In Europe, CV-A6 was responsible for HFMD outbreaks in Finland, France and Spain from 2009 to 2011^{3,13} and more recently in Edinburgh, United Kingdom, in 2011–2013¹⁸. Argentinean CV-A6 strains show 95.5% similarity to strains detected in Scotland, suggesting a European source.

Classical HFMD has the highest incidence in infants and children. However, during the outbreak of atypical HFMD in Finland in 2008, a high incidence in adults was notified¹⁶. In Europe and North America, no significant differences in age were found among the CV-A6 cases. In our patients, the age range in Chacay Oeste and General San Martín was 2–28 years (average = 10.1) and 1–58 years (average = 12.2) respectively. We observed a wide age distribution although it would be necessary to study a larger population to estimate the real impact of this virus infection.

In this study, it was not possible to investigate material obtained from vesicular lesions and only stool samples were available. No virus was isolated in any of the samples tested possibly because Coxsackie A viruses are difficult to recover in cell culture systems¹¹. In Argentina, there is no active public health surveillance for HFMD and it is not a notifiable disease. Studies conducted in patients with meningitis, sepsis and in environmental samples in our country highlight the predominant role of *Enterovirus B* species (EV-B),

mainly represented by *Echovirus 30* (E-30), E-9, E-14, E-6, E-16, CV-A9 and CV-B1-3^{8,9,14}. Epidemiological information generated in our laboratory confirms that the circulation of EV-A is poorly detected (Table 1). Consequently, CV-A6 infections are probably underdiagnosed as a result of the lack of active public health surveillance for HFMD. Finally, the understanding of the variability of this atypical form of HFMD should help to avoid confusion with other skin conditions such as *eczema herpeticum* or chickenpox^{5,12}. In patients with non-specific clinical manifestations, it is important to rule out the presence of enteroviruses, which would avoid unnecessary treatment with acyclovir and possible hospitalization.

To our knowledge, this is the first report of CV-A6 infections associated with atypical HFMD in Argentina and South America. This information could be useful to establish an adequate surveillance system for enteroviral disease to implement more appropriate public health interventions.

Funding

This work was supported by INEI-ANLIS “Dr. Carlos G. Malbrán”, Buenos Aires, Argentina.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Acknowledgments

We thank to Gabriela Elbert, DiNaCEI, Ministerio de Salud de la Nación for collecting cases from San Luis province.

References

- Bonaventura R, Lema C, Martinez L, Jurado V, Girard D, Cisterna D, Freire MC. Enterovirus asociados a enfermedades neurológicas en Argentina durante el período 2005–2010. IX Congreso Argentino de Virología, Resumen 172. Rev Argent Microbiol. 2011;42 Suppl. 1:94.
- Brian L, Wang Y, Yao X, Mao Q, Xu M, Liang Z. *Coxsackievirus A6*: a new emerging pathogen causing hand, foot and mouth disease outbreaks worldwide. Expert Rev Anti Infect Ther. 2015;13:1061–71.
- Cabrero M, De Miguel T, Armada A, Martinez-Risco R, Pousa A, Trallero G. Onychomadesis after a hand, foot, and mouth disease outbreak in Spain, 2009. Epidemiol Infect. 2010;138:1775–8.
- Casas I, Klapper PE, Cleator GM, Echevarria JE, Tenorio A, Echevarria JM. Two different PCR assays to detect enteroviral RNA in CSF samples from patients with acute aseptic meningitis. J Med Virol. 1995;47:378–85.
- Chung WH, Shih SR, Chang CF, Lin TY, Huang YC, Chang SC, Liu MT, Ko YS, Deng MC, Liao YL, Lin LH, Chen TH, Yang CH, Ho HC, Lin JW, Lu CW, Lu CF, Hung SI. Clinicopathologic analysis of *Coxsackievirus A6* new variant induced widespread mucocutaneous bullous reactions mimicking severe cutaneous adverse reactions. J Infect Dis. 2013;208:1968–78.
- Cisterna DM, Palacios G, Rivero K, Girard D, Lema C, Freire MC. Epidemiology of enterovirus associated with neurologic diseases. Medicina (B Aires). 2007;67:113–9.
- Cisterna D, Lema C, Riviello Lopez G, Martinez L, Torres C, Mbayed V, Freire MC. Detección y caracterización de enterovirus y hepatitis A en el Riachuelo y Río de la Plata. IX Congreso Argentino de Virología Resumen 22999. Rev Argent Microbiol. 2008;40 Suppl. 1:49.
- Farias A, Cabrerizo M, Re V, Glatstein N, Pisano B, Spinsanti L, Contigiani MS. Molecular identification of human enteroviruses in children with neurological infections from the central region of Argentina. Arch Virol. 2011;156:129–33.
- Farias AA, Mojsiejczuk LN, Pisano MB, Flores FS, Aguilar JJ, Jean A, Yanes AN, Masachessi LA, Prez G, Isa VE, Campos MB, Re RH, Nates VESV. Environmental surveillance of Enteroviruses in Central Argentina: first detection and evolutionary analyses of E14. Food Environ Virol. 2018;10:121–6.
- Ganorkar NN, Patil PR, Tikute SS, Gopalakrishna V. Genetic characterization of enterovirus strains identified in Hand, Foot and Mouth Disease (HFMD): emergence of B1c, C1 subgenotypes, E2 sublineage of CV-A16 EV-71 and CV-A6 strains in India. Infect Genet Evol. 2017;54:192–9.
- Lipson SM, Walderman R, Costello P, Szabo K. Sensitivity of rhabdomyosarcoma and guinea pig embryo cell cultures to field isolates of difficult-to-cultivate group A coxsackieviruses. J Clin Microbiol. 1988;26:1298–303.
- Mathes EF, Oza V, Frieden IJ, Cordoro KM, Yagi S, Howard R, Kristal L, Ginocchio CC, Schaffer J, Maguiness S, Bayliss S, Lara-Corrales I, Garcia-Romero MT, Kelly D, Salas M, Oberste MS, Nix WA, Glaser C, Antaya R. “Eczema coxsackium” and unusual cutaneous findings in an enterovirus outbreak. Pediatrics. 2013;132:e149–57.
- Mirand A, Henquell C, Archimbaud C, Ughetto S, Antona D, Bailly JL, Peigue-Lafeuille H. Outbreak of hand, foot and mouth disease/herpangina associated with coxsackievirus A6 and A10 infections in 2010, France: a large citywide, prospective observational study. Clin Microbiol Infect. 2012;18:E110–8.
- Mistchenko AS, Viegas M, Latta MP, Barrero PR. Molecular and epidemiologic analysis of enterovirus B neurological infection in Argentine children. J Clin Virol. 2006;37:293–9.
- Oberste MS, Maher K, Kilpatrick DR, Pallansch MA. Molecular evolution of the human enteroviruses: correlation of serotype with VP1 sequence and application to picornavirus classification. J Virol. 1999;73:1941–8.
- Osterback R, Vuorinen T, Linna M, Susi P, Hyypia T, Waris M. *Coxsackievirus A6* and hand, foot, and mouth disease, Finland. Emerg Infect Dis. 2009;15:1485–8.
- Shoja ZO, Tabatabaie H, Shahmahmoudi S, Nategh R. Comparison of cell culture with RT-PCR for enterovirus detection in stool specimens from patients with acute flaccid paralysis. J Clin Lab Anal. 2007;21:232–6.
- Sinclair C, Gaunt E, Simmonds P, Broomfield D, Nwafor N, Wellington L, Templeton K, Willocks L, Schofield O, Harvala H. Atypical hand, foot, and mouth disease associated with *Coxsackievirus A6* infection Edinburgh, United Kingdom, January to February 2014. Euro Surveill. 2014;19:20745.
- Xu M, Su L, Cao L, Zhong H, Dong N, Dong Z, Xu J. Genotypes of the Enterovirus causing Hand Foot and Mouth Disease in Shanghai China, 2012–2013. PLoS ONE. 2015;10.