




ORIGINAL ARTICLE

Endophytic *Fusarium* isolates from *Ceratozamia mirandae* enhance tomato growth, suppress pathogenic fungi, and induce protection against *Botrytis cinerea*



Nidia del C. Ríos-De León^a, Miguel Á. Salas-Marina ^{a,*}, Vidal Hernández-García^a, Brenda del R. Saldaña-Morales^a, Luis A. Rodríguez-Larramendi^a, Rubén Martínez-Camilo^a, Claudio Ríos-Velasco^b, Daniel A. Pérez-Corral^b, Sergio Casas-Flores^c

^a Laboratorio de Biofertilizantes y Bioinsecticidas, Universidad de Ciencias y Artes de Chiapas, Facultad de Ingeniería, Sede Villa Corzo, Km 3.0 Carretera Villa Corzo, Ejido Monterrey, C.P. 30520, Chiapas, Mexico

^b Centro de Investigación en Alimentación y Desarrollo, Av. Río Conchos, Parque Industrial C.P. 31500, Ciudad Cuauhtémoc, Chihuahua, Mexico

^c IPICYT, División de Biología Molecular, Laboratorio de Genómica Funcional y Comparativa, Camino a la Presa San José 2055, Col. Lomas 4 Sección, 78216 San Luis Potosí, SLP, Mexico

Available online 6 September 2025

KEYWORDS

Arabidopsis;
Biofertilizer;
Colletotrichum;
Neopestalotiopsis;
Solanum lycopersicum

Abstract Fungal diseases in agricultural crops cause economic losses, with chemical control being the conventional method to manage them. However, this approach negatively impacts both the environment and human health. This study focused on endophytic fungi isolated from the roots of *Ceratozamia mirandae* in the Mexican locality of Juan Sabines (Villa Corzo, Chiapas). These fungi were identified morphologically and molecularly, biochemically characterized, and evaluated for their antagonistic activity against *Colletotrichum karstii*, *Neopestalotiopsis* sp. and *Fusarium oxysporum*. Their potential for promoting growth in *Arabidopsis thaliana* and tomato, as well as protecting against *Botrytis cinerea*, was also assessed. Fourteen fungal isolates were identified and grouped into six genera: *Fusarium*, *Pestalotiopsis*, *Trichoderma*, *Umbelopsis*, *Nectria* and *Podospira*. Among these *Fusarium proliferatum* JS311 and *F. oxysporum* JS239 exhibited strong inhibitory effects against the tested pathogens. Eight isolates were found to produce indole-3-acetic acid, promoting the growth of *A. thaliana* and tomato plants. Notably, *F. oxysporum* JS439 and *F. solani* (JS240, JS256 and JS4101) exhibited additional capabilities, including siderophore production and growth in nitrogen-free media. All *Fusarium* endophytic isolates induced systemic resistance against *B. cinerea* in tomato. Endophytic fungi

* Corresponding author.

E-mail address: miguel.salas@unicach.mx (M.Á. Salas-Marina).

<https://doi.org/10.1016/j.ram.2025.07.004>

0325-7541/© 2025 The Author(s). Published by Elsevier España, S.L.U. on behalf of Asociación Argentina de Microbiología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

from *C. mirandae* show promising potential as biofertilizers due to their combined mechanisms of anti-phytopathogenic activity, plant growth promotion, and systemic resistance induction through the production of beneficial metabolites.

© 2025 The Author(s). Published by Elsevier España, S.L.U. on behalf of Asociación Argentina de Microbiología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Arabidopsis;
Biofertilizante;
Colletotrichum;
Neopestalotiopsis;
Solanum
lycopersicum

Aislados endófitos obtenidos de *Fusarium* de *Ceratozamia mirandae* promueven el crecimiento de tomate, inhiben hongos patógenos e inducen protección contra *Botrytis cinerea*

Resumen Las enfermedades fúngicas causan pérdidas económicas en los cultivos agrícolas, y el control químico es el método principal para reducirlas. Sin embargo, este enfoque repercute negativamente tanto en el ambiente como en la salud humana. Este estudio se centró en hongos endófitos aislados de las raíces de *Ceratozamia mirandae* de muestras obtenidas en la localidad mexicana de Juan Sabines (Villa Corzo, Chiapas). Los hongos aislados se identificaron morfológicamente y molecularmente, se caracterizaron bioquímicamente y se evaluó su actividad antagonista frente a *Colletotrichum karstii*, *Neopestalotiopsis* sp. y *Fusarium oxysporum*. También se estudió su potencial para promover el crecimiento de *Arabidopsis thaliana* y tomate, así como el efecto protector frente a *Botrytis cinerea*. Se identificaron catorce hongos agrupados en seis géneros: *Fusarium*, *Pestalotiopsis*, *Trichoderma*, *Umbelopsis*, *Nectria* y *Podospora*. Entre ellos, *Fusarium proliferatum* JS311 y *F. oxysporum* JS239 mostraron fuertes efectos inhibidores contra los patógenos evaluados. Ocho aislados produjeron ácido indol-3-acético y promovieron el crecimiento de *A. thaliana* y tomate. En particular, *F. oxysporum* JS439 y *F. solani* JS240, JS256 y JS4101 mostraron capacidades adicionales, incluyendo la producción de sideróforos y el crecimiento en medios libre de nitrógeno. Todos los aislados de *Fusarium* indujeron resistencia sistémica contra *B. cinerea* en tomate. Los hongos endófitos de raíces de *C. mirandae* presentan un potencial prometedor como biofertilizantes debido a sus mecanismos combinados de actividad antifitopatogénica, promoción del crecimiento vegetal e inducción de resistencia sistémica mediante la producción de metabolitos benéficos.

© 2025 Los Autores. Publicado por Elsevier España, S.L.U. en nombre de Asociación Argentina de Microbiología. Este es un artículo Open Access bajo la CC BY-NC-ND licencia (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Agricultural pests and diseases annually reduce global crop yields by 30–50%, with chemical control remaining the most widely used method in modern agriculture¹³. However, excessive reliance on chemical inputs has led to significant challenges, including the development of resistance in pathogens and pests, loss of biodiversity, soil degradation, harm to beneficial fauna, and adverse effects on human and animal health²².

The use of endophytic microorganisms (EMs) in biofertilizers has emerged as an effective and sustainable strategy to enhance agricultural productivity. EMs control diseases caused by pathogens and suppress pest damage, reduce dependence on chemical inputs, mitigate ecological impacts, and improve crop yield and quality¹⁵. Recent studies highlight the potential of EMs as biofertilizers in agriculture. For instance, Khan et al.¹⁷ demonstrated that *Acremonium* strains isolated from *Lilium davidii* exhibited antifungal activity, promoted the growth of *Allium tuberosum*, and produced plant growth-promoting substances such as 3-indolacetic acid (IAA), siderophores, organic acids

and solubilized phosphates. Similarly, EMs from the genera *Alternaria*, *Didymella*, *Fusarium* and *Xylogone* isolated from *Sophora flavescens* were shown to enhance primary root growth in *Arabidopsis thaliana* while producing IAA, siderophores and solubilized phosphates⁴³. Species of the genus *Trichoderma* have also been widely utilized as biological control agents due to their ability to confer resistance against plant pathogens and increase the productivity of economically important crops³². These findings underscore the promising role of EMs in sustainable agriculture, offering an eco-friendly alternative to conventional chemical-based practices.

EMs reside within plant tissues during part or all of their life cycle, offering their host plants a wide range of biological benefits¹⁵. The primary effects of EMs include enhanced stress tolerance by increasing resilience to abiotic stressors, such as acidity, salinity, heavy metals, drought, floods, extreme temperatures, as well as biotic stressors like pathogens, insects and herbivores⁴¹. EMs also promote plant growth through the production and secretion of plant hormones such as gibberellins and auxins, solubilizing essential nutrients in the soil including phosphate and iron, to

make them more available to plants^{11,13}, and facilitating nitrogen transfer to plants through strains that express ACC deaminase, which are up to 40% more efficient at forming nitrogen-fixing nodules¹². Additionally, EMs induce systemic defense mechanisms in plants by producing anti-fungal and antibiotic secondary metabolites that inhibit pathogen development and compete for space and nutrients, thereby protecting the plant^{13,15}.

The rhizosphere serves as a prolific source of rhizospheric or endophytic microorganisms; it is estimated that over one million EMs species, including bacteria and fungi, are associated with approximately 300 000 plant species¹⁰. However, studies suggest that the diversity of endophytic communities associated with various plant species remains significantly underestimated^{6,42}. For example, Ulloa-Muñoz et al.⁴⁵ isolated EM species from native and endemic plants of the Cordillera Blanca in the Peruvian Andes, demonstrating their potential to promote crop growth and control pathogen.

These findings suggest that wild plants endemic to conserved areas harbor a vast diversity of EMs. Therefore, this study aims to characterize endophytic fungi associated with the roots of *Ceratozamia mirandae* (Pérez, Vovides & Iglesias Amenduái)²⁹ (Cycadales: Zamiaceae), an endemic species of Chiapas, Mexico, focusing on their biocontrol potential against fungal phytopathogens, their ability to promote plant growth, and their capacity to induce systemic resistance in tomato seedlings against *Botrytis cinerea*.

Materials and methods

Study area

Samples of healthy secondary roots of *C. mirandae* were collected in the Natural Resource Protection Reserve "La Frailescana", located in the locality of "Juan Sabines" within the municipality of Villa Corzo, Chiapas, Mexico (16°03'20.5"N; 93°28'41.6"W, at an altitude of 1817 m above sea level).

Sampling and isolation of endophytic fungi

Healthy roots of *C. mirandae* were collected from a depth of 10 cm, carefully wrapped in kraft paper, and transported in polyethylene bags to the Laboratory of Biofertilizers and Bioinsecticides at the University of Sciences and Arts of Chiapas (UNICACH-Villa Corzo campus). The roots were cut into 10 mm explants using a scalpel, then surface-disinfected with 70% ethanol for 3 min, followed by 10% sodium hypochlorite for 10 min, and finally rinsed twice with sterile distilled water. The disinfected explants were placed on potato dextrose agar (PDA) medium and incubated at 28 °C for 48 h. Emerging fungal colonies were isolated and purified using the hyphal tip technique.

Morphological and molecular identification

Fungal isolates were first identified morphologically by observing their reproductive structures using an Axiolab Carl Zeiss® compound microscope and following dichotomous keys for fungal taxonomy⁴⁶. For

molecular identification, genomic DNA was extracted from the isolates according to the protocol described by Raeder and Broda³³. The internal transcribed spacer (ITS) region of 18s rDNA was amplified and sequenced using the ITS5 (5'-GGAAGTAAAAGTCGTAACAAGG-3') and ITS4 (5'-TCTCTCCGCTTATTGATATATGC-3') primers⁴⁷. Polymerase chain reaction (PCR) amplicons were sequenced using the Sanger method on an ABI sequencer (Applied Biosystem)¹. The resulting sequences were compared with the National Center for Biotechnology Information (NCBI) database using the BLAST algorithm and aligned using CLUSTAL W in MEGA7® software. GenBank accession numbers were obtained for the submitted sequences⁴².

In vitro dual confrontation bioassays

First, a screening bioassay was performed to evaluate the antagonistic activity of endophytic fungi (EF) using *Colletotrichum karstii* as a model; the isolates that exhibited notable antagonistic responses were used for further assays against other fungi. The antagonistic EF were evaluated against *C. karstii* and *Neopestalotiopsis* sp., foliar pathogens of camedor palm³⁸, and *Fusarium oxysporum* coffee (FoC), a coffee seedling pathogen donated by the Laboratory of Biofertilizers and Bioinsecticides of UNICACH-Villa Corzo. Pathogen explants were placed at one end of a Petri dish containing PDA, while EF explants were placed at the opposite end. The plates were incubated at 28 °C for 8 days (d). Mycelial growth inhibition (MGI) of the pathogen was calculated using the formula $[(R - r)/R] \times 100$, where *R* is the control pathogen growth and *r* is the pathogen growth in confrontation³⁴. A completely randomized design with four replicates per EF was used.

Characterization of fungi with plant growth-promoting activity

The plant growth-promoting activity of *C. mirandae* endophytic isolates were assessed on three-day-old *A. thaliana* Ecotype Col-0 seedlings. Each experiment included four replicates, with five seedlings per replicate. Seedlings were placed on the surface of Petri dishes containing Murashige and Skoog (MS) culture medium, with EF explants positioned at the bottom. Plates were maintained under a 16 h light/8 h dark photoperiod at 25 °C for 12 d and monitored daily until the roots contacted the fungal growth³⁶. Morphological changes in root architecture and aerial parts were considered positive indicators of growth promotion.

Biochemical characterization of fungi with plant growth-promoting activity

Detection of 3-indolacetic acid

IAA production was determined using the protocol by Ponce de León³¹. Five-day-old EF explants pre-cultured on PDA were incubated in test tubes with nutrient broth supplemented with L-tryptophan (150 g/l) under constant shaking (1800 rpm) at 28 ± 1 °C for 7 d. Supernatants from centrifuged cultures (13 000 g, 15 min) were mixed with Salkowski's reagent (100 ml of 35% perchloric acid and

2 ml of 0.5M ferric chloride) and incubated in the dark for 30 min. A color change from yellow to pink indicated IAA production, which was quantified at 530 nm using a Thermo Scientific® Evolution™ 300 UV-Vis spectrophotometer. Absorbance values were compared to an IAA calibration curve (0–0.1 mg/ml)³¹.

Detection of siderophores

EF were inoculated on plates containing solidified Cro-mozurol S (CAS) medium, as described by Loudon et al.²⁰. Plates were incubated at $28 \pm 1^\circ\text{C}$ for 7 d in darkness. Isolates displaying orange halos around the colonies were identified as siderophore producers.

Growth in nitrogen-free media assay

EF were grown on nitrogen-free Ashby culture medium (20 g sucrose, 0.2 g K_2HPO_4 , 0.2 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g NaCl, 0.1 g K_2SO_4 , 5.0 g CaCO_3 , and 15 g agar per liter). Plates were incubated at $28 \pm 1^\circ\text{C}$ for 7 d. EF that grew on this medium were considered positive for growth in nitrogen-free media⁷.

Phosphate solubilization assay

Phosphate solubilization was evaluated using NBTRIP medium (10 g glucose, 5 g $\text{Ca}_3(\text{PO}_4)_2$, 5 g $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, 0.25 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.20 g KCl, 0.1 g $(\text{NH}_4)_2\text{SO}_4$ and 15 g agar per liter). EF explants were inoculated and incubated in darkness at $28 \pm 1^\circ\text{C}$ for 7 d. Clear halos around the colonies indicated phosphate solubilization²⁴.

Tomato growth promotion bioassay

Black 96-cavity Hydroenvironment® germination trays filled with CosmoPeat® peat moss substrate were used. *Solanum lycopersicum* var. Pomodoro (USAgriSeeds®) seeds were sown at a 3 mm depth and subjected to the following treatments: T1 – absolute control (water), T2 – negative control with *F. oxysporum* tomato strain (FoT), T3 – positive control I (50% N, 1.63 kg/ha), T4 – positive control II (100% N, 3.23 kg/ha), T5–T12 – treatments with isolates JS286, JS240, JS414, JS439, JS4101, JS296, JS285, and JS256, respectively.

A non-growth-promoting pathogenic tomato strain of *F. oxysporum* (FoT) was used as a negative control. Seeds were inoculated with 20 μL of 1×10^6 conidia/ml of each EF isolate at sowing, applied directly to the seeds, followed by a second inoculation 5 d post-emergence, applied directly to the base of the stem using a micropipette.

Fertilization with Fertiquim® urea, based on a planting density of 41 666 plants/ha, was applied using a 50% nitrogen dose in plant-EF interactions (T2, T5–T12)²⁵ to evaluate the capacity of EF isolates to promote growth under nitrogen deficiency. Seedlings were maintained under semi-controlled conditions with 50% shade and irrigated to prevent leaching.

A completely randomized design with eight replicates per treatment was used. At 30 d post-planting (dpp), destructive analyses were performed to measure stem diameter (SD, mm), root length (RL, cm), seedling height (SH, cm), seedling fresh weight (SFW, mg), root fresh weight (RFW, mg), seedling dry weight (SDW, mg), root dry weight (RDW, mg), and leaf chlorophyll content ($\mu\text{mol}/\text{m}^2/\text{s}$).

Morphological traits (SD, RL, SH) were measured using an Adeske® electronic vernier caliper. SD was measured 3 cm

from the stem base; RL from the stem base to the main root tip; and SH from the stem base to the terminal bud. Biomass variables (SFW and RFW) were recorded by separately weighing the stem and roots using a Velab® analytical scale. Dry biomass variables (SDW and RDW) were determined after drying samples at 70°C for 72 h in a Thermo Scientific® oven. Chlorophyll content was measured using a Quantum Flux Apogee® sensor placed at the center of seedling leaves²⁵.

Bioassays of systemic protection against *B. cinerea* induced by endophytic fungi of *C. mirandae* on tomato plants

Thirty-day-old leaves of *S. lycopersicum* var. Pomodoro (USAgriSeeds) seedlings, produced in an independent experiment under the same conditions as described for the growth promotion bioassay, were inoculated with 20 μL of the necrotrophic fungus *B. cinerea* (a pathogenic tomato strain obtained from the Centro de Investigación en Alimentación y Desarrollo CIAD-Unidad Cuauhtémoc, Chihuahua) at a concentration of 1×10^6 conidia/ml. The conidia were resuspended in water containing Break-Thru® (BASF) at 0.1% and applied directly to the leaves with a micropipette³⁷. A completely randomized design with six replicates per treatment was employed, with three leaves inoculated per seedling. Seedlings were maintained in a greenhouse for 7 d at $26 \pm 2^\circ\text{C}$. Leaf lesions were measured using ImageJ software, calibrated with a scale for accuracy³⁹. The percentage of disease severity was calculated using the formula (% severity = damaged area/total area \times 100)²⁷.

Statistical analysis

Data were subjected to one-way analysis of variance (ANOVA), followed by Tukey's multiple comparison test ($p \leq 0.05$), using the MINITAB19® statistical software. Experiments were conducted in duplicate at separate times to ensure reproducibility.

Results

Identified endophytic fungi

The ITS sequences (ITS4–ITS5) of the EF showed similarities ranging from 94 to 100% in the GenBank database. These sequences were submitted with the National Center for Biotechnology Information (NCBI), where unique accession numbers were assigned (Table 1). A total of 14 EF isolates associated with *C. mirandae* were identified, representing the genera *Fusarium*, *Pestalotiopsis*, *Trichoderma*, *Umberlopsis*, *Nectria* and *Podospira*. Based on morphological and molecular analyses, these fungi were classified into three classes: Sordariomycetes, Euascomycetes and Zygomycetes (Table 1).

Endophytic *Fusarium* isolates inhibit pathogen growth

In the *in vitro* screening bioassay, *F. proliferatum* JS311 and *F. oxysporum* JS439 exhibited notable antagonistic responses

Table 1 Diversity of root endophytic fungi associated with *C. mirandae*: morphological and molecular identification using ITS marker sequencing and GenBank accession.

Endophytic fungal isolates	GenBank access number	Macro and microscopic identification	Related species in GenBank	Similarity (%)	Class
JS311	OR843338	<i>F. proliferatum</i>	<i>F. proliferatum</i>	99	Sordariomycetes
JS312	OR842899	<i>Pestalotiopsis</i> sp.	<i>Pestalotiopsis</i> sp.	99	Sordariomycetes
JS414	OR843339	<i>F. oxysporum</i>	<i>F. oxysporum</i>	100	Sordariomycetes
JS439	OR843341	<i>F. oxysporum</i>	<i>F. oxysporum</i>	99	Sordariomycetes
JS240	OR843340	<i>F. solani</i>	<i>F. solani</i>	99	Sordariomycetes
JS455	PP722765	<i>Trichoderma spirale</i>	<i>Trichoderma spirale</i>	98	Euascomycetes
JS256	OR852613	<i>F. solani</i>	<i>F. solani</i>	99	Sordariomycetes
JS469	OR843989	<i>Umbelopsis nana</i>	<i>Umbelopsis nana</i>	99	Zygomycetes
JS285	OR843990	<i>Nectria</i> sp.	<i>Nectria</i> sp.	99	Sordariomycetes
JS286	PP722977	<i>Fusarium</i> sp.	<i>Fusarium</i> sp.	100	Sordariomycetes
JS296	PP586144	<i>F. solani</i>	<i>F. solani</i>	94	Sordariomycetes
JS4101	PP724710	<i>F. solani</i>	<i>F. solani</i>	99	Sordariomycetes
JS3102	PP586187	<i>Podospira dennisiae</i>	<i>Podospira dennisiae</i>	99	Sordariomycetes
JS2107	PP724711	<i>F. solani</i>	<i>F. solani</i>	100	Sordariomycetes

Table 2 Radial growth inhibition of *C. karstii*, *Neopestalotiopsis* sp. and *F. oxysporum* coffee (FoC) by EF isolates *F. oxysporum* JS439 and *F. proliferatum* JS311.

Endophytic fungi	<i>C. karstii</i>	<i>Neopestalotiopsis</i> sp.	FoC
<i>F. oxysporum</i> (JS439)	40.62 ± 2.04 ^A	29.25 ± 0.62 ^B	20.93 ± 0.80 ^C
<i>F. proliferatum</i> (JS311)	39.13 ± 1.61 ^A	24.87 ± 1.82 ^B	−4.62 ± 0.98 ^C

Each result represents the mean ± standard deviations (SD) of four replicates for each pathogenic fungus. Different letters within rows indicate statistically significant differences (Tukey's test, $p \leq 0.05$).

against *C. karstii*. In the bioassay involving three pathogens, *F. oxysporum* JS439 inhibited the mycelial growth of *C. karstii* (40.62%), *Neopestalotiopsis* sp. (29.25%) and *F. oxysporum* (20.93%), respectively. In contrast, *F. proliferatum* JS311 inhibited the growth of *C. karstii* (39.13%) and *Neopestalotiopsis* sp. (24.87%) but showed no inhibition against *F. oxysporum* (Table 2).

Endophytic *Fusarium* isolates promote *A. thaliana* growth

Eight of the 14 EF isolates associated with *C. mirandae* demonstrated growth-promoting effects of *A. thaliana* seedlings. Isolates of *F. oxysporum* (JS414 and JS439), *F. solani* (JS240, JS256, JS296 and JS4101), *Nectria* sp. (JS285) and *Fusarium* sp. (JS286) significantly enhanced root architecture, root hair development, and aerial growth of *A. thaliana* seedlings compared to the control group (Table 3 and Fig. 1).

Endophytic *Fusarium* isolates produce auxins, siderophores and fix nitrogen

The eight EF isolates that promoted *A. thaliana* growth were found to produce IAA at varying concentrations (Table 3). Among these, *F. oxysporum* JS439, *F. solani* JS240, JS256 and JS4101 isolates demonstrated siderophore production,

with *F. oxysporum* JS439 producing the largest orange halo on CAS medium (Table 3). Six isolates, including *F. oxysporum* (JS414, JS439), *F. solani* (JS240, JS296, JS4101) and *Nectria* sp. (JS285), exhibited the ability to grow in nitrogen-free medium (Table 3). However, none of the tested isolates showed positive results for phosphate solubilization (Table 3).

Endophytic *Fusarium* isolates promote tomato growth

Tomato seedlings inoculated with the EF isolates of *C. mirandae*, *F. oxysporum* JS414, JS439; *F. solani* JS240, JS256, JS296, JS4101; *Nectria* sp. JS285 and *Fusarium* sp. JS286, exhibited significantly enhanced growth across several variables, including stem diameter (SD), root length (RL), seedling height (SH), seedling dry weight (SDW) and root dry weight (RDW), compared to the control treatments (T1, T2, T3, T4) (Table 4).

In the SD variable, seedlings inoculated with EF and supplemented with 50% nitrogen fertilization recorded higher values than the control groups, including the positive control II (T4, 100% nitrogen). The highest SD values were observed in seedlings treated with *Nectria* sp. JS285 and *F. solani* JS296 at 2.02 and 1.96 mm, respectively (Table 4).

For RL, the positive control T3 treatment (50% nitrogen) produced the longest roots at 93.42 mm. However, seedlings

Table 3 Endophytic fungi from *C. mirandae* roots: growth promoters of *A. thaliana*, producers of IAA and siderophores, nitrogen (N) fixers, and phosphate (P) solubilizers.

Isolate	Species	Growth promotion	IAA (mg/ml)	Siderophores	Fixing N	Solubilization P
JS311	<i>F. proliferatum</i>	—	—	—	—	—
JS312	<i>Pestalotiopsis</i> sp.	—	—	—	—	—
JS414	<i>F. oxysporum</i>	+	0.06 ± 0.04 ^{BC}	—	+	—
JS439	<i>F. oxysporum</i>	+	0.08 ± 0.11 ^{ABC}	+	+	—
JS240	<i>F. solani</i>	+	0.01 ± 0.00 ^D	+	+	—
JS455	<i>Trichoderma spirale</i>	—	—	—	—	—
JS256	<i>F. solani</i>	+	0.04 ± 0.05 ^{CD}	+	—	—
JS469	<i>Umbelopsis nana</i>	—	—	—	—	—
JS285	<i>Nectria</i> sp.	+	0.05 ± 0.03 ^{CD}	—	+	—
JS286	<i>Fusarium</i> sp.	+	0.12 ± 0.19 ^A	—	—	—
JS296	<i>F. solani</i>	+	0.01 ± 0.00 ^D	—	+	—
JS4101	<i>F. solani</i>	+	0.10 ± 0.04 ^{AB}	+	+	—
JS3102	<i>Podospora dennisiae</i>	—	—	—	—	—
JS2107	<i>F. solani</i>	—	—	—	—	—

Based on the qualitative analysis, results were categorized as positive (+) or negative (—). For the quantitative analysis of IAA, each result represents the mean of four replicates ± standard deviation (SD). Values with different letters indicate statistically significant differences according to Tukey's test ($p \leq 0.05$).




Figure 1 Schematic representation of growth promotion of *A. thaliana* by endophytic fungi of *C. mirandae*. (a) Control, (b) *F. solani* (JS240) and (c) *F. solani* (JS296).

inoculated with *F. solani* JS4101 plus 50% nitrogen and *Nectria* sp. JS285 plus 50% nitrogen recorded 92.65 mm and 91.65 mm, respectively, surpassing the control treatments T1, T2 and T4 (Table 4).

In the SH variable, EF-inoculated treatments promoted seedlings height compared to controls (T1, T2 and T3). The highest SH values were recorded in positive control II T4 (100% nitrogen control, 72.88 mm) and *F. solani* JS256 plus 50% nitrogen (72.77 mm), followed by *Nectria* sp. JS285 at 69.25 mm (Table 4).

For seedling fresh weight (SFW) and root fresh weight (RFW), EF-inoculated seedlings with 50% nitrogen displayed comparable values to the 100% nitrogen control (T4), and outperformed treatments T1, T2 and T3. The highest SFW (0.53 mg) and RFW (0.37 mg) were achieved with *F. solani* JS256 plus 50% nitrogen (Table 4).

In the SDW and RDW variables, EF-inoculated seedlings with 50% nitrogen fertilization showed equivalent results to the 100% nitrogen positive control II (T4) and outperformed T1, T2 and T3. The highest SDW values (0.05 mg) were observed in seedlings treated with *F. solani* JS256 plus 50% N and *Fusarium* sp. JS286 plus 50% N. For RDW, *F. solani*

JS256 presented the highest value (0.05 mg), exceeding all control treatments (Table 4).

Chlorophyll concentration in tomato leaves was highest in seedlings treated with *Nectria* sp. JS285 and *F. solani* JS296, with values of 224.60 $\mu\text{mol}/\text{m}^2/\text{s}$ and 221.76 $\mu\text{mol}/\text{m}^2/\text{s}$, respectively. In comparison, the absolute water control treatment (T1) recorded a chlorophyll concentration of 192.21 $\mu\text{mol}/\text{m}^2/\text{s}$ (Table 4).

Colonization of tomato roots by endophytic *Fusarium* isolates induces resistance against *B. cinerea*

Seven days after the foliar plant–pathogen interaction, diverse levels of severity caused by *B. cinerea* were observed. Tomato seedling leaves treated with different endophytic isolates of *C. mirandae* showed significant lower severity percentages compared to the control groups (negative control (T2), positive control I (T3) and positive control II (T4)). Among the treatments, *Nectria* sp. JS285 demonstrated the highest systemic resistance, with a severity value of 2.89%, which was statistically comparable to the abso-

Table 4 Impact of endophytic *Fusarium* isolates on tomato seedling growth.

Treatments	SD (mm)	RL (mm)	SH (mm)	SFW (mg)	SFW (mg)	SDW (mg)	RDW (mg)	Cl ($\mu\text{mol}/\text{m}^2/\text{s}$)
Control	1.61 \pm 0.06 ^E	89.88 \pm 1.32 ^{BC}	64.10 \pm 0.58 ^C	0.33 \pm 0.00 ^D	0.11 \pm 0.00 ^G	0.03 \pm 0.00 ^{EF}	0.01 \pm 0.00 ^{EF}	192.21 \pm 1.56 ^{EF}
FoT	1.62 \pm 0.08 ^E	75.30 \pm 1.59 ^{GH}	64.11 \pm 1.20 ^C	0.36 \pm 0.04 ^{CD}	0.12 \pm 0.01 ^G	0.03 \pm 0.00 ^F	0.01 \pm 0.00 ^F	203.89 \pm 1.67 ^D
N 50%	1.43 \pm 0.05 ^F	93.42 \pm 1.17 ^A	61.65 \pm 0.70 ^D	0.37 \pm 0.02 ^C	0.16 \pm 0.01 ^F	0.04 \pm 0.00 ^{BCDE}	0.01 \pm 0.00 ^{CD}	187.95 \pm 3.02 ^F
N 100%	1.83 \pm 0.05 ^{BC}	77.57 \pm 1.55 ^{FG}	72.88 \pm 0.28 ^A	0.48 \pm 0.01 ^B	0.15 \pm 0.01 ^F	0.04 \pm 0.00 ^{ABC}	0.02 \pm 0.00 ^{CD}	208.95 \pm 1.79 ^{BC}
<i>Fusarium</i> sp. (JS286)	1.86 \pm 0.05 ^{BC}	81.00 \pm 3.46 ^E	65.55 \pm 1.06 ^C	0.46 \pm 0.02 ^B	0.23 \pm 0.03 ^{CD}	0.05 \pm 0.00 ^A	0.02 \pm 0.00 ^B	211.79 \pm 3.63 ^B
<i>F. solani</i> (JS240)	1.78 \pm 0.06 ^{CD}	72.77 \pm 1.80 ^H	64.21 \pm 0.90 ^C	0.47 \pm 0.02 ^B	0.23 \pm 0.02 ^{CD}	0.05 \pm 0.00 ^{AB}	0.02 \pm 0.00 ^{CD}	189.39 \pm 3.96 ^{EF}
<i>F. oxysporum</i> (JS414)	1.68 \pm 0.06 ^{DE}	79.28 \pm 1.53 ^{EF}	59.30 \pm 0.35 ^E	0.37 \pm 0.01 ^C	0.18 \pm 0.01 ^{EF}	0.03 \pm 0.00 ^{DEF}	0.01 \pm 0.00 ^{EF}	206.13 \pm 2.07 ^{CD}
<i>F. oxysporum</i> (JS439)	1.88 \pm 0.08 ^{BC}	87.30 \pm 1.31 ^{CD}	61.86 \pm 1.58 ^D	0.45 \pm 0.01 ^B	0.20 \pm 0.02 ^{DE}	0.04 \pm 0.00 ^{CDEF}	0.02 \pm 0.00 ^C	193.07 \pm 2.29 ^E
<i>F. solani</i> (JS4101)	1.81 \pm 0.08 ^{CD}	92.61 \pm 1.35 ^{AB}	62.05 \pm 0.81 ^D	0.44 \pm 0.01 ^B	0.23 \pm 0.01 ^{CD}	0.04 \pm 0.00 ^{ABCD}	0.01 \pm 0.00 ^{DE}	211.53 \pm 1.28 ^B
<i>F. solani</i> (JS296)	1.96 \pm 0.05 ^{AB}	87.81 \pm 1.60 ^{CD}	65.18 \pm 0.80 ^C	0.48 \pm 0.01 ^B	0.25 \pm 0.00 ^C	0.04 \pm 0.00 ^{ABC}	0.01 \pm 0.00 ^{CD}	221.76 \pm 1.46 ^A
<i>Nectria</i> sp. (JS285)	2.02 \pm 0.08 ^A	91.65 \pm 1.35 ^{AB}	69.25 \pm 0.88 ^B	0.47 \pm 0.00 ^B	0.33 \pm 0.01 ^B	0.04 \pm 0.00 ^{ABC}	0.02 \pm 0.00 ^B	224.60 \pm 1.94 ^A
<i>F. solani</i> (JS256)	1.86 \pm 0.13 ^{BC}	85.85 \pm 0.84 ^D	72.77 \pm 2.75 ^A	0.53 \pm 0.01 ^A	0.37 \pm 0.01 ^A	0.05 \pm 0.00 ^A	0.05 \pm 0.00 ^A	210.26 \pm 2.23 ^{BC}

SD: stem diameter; RL: root length; SH: seedling height; SFW: seedling fresh weight; RFW: root fresh weight; SDW: seedling dry weight; RDW: root dry weight; Cl: chlorophyll content. *F. oxysporum* tomato (FoT), nitrogen (N). Each result represents the mean of eight seedlings \pm SD. Values within columns followed by the same letter are not significantly different according to Tukey's test ($p \leq 0.05$).




Figure 2 Systemic protection induced by endophytic fungi of *C. mirandae* in tomato seedlings against the necrotrophic pathogen *B. cinerea*. The bars in the graph represent the percentage of severity caused by *B. cinerea*. Each result is the mean of six replicates \pm SD. Values sharing the same letter are not significantly different according to Tukey's test ($p \leq 0.05$).

lute water control treatment (T1), with a severity of 0.07% (Fig. 2). These findings suggest that the inoculation of endophytic *Fusarium* isolates can effectively induce systemic resistance in tomato plants against *B. cinerea*.

Discussion

Fusarium was the most abundant genus (eight isolates) among the 14 EF isolated from *C. mirandae* roots. *Fusarium* is one of the three most dominant genera worldwide, with more than 1500 rhizosphere-associated species described⁴⁸. Its strong ability to colonize tissues of a wide variety of host plants has also been documented²⁸. Some strains are plant pathogens, while others can survive as saprophytes⁵. Due to its phytopathogenic nature, it is ranked among the ten most devastating fungal genera globally⁸. However, in recent years, non-pathogenic endophytic *Fusarium* strains have been reported to play beneficial roles in agriculture and forestry, owing to their ability to colonize root surfaces and protect even susceptible plant varieties from the highly virulent pathogenic isolates²⁸.

Endophytic *Fusarium* isolates from *C. mirandae* showed *in vitro* antagonism against *C. karstii*, *Neopestalotiopsis* sp. and FoC. These results align with those reported by Hamzah et al.¹⁴, where *Fusarium* strains, such as mangrove endophytes, inhibited the growth of the pathogen *F. solani* 45–66%. It has been reported that non-pathogenic *Fusarium* isolates produce compounds such as 5-hexenoic acid, limonene, octanoic acid, 3,4-dihydropyran, saponins, phenols, flavonoids, tannins, alkaloids, anthroquinones, and terpenoids, some of which are toxic to fungal and bacterial pathogens²⁸.

Additionally, *Fusarium* isolates produced the highest amount of IAA (0.12 mg/ml). Previous studies have reported that endophytic *Fusarium* isolates can produce up to 0.42 mg/ml IAA.²¹ Other EF have been reported

to have different amounts of IAA, such as *Acremonium* (53.12–167.71 mg/ml)¹⁷, *Didymella*, *Alternaria* and *Fusarium*, which produced IAA levels of 0.5–16.0 mg/ml⁴⁴. Therefore, the amount of IAA produced by a fungus depends on the species and the nutritional and environmental conditions to which it is exposed^{13,28}.

In this study, endophytic *Fusarium* isolates of *C. mirandae* produced siderophores; siderophore-producing fungi are generally considered to promote plant growth¹⁷. Six of the 14 EF isolates were able to grow in nitrogen-free medium, likely due to their association with diazotrophic microorganisms capable of nitrogen fixation, endohyphal bacteria⁴⁰, or the production and secretion of enzymes such as ACC deaminase, which facilitate nitrogen fixation and transfer to plants¹⁷. However, EF isolates of *C. mirandae* were unable to solubilize phosphorus (P). In this regard, the primary EF isolates reported as P solubilizers belong to the genera *Penicillium*, *Aspergillus*, *Piriformospora*, *Curvularia*, and arbuscular mycorrhizal fungi²³. Nevertheless, endophytic *Fusarium* isolates capable of phosphate solubilization have been documented⁹. Therefore, the antagonistic and growth-promoting effects of beneficial *Fusarium* strains in plants are probably due to the production and secretion of various metabolites.

EF of *C. mirandae* positively modified root and shoot architecture in *A. thaliana* seedlings. It also significantly increased growth parameters (SD, RL, SH, SFW, RFW, SDW and RDW) in tomato seedlings. The use of endophytic *Fusarium* isolates has been documented in various agriculturally important crops. Cheng et al.⁴ reported that native *Fusarium* strains, along with their endobacterium *Klebsiella aerogenes*, promoted tomato seedling growth by increasing RL and SH compared to control groups. Similarly, Fauriah et al.⁹ demonstrated that endophytic *Fusarium* strains enhanced RL and SH in maize compared to control seedlings. Kisaakye et al.¹⁸ showed that non-pathogenic *Fusarium* strains improved banana yields during the first cycle, increasing SH and pseudostem circumference while reducing *Radopholus similis* nematode densities. Likewise, Moročko-Bičevska et al.²⁶ demonstrated that endophytic *Fusarium* strains promoted strawberry growth, with significantly higher RFW and shoots weights compared to control plants. They also documented that pre-inoculating soil with *Fusarium* strains significantly reduced disease severity caused by *Gnomonia fragariae*. Glick¹² showed that crude extracts from endophytic *Fusarium* strains of *Biserula pelecinus* increased SDW and RDW values in *Lolium multiflorum* (a forage plant) seedlings compared to controls. Moreover, tomato plants inoculated with *Fusarium* endophytes and fertilized with 50% nitrogen showed growth variables comparable to the positive control II T4 (100%). This suggests that the 50% nitrogen deficiency may be compensated by the production of various metabolites or through interactions between endophytes and other microorganisms²⁸.

EF of *C. mirandae* significantly increased chlorophyll content in tomato seedlings. Hatamzadeh et al.¹⁵ reported that corn seedlings treated with *Alternaria consortiale*, an endophyte of *Anthemis altissima*, showed up to a 60% increase in chlorophyll levels compared to untreated seedlings. Higher chlorophyll concentrations enhance light energy uptake, thereby maximizing photosynthesis and

accelerating plant growth³⁵. It is important to note that although isolated endophytes of *Fusarium* have been shown to promote the growth of *Arabidopsis* and tomato seedlings, further studies are needed. This is particularly relevant given that some *Fusarium* species are pathogenic and produce mycotoxins during plant interactions, which can negatively affect plants and pose risks to human and animal health³⁰.

EF of *C. mirandae* also induced systemic protection in tomato seedlings, reducing disease severity caused by necrotrophic pathogen *B. cinerea*. Comparable results were observed with endophytic strains of *Beauveria* spp., which induced resistance in chili bell pepper and tomato plants against *B. cinerea*³. Additionally, *Trichoderma asperellum* has been reported to induce systemic protection against *B. cinerea* in tomato plants¹⁶. The confrontation assays in this study demonstrated the antifungal potential of EF from *C. mirandae* against *B. cinerea*. This systemic defense induction may result from the direct and/or indirect effects of EF on the plant. The application of non-pathogenic *Fusarium* strains to seeds, roots, or leaves has been shown to induce systemic resistance to pathogens, mediated by elicitors that trigger the expression of defense proteins such as glucanases, chitinases, peroxidases and PR proteins²⁸. Li et al.¹⁹ attributed the biocontrol potential of *Albifimbria verrucaria* against *B. cinerea* in *Vitis vinifera* to its chitinase activity. Similarly, Asim et al.² reported that endophytic *Fusarium* strains showed potential as bioherbicides in wheat (*Avena fatua*) due to their enzymatic activities, such as catalase and peroxidase. Therefore, these beneficial *Fusarium* isolates, in addition to producing growth-promoting metabolites, may also synthesize molecules that elicit induced systemic resistance.

This work documents non-pathogenic endophytic *Fusarium* isolates from *C. mirandae* with potential as plant growth promoters and biocontrol agents against fungal phytopathogens. These findings highlight the importance of endemic plants from protected natural areas as reservoirs of endophytic microorganisms with potential applications as biofertilizers or biocontrol agents in agriculture and forestry. Notably, this is the first report from Mexico of beneficial *Fusarium* isolates that promote plant growth and induce plant defense systems.

The advantages of beneficial *Fusarium* strains for plants have been reported²⁸. In the particular case of these isolates, agronomic, biochemical and molecular characterization will continue to confirm the safety of cultivating these fungi, specifically ensuring that they do not produce mycotoxins in food. Moreover, identifying the metabolites responsible for their beneficial effects on plants is essential. It will also be important to investigate the potential horizontal transmission of virulence genes between *Fusarium* species, to prevent the risk of these beneficial strains becoming pathogenic.

ORCID ID

Nidia del C. Ríos-De León: 0009-0000-5013-2691

Brenda del R. Saldaña-Morales: 0009-003-9882-0567

Daniel A. Pérez-Corral: 0000-0001-8298-2738

Vidal Hernández-García: 0000-0002-9383-6426

Claudio Ríos-Velasco: 0000-0002-3820-2156

Sergio Casas-Flores: 0000-0002-9612-9268

Luis A. Rodríguez-Larramendi: 0000-0001-8805-7180

Rubén Martínez-Camilo: 0000-0002-9057-8601

CRedit authorship contribution statement

NCRL conducted field and laboratory work, and contributed to writing the paper; BRSM and DPC performed laboratory work; VH, CRV provided equipment and reagents for the experiments; SCF provided equipment and reagents for the experiments and contribute to the writing of the paper; LARL performed data analysis; RMC conducted field work and data analysis; MASM design the experiments, write and reviewed the paper.

Ethical approval

This article does not include any studies involving human participants or animals conducted by the authors.

Informed consent

Informed consent was obtained from all individual participants included in the study.

Funding

This study was funded by University of Sciences and Arts of Chiapas (UNICACH).

Conflict of interest

The authors declare no conflict of interest.

Acknowledgments

We thank the Centro de Investigación en Alimentación y Desarrollo A.C. (CIAD-Unidad Cuauhtémoc) for their support in providing the facilities to carry out some experiments. We also thank Consejo Nacional de Humanidades, Ciencias y Tecnologías (CONAHCYT) for their contribution to the first author's master's scholarship.

References

1. Altschul SF, Gish W, Miller W, Myers EW, Lipman JD. Basic local alignment search tool. *J Mol Biol.* 1990;215:403–10, [http://dx.doi.org/10.1016/S0022-2836\(05\)80360-2](http://dx.doi.org/10.1016/S0022-2836(05)80360-2).
2. Asim S, Hussain A, Murad W, Hamayun M, Iqbal A, Rehman H, Tawab A, Irshad M, Alataway A, Dewidar AZ, Elansary HO, Lee IJ. Endophytic *Fusarium oxysporum* GW controlling weed and an effective biostimulant for wheat growth. *Front Plant Sci.* 2022;13:1–2, <http://dx.doi.org/10.3389/fpls.2022.922343>.
3. Barra-Bucarei L, France IA, Gerding GM, Silva AG, Carrasco-Fernández J, Franco CJ, Ortiz CJ. Antifungal activity of *Beauveria bassiana* endophyte against *Botrytis cinerea* in two Solanaceae crops. *Microorganisms.* 2019;8:1–15, <http://dx.doi.org/10.3390/microorganisms8010065>.

4. Cheng S, Jiang JW, Tan LT, Deng JX, Liang PY, Su H, Sun ZX, Zhou Y. Plant growth-promoting ability of mycorrhizal *Fusarium* strain KB-3 enhanced by its IAA producing endohyphal bacterium, *Klebsiella aerogenes*. *Front Microbiol.* 2022;13:1–12, <http://dx.doi.org/10.3389/fmicb.2022.855399>.
5. Coleman JJ. The *Fusarium solani* species complex: ubiquitous pathogens of agricultural importance. *Mol Plant Pathol.* 2016;17:146–58, <http://dx.doi.org/10.1111/mpp.12289>.
6. Coleman-Derr D, Desgarennes D, Fonseca-Garcia C, Gross S, Clingenpeel S, Woyke T, North G, Visel A, Partida-Martinez LP, Tringe SG. Plant compartment and biogeography affect microbiome composition in cultivated and native *Agave* species. *New Phytol.* 2016;209:798–811, <http://dx.doi.org/10.1111/nph.13697>.
7. Cun H, Munir S, He P, Wu Y, He P, Ahmed A, Che H, Li J, He Y. Diversity of root endophytic bacteria from maize seedling involved in biocontrol and plant growth promotion. *Egypt J Biol Pest Control.* 2022;32:1–9, <http://dx.doi.org/10.1186/s41938-022-00622-7>.
8. Dean R, Van Kan JAL, Pretorius ZA, Hammond-Kosack KE, Di Pietro A, Spanu PD. The top 10 fungal pathogens in molecular plant pathology. *Mol Plant Pathol.* 2012;13:414–30, <http://dx.doi.org/10.1111/j.1364-3703.2011.00783.x>.
9. Fauriah R, Djaya E, Djaenuddin N, Muis A, Nonci N. Potential of endophytic fungi as a pathogenic biocontrol agent and growth promoters in corn seedlings. *Egypt J Biol Pest Control.* 2023;33:1–7, <http://dx.doi.org/10.1186/s41938-023-00728-6>.
10. Fouda AM, Hassad SED, Eid AM, Eweis EED. Biotechnological applications of fungal endophytes associated with medicinal plant *Asclepias sinaica* (Bioss.). *Ann Agric Sci.* 2015;60:95–104, <http://dx.doi.org/10.1016/j.aosas.2015.04.001>.
11. García-Latorre C, Rodrigo S, Marin-Felix Y, Stadler M, Santamaria O. Plant-growth promoting activity of three fungal endophytes isolated from plants living in dehesas and their effect on *Lolium multiflorum*. *Sci Rep.* 2023;13:1–15, <http://dx.doi.org/10.1038/s41598-023-34036-8>.
12. Glick BR. Bacteria with ACC deaminase can promote plant growth and help to feed the world. *Microbiol Res (Pavia).* 2014;169:30–9, <http://dx.doi.org/10.1016/j.micres.2013.09.009>.
13. Grabka R, d'Entremont TW, Adams SJ, Walker AK, Tanney JB, Abbasi PA, Ali S. Fungal endophytes and their role in agricultural plant protection against pests and pathogens. *Plants (Basel).* 2021;11:1–29, <http://dx.doi.org/10.3390/plants11030384>.
14. Hamzah T, Lee S, Hidayat A, Terhem R, Faridah-Hanum I, Mohamed R. Diversity and characterization of endophytic fungi isolated from the tropical mangrove species, *Rhizophora mucronata*, and identification of potential antagonists against the soil-borne fungus, *Fusarium solani*. *Front Microbiol.* 2018;9:1–17, <http://dx.doi.org/10.3389/fmicb.2018.01707>.
15. Hatamzadeh S, Rahnema K, White JF, Oghaz NO, Nasrolahnejad S, Hemmati K. Investigation of some endophytic fungi from five medicinal plants with growth promoting ability on maize (*Zea mays* L.). *J Appl Microbiol.* 2022;134:1–15, <http://dx.doi.org/10.1093/jambio/lxac015>.
16. Herrera-Téllez VI, Cruz-Olmedo AK, Plasencia J, Gavilanes-Ruiz M, Arce-Cervantes O, Hernández-León S, Saucedo-García M. The protective effect of *Trichoderma asperellum* on tomato plants against *Fusarium oxysporum* and *Botrytis cinerea* diseases involves inhibition of reactive oxygen species production. *Int J Mol Sci.* 2019;20:1–13, <http://dx.doi.org/10.3390/ijms20082007>.
17. Khan MS, Gao J, Munir I, Zhang M, Liu Y, Moe TS, Xue J, Zhang X. Characterization of endophytic fungi, *Acremonium* sp., from *Lilium davidii* and analysis of its antifungal and plant growth-promoting effects. *Hindawi BioMed Res Int.* 2021;2021:1–14, <http://dx.doi.org/10.1155/2021/9930210>.
18. Kisaakye J, Fourie H, Haukeland S, Kisitu J, Nakimera S, Cortada L, Subramanian S, Coyne D. Endophytic non-pathogenic *Fusarium oxysporum*-derived dual benefit for nematode management and improved banana (*Musa* spp.) productivity. *Agriculture.* 2022;12:1–17, <http://dx.doi.org/10.3390/agriculture12020125>.
19. Li Z, Chang P, Gao L, Wang X. The endophytic fungus *Albifimbria verrucaria* from wild grape as an antagonist of *Botrytis cinerea* and other grape pathogens. *Phytopathology.* 2020;110:843–50, <http://dx.doi.org/10.1094/phyto-09-19-0347-r>.
20. Loudon BC, Haarmann D, Lynne AM. Use of blue agar CAS assay for siderophore detection. *J Microbiol Biol Educ.* 2011;1:51–3, <http://dx.doi.org/10.1128/jmbe.v12i1.249>.
21. Lü ZW, Liu HY, Wang CL, Chen X, Huang YX, Zhang MM, Huang QL, Zhang GF. Isolation of endophytic fungi from *Cotoneaster multiflorus* and screening of drought-tolerant fungi and evaluation of their growth-promoting effects. *Front Microbiol.* 2023;14:1–14, <http://dx.doi.org/10.3389/fmicb.2023.1267404>.
22. Marfetan JA, Vélez ML, Comerio R, Gallo A, Romero S. Rhizospheric *Penicillium* and *Talaromyces* strains in *Austrocedrus chilensis* native forests: identification and evaluation of biocontrol candidates against the pathogen *Phytophthora austrocedri*. *J Plant Prot Res.* 2023;63:239–53, <http://dx.doi.org/10.24425/jppr.2023.145757>.
23. Mehta P, Sharma R, Putatunda C, Walia A. Endophytic fungi: role in phosphate solubilization. In: Singh B, editor. *Advances in endophytic fungal research fungal biology*. Springer Cham; 2019. p. 183–209, http://dx.doi.org/10.1007/978-3-030-03589-1_9.
24. Mehta S, Nautiyal CS. An efficient method for qualitative screening of phosphate-solubilizing bacteria. *Curr Microbiol.* 2001;43:51–6, <http://dx.doi.org/10.1007/s002840010259>.
25. Moreno-Pérez JA. Efecto de los biofertilizantes sobre la calidad de plántulas de tomate (*Solanum lycopersicum* L.). Tesis de licenciatura en Ingeniería Agroforestal 2019. ANLIS “Dr. Miguel Ángel Salas Marina” y Universidad de Ciencias y Artes de Chiapas.
26. Moročko-Bičevska I, Fatehi J, Gerhardson B. Biocontrol of strawberry root rot and petiole blight by use of non-pathogenic *Fusarium* sp. strains. *Acta Hort.* 2014;1049:599–605, <http://dx.doi.org/10.17660/actahortic.2014.1049.93>.
27. Nutter FW, Esker PD, Netto RAC. Disease assessment concepts and the advancements made in improving the accuracy and precision of plant disease data. *Eur J Plant Pathol.* 2006;115:95–103, <http://dx.doi.org/10.1007/s10658-005-1230-z>.
28. Patil S, Sriram S. Biological control of *Fusarium* wilt in crop plants using non-pathogenic isolates of *Fusarium* species. *Indian Phytopathol.* 2020;73:11–9, <http://dx.doi.org/10.1007/s42360-020-00202-5>.
29. Pérez-Farrera MÁ. *Cycadales* spp. In: Chiapas, Mexico (*Ceratozamia mirandae*). Wg 3 – succulents and cycads. Case study 2. NDF workshop case studies; 2008. p. 1–12 [cited 2025 Mar 18]. Available from: https://cites.org/sites/default/files/ndf_material/WG3-CS2.pdf.
30. Perincherry L, Lalak-Kańczugowska J, Stępień Ł. *Fusarium*-produced mycotoxins in plant-pathogen interactions. *Toxins.* 2019;11:664, <http://dx.doi.org/10.3390/toxins11110664>.
31. Ponce-de León YM. Biocontrol *in vitro* de *Phytophthora capsici* y promoción del crecimiento vegetal de *Capsicum annuum* por actinobacterias no estreptomicetos. Tesis de maestría en Biotecnología 2022. ANLIS “Dr. Claudio Rios Velasco” y Centro de Investigación en Alimentación y Desarrollo, A.C. Cuauhtémoc, Chihuahua.
32. Poveda J, Hermosa R, Monte E, Nicolás C. *Trichoderma harzianum* favours the access of arbuscular mycorrhizal fungi to non-host Brassicaceae roots and

- increases plant productivity. *Sci Rep*. 2019;9:1–11, <http://dx.doi.org/10.1038/s41598-019-48269-z>.
33. Raeder U, Broda P. Rapid preparation of DNA from filamentous fungi. *Lett Appl Microbiol*. 1985;1:17–20, <http://dx.doi.org/10.1111/j.1472-765X.1985.tb01479.x>.
34. Ríos-Velasco C, Caro-Cisneros JM, Berlanga-Reyes DI, Ruiz-Cisneros MF, Ornelas-Paz JJ, Salas-Marina MA, Villalobos-Pérez E, Guerrero-Prieto VM. Identificación y actividad antagónica *in vitro* de aislados de *Bacillus* spp. Y *Trichoderma* spp. Contra hongos fitopatógenos comunes. *Rev Mex Fitopatol*. 2016;34:84–99, <http://dx.doi.org/10.18781/R.MEX.FIT.1507-1>.
35. Rodríguez-Larramendi LA, Salas-Marina MÁ, Hernández-García V, Campos-Saldaña RA, Cruz-Macías WO, de la Cruz-Morales M, Gordillo-Curiel A, Guevara-Hernández F. Physiological effect of water and nitrogen availability on guava plants. *Trop Subtrop Agroecosyst*. 2021;24:1–10, <http://dx.doi.org/10.56369/tsaes.3391>.
36. Salas-Marina MÁ, Silva-Flores MÁ, Cervantes-Badillo MG, Rosales-Saavedra MT, Islas-Osuna MA, Casas-Flores S. The plant growth-promoting fungus *Aspergillus ustus* promotes growth and induces resistance against different lifestyle pathogens in *Arabidopsis thaliana*. *J Microbiol Biotechnol*. 2011;21:686–96, <http://dx.doi.org/10.4014/jmb.1101.01012>.
37. Salas-Marina MÁ, Silva-Flores MÁ, Uresti-Rivera EE, Castro-Longoria E, Herrera-Estrella A, Casas-Flores S. Colonization of *Arabidopsis* roots by *Trichoderma atroviride* promotes growth and enhances systemic disease resistance through jasmonic acid/ethylene and salicylic acid pathways. *Eur J Plant Pathol*. 2011;131:15–26, <http://dx.doi.org/10.1007/s10658-011-9782-6>.
38. Sarmiento-Chacón M, Hernández-García V, Rodríguez-Larramendi LA, Salas-Marina MÁ, Ríos-Velasco C. *Neopestalotiopsis* sp. And *Colletotrichum karstii*, causal agents of leaf spots on camedoro palm (*Chamaedorea quezalteca*) in Mexico. *Rev Mex Fitopatol*. 2023;41:165–81, <http://dx.doi.org/10.18781/R.MEX.FIT.2302-7>.
39. Schneider C, Rasband W, Eliceiri K. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods*. 2012;9:671–5, <http://dx.doi.org/10.1038/nmeth.2089>.
40. Shaffer JP, Sarmiento C, Zalamea PC, Gallery RE, Davis AS, Baltrus DA, Arnold AE. Diversity, specificity, and phylogenetic relationships of endophytic bacteria in fungi that inhabit tropical seeds and leaves. *Front Ecol Evol*. 2016;4:116, <http://dx.doi.org/10.3389/fevo.2016.00116>.
41. Su Z, Zeng Y, Li X, Perumal AB, Zhu J, Lu X, Dai M, Liu X, Lin F. The endophytic fungus *Piriformospora indica*-assisted alleviation of cadmium in Tobacco. *J Fungi (Basel)*. 2021;7:1–17, <http://dx.doi.org/10.3390/jof7080675>.
42. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol*. 2011;28:2731–9, <http://dx.doi.org/10.1093/molbev/msr121>.
43. Turbat A, Rakk D, Vigneshwari A, Kocsabé S, Thu H, Szepesi A, Bakacsy L, Škrbić BD, Jigjiddorj EA, Vágvolgyi C, Szekeres A. Characterization of the plant growth-promoting activities of endophytic fungi isolated from *Sophora flavescens*. *Microorganisms*. 2020;8:1–15, <http://dx.doi.org/10.3390/microorganisms8050683>.
44. U'Ren JM, Arnold AE. Diversity, taxonomic composition, and functional aspects of fungal communities in living, senesced, and fallen leaves at five sites across. *PeerJ*. 2016;4:1–31, <http://dx.doi.org/10.7717/peerj.2768>.
45. Ulloa-Muñoz R, Olivera-Gonzales P, Castañeda-Barreto A, Villena GK, Tamariz-Angeles C. Diversity of endophytic plant-growth microorganisms from *Gentianella weberbaueri* and *Valeriana pycnantha*, highland Peruvian medicinal plants. *Microbiol Res (Pavia)*. 2020;233:1–11, <http://dx.doi.org/10.1016/j.micres.2020.126413>.
46. Watanabe T. Pictorial Atlas of soil and seed fungi: morphologies of cultured fungi and key to species. Washington, DC, USA: Taylor & Francis; 2002.
47. White TJ, Bruns T, Lee S, Taylor JW. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, editors. *PCR protocols: a guide to methods and applications*. New York: Academic Press; 1990. p. 315–22.
48. Yadav G, Meena M. Bioprospecting of endophytes in medicinal plants of Thar Desert: an attractive resource for biopharmaceuticals. *Biotechnol Rep (Amst)*. 2021;30:1–30, <http://dx.doi.org/10.1016/j.btre.2021.e00629>.