



REVISTA COLOMBIANA DE PSIQUIATRÍA

www.elsevier.es/rcp



Original article

Incidence of confusional syndrome (delirium) in a Latin American university hospital



Gabriel Fernando Oviedo Lugo^{a,b,c,*}, Paola Andrea García Martínez^{a,b,c},
Andrés Duarte Osorio^d, Carlos Javier Rincón^e, Alejandra López^f, Roberto Chavarría^f,
Catalina Pineda Betancur^f, Natalia García Gil^g, María Natalia Duarte Ariza^g,
Juan Agustín Patiño Trejos^h, Juan Sebastián Insignares^h, Carlos Gómez-Restrepo^{a,b}

^a Departamento de Psiquiatría y Salud Mental, Facultad de Medicina, Pontificia Universidad Javeriana, Bogotá, Colombia

^b Servicio de Psiquiatría, Hospital Universitario San Ignacio, Bogotá, Colombia

^c Centro de Memoria y Cognición Intellectus, Bogotá, Colombia

^d Departamento de Medicina Preventiva y Social, Facultad de Medicina, Pontificia Universidad Javeriana, Bogotá, Colombia

^e Departamento de Epidemiología Clínica y Bioestadística, Facultad de Medicina, Pontificia Universidad Javeriana, Bogotá, Colombia

^f Especialista en Psiquiatría y en Psiquiatría de enlace

^g Departamento de Psiquiatría, Pontificia Universidad Javeriana, Bogotá, Colombia

^h Facultad de Medicina, Pontificia Universidad Javeriana, Bogotá, Colombia

ARTICLE INFO

Article history:

Received 5 September 2021

Accepted 12 January 2022

Available online 22 April 2024

Keywords:

Delirium

Delirium assessment

Incidence

CAM

CAM ICU

Delirium subtypes

Geriatric Unit

ABSTRACT

Background: Little is known about the incidence of delirium and its subtypes in patients admitted to different departments of university hospitals in Latin America.

Objective: To determine the incidence of delirium and the frequency of its subtypes, as well as its associated factors, in patients admitted to different departments of a university hospital in Bogotá, Colombia.

Methods: A cohort of patients over 18 years of age admitted to the internal medicine (IM), geriatrics (GU), general surgery (GSU), orthopaedics (OU) and intensive care unit (ICU) services of a university hospital was followed up between January and June 2018. To detect the presence of delirium, we used the CAM (Confusion Assessment Method) and the CAM-ICU if the patient had decreased communication skills. The delirium subtype was characterised using the RASS (Richmond Agitation and Sedation Scale). Patients were assessed on their admission date and then every two days until discharged from the hospital. Those in whom delirium was identified were referred for specialised intra-institutional interdisciplinary management.

Results: A total of 531 patients admitted during the period were assessed. The overall incidence of delirium was 12% (95% CI, 0.3–14.8). They represented 31.8% of patients in the GU, 15.6% in the ICU, 8.7% in IM, 5.1% in the OU, and 3.9% in the GSU. The most frequent clinical display was the mixed subtype, at 60.9%, followed by the normoactive subtype (34.4%) and

DOI of original article: <https://doi.org/10.1016/j.rcp.2022.01.001>.

* Corresponding author.

E-mail address: goviedo@javeriana.edu.co (G.F. Oviedo Lugo).

2530-3120/© 2022 Asociación Colombiana de Psiquiatría. Published by Elsevier España, S.L.U. All rights reserved.

the hypoactive subtype (4.7%). The factors most associated with delirium were age (adjusted RR = 1.07; 95% CI, 1.05–1.09), the presence of four or more comorbidities (adjusted RR = 2.04; 95% CI, 1.31–3.20), and being a patient in the ICU (adjusted RR = 2.02; 95% CI, 1.22–3.35).

Conclusions: The incidence of delirium is heterogeneous in the different departments of the university hospital. The highest incidence occurred in patients that were admitted to the GU. The mixed subtype was the most frequent one, and the main associated factors were age, the presence of four or more comorbidities, and being an ICU patient.

© 2022 Asociación Colombiana de Psiquiatría. Published by Elsevier España, S.L.U. All rights reserved.

Incidencia del Síndrome Confusional (delirio) en un Hospital Universitario Latinoamericano

R E S U M E N

Palabras clave:

Delirio

Incidencia

CAM

CAM UCI

Subtipos de delirio

Unidad geriátrica

Introducción: Poco se conoce acerca de la incidencia del delirio en pacientes hospitalizados en diferentes servicios de hospitales universitarios en Latinoamérica y de los subtipos que se presentan.

Objetivo: Determinar la incidencia del delirio, la frecuencia de los subtipos motores y los factores asociados en pacientes hospitalizados en diferentes servicios de un hospital universitario en Bogotá, Colombia.

Métodos: Se dio seguimiento a una cohorte de pacientes mayores de 18 años hospitalizados en los servicios de Medicina Interna, Geriátrica, Cuidado Intensivo, Cirugía General y Ortopedia de un hospital universitario entre enero y junio de 2018. Para identificar la presencia de delirio, se utilizó la escala CAM (*Confusion Assessment Method*) y la CAM-ICU si el paciente presentaba disminución de las capacidades de comunicación. El subtipo de delirio se caracterizó utilizando la escala RASS (*Richmond Agitation and Sedation Scale*). Los pacientes fueron valorados el día de ingreso y luego cada 2 días hasta su alta hospitalaria. Se derivó a los pacientes en quienes se identificó delirio para tratamiento especializado interdisciplinario intrahospitalario.

Resultados: Se evaluó a 531 pacientes que ingresaron durante ese periodo a los servicios mencionados. La incidencia global del delirio fue del 12% (IC95%, 0,3–14,8). En orden descendiente, el 31,8% de los pacientes hospitalizados en el servicio de Geriátrica, el 15,6% en Cuidado Intensivo, el 8,7% en Medicina Interna, el 5,1% en Ortopedia y el 3,9% en Cirugía. El subtipo motor más frecuente fue el mixto (60,9%), seguido por el normoactivo (34,4%) y el hipoactivo (4,7%). Los factores asociados con la incidencia del delirio fueron la edad (RR ajustada = 1,07; IC95%, 1,05–1,09), la presencia de 4 o más comorbilidades (RR ajustada = 2,04; IC95%, 1,31–3,20) y la hospitalización en Cuidado Intensivo (RR ajustada = 2,02; IC95%, 1,22–3,35).

Conclusiones: La incidencia del delirio es heterogénea en los diferentes servicios del hospital universitario. La mayor incidencia se presentó en pacientes ingresados en el servicio de Geriátrica; el subtipo más frecuente fue el mixto y los principales factores asociados fueron la edad, la presencia de 4 o más comorbilidades y la hospitalización en Cuidado Intensivo.

© 2022 Asociación Colombiana de Psiquiatría. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Delirium, included for the first time as a clinical entity in the DSM-III of 1980, is a disturbance in attention and awareness that develops over a short period of time, usually hours or days,^{1,2} tends to fluctuate in severity during the course of the day, and may be accompanied by disorientation, cognitive impairment, memory deficit, impaired language and visuospatial ability or perception.^{1–3}

This condition is common in hospitalised patients, especially those admitted to intensive care units (ICU), where frequencies of up to 80% have been reported,³ and in elderly patients.^{4–6} Despite its known association with longer hospital stays, re-hospitalisations, increased hospital costs, risk of long-term cognitive impairment and increased mortality,^{7–9} it tends to be an underdiagnosed condition, probably due to its fluctuating nature, the overlap with depression or neurocognitive disorders and the lack of systematic application of formal hospital assessment instruments.

In Colombia, only a few studies have been performed that measure the incidence of this disorder, and the validity of many of these studies is questionable, given that no active searches for delirium are performed, or patients are not monitored throughout their full hospital stay.^{10–14} In a study conducted at our institution, a global incidence of delirium of 18.6% was found, with a higher percentage (60%) among patients admitted to the intensive care unit (ICU). However, as in many of the published studies, the motor subtype was not classified and the sample size (160 patients) limited the likelihood of being able to compare the incidence between different departments.¹⁵

There are four delirium motor subtypes: hyperactive, hypoactive, mixed and no motor subtype. Despite methodological problems faced when defining the delirium subtypes, the motor-activity profile is considered clinically relevant for their detection, and their relationship with other neuropsychiatric diagnoses, comorbidities, aetiology and treatment. Several studies have shown that the presence of a neurocognitive disorder, a prolonged duration and a hypoactive motor subtype are associated with a worse prognosis.^{6,16–18}

The aim of this study is to determine the incidence of delirium, to classify the motor subtype, to explore associated factors and to establish comparisons between the Surgery, Orthopaedics, Geriatrics and Internal Medicine departments and the ICU of Hospital San Ignacio (HUSI) in Bogotá.

Methods

This study describes a unique cohort of patients over 18 years of age admitted to the ICU and the Internal Medicine, Geriatrics, Orthopaedics and General Surgery departments of a high-complexity hospital during the first half of 2018.

Patients who were *delirious* or unconscious, or who were receiving antipsychotics, at the time of admission, were excluded. On day one of admission and then every 2 days while in the hospital, patients were assessed using the CAM (*Confusion Assessment Method*) or the CAM-ICU scales validated in Spanish and for Colombia.¹⁹ The delirium motor subtype was characterised using the RASS (*Richmond Agitation Sedation Scale*).^{20,21} Trained personnel were responsible for applying the scales and for collecting data. Most were residents from the general psychiatry and liaison psychiatry specialties. An institutional interdisciplinary group was responsible for treating the patients.

The cumulative incidence of delirium and the corresponding 95% confidence intervals (95% CI) were estimated, classified by sex, age, marital status, profession, hospital department and number of comorbidities. If a patient had more than one episode of delirium during their time in the hospital, only the first episode was quantified to calculate the incidence. For each variable, the relative risk (RR) was estimated, which was adjusted using a Poisson model for the variables of age, widowhood status, hospital department and number of comorbidities. The Stata 15.0 software tool was used for data processing and analysis.

The research protocol was submitted to the Ethics and Research Committees of the Faculty of Medicine and the hospital. All patients provided their written consent.

Table 1 – Sociodemographic characteristics and number of comorbidities in 531 hospitalised patients during the first half of 2018.

Variables	
Age	65 [48–79]
Female	261 (49.2)
Marital status	
Single	105 (19.8)
Married	206 (38.8)
Divorced	45 (8.5)
Widowed	95 (17.9)
Cohabiting	80 (15.1)
Profession	
Student	18 (3.4)
Unemployed	140 (26.4)
Vocational	169 (31.8)
Undergraduate	63 (11.9)
Retired	141 (26.6)
Education (years)	10 [5–11]
Comorbidities	
None	26 (4.9)
1	179 (33.7)
2	168 (31.6)
3	102 (19.2)
4 or more	56 (10.5)

Values are expressed as n (%) or median [interquartile range].

Results

A total of 531 patients who were admitted to the ICU and to the Internal Medicine (IM), Geriatrics, General Surgery and Orthopaedics departments were assessed and voluntarily gave their consent to participate. The sociodemographic data, hospital department and number of comorbidities are shown in [Table 1](#).

The age of the patients ranged between 18 and 96 years, with differences in medians between the different departments (53.9 years for patients in the Surgery department, 56.8 for those in Orthopaedics, 58.4 for those in Internal Medicine, 59.4 for patients in the ICU and 84.6 for those in Geriatrics). Half of the patients were elderly adults and 25% were 80 years of age or above. Although more men were admitted than women, this difference was small (50.8 compared to 49.2%). With regards to marital status, the frequency with which the patients reported being single, widowed or divorced is noteworthy (46.1%), and more so in women (59.8%) than in men (33%). Most patients had one or more comorbidities (95.1%).

The overall incidence of delirium, with its confidence intervals and broken down by department are shown in [Table 2](#). Statistically significant differences were evident in the incidence of delirium in patients admitted to the Geriatrics department compared to those admitted to the Orthopaedics, Internal Medicine and Surgery departments. However, despite a higher relative risk for patients admitted to the Geriatrics unit, no statistically significant differences were found with respect to the other departments. The mixed motor subtype was the most common (60.9%), followed by no motor subtype (34.4%) and the hypoactive motor subtype (4.7%). No statis-

Table 2 – Incidence of delirium and relative risks per hospital department.

Department	Patients, n		Incidence (95% CI), %	RR (95% CI)
	Delirium	Total		
Surgery (comparator)	4	102	3.92 (0.15–7.69)	—
Internal Medicine	13	150	8.67 (4.16–13.17)	2.21 (0.74–6.59)
Intensive Care	15	96	15.63 (8.36–22.89)	3.98 (1.37–11.58)
Orthopaedics	5	98	5.10 (0.75–9.46)	1.30 (0.36–4.70)
Geriatrics	27	85	31.76 (21.87–41.66)	8.10 (2.95–22.24)
Total	64	531	12.05 (9.28–14.82)	

95% CI: 95% confidence interval; RR: relative risk.

Table 3 – Incidence of delirium and relative risks according to age, sex, marital status, professional status and number of comorbidities.

Variables	Cases (n/N)	Incidence of delirium (95% CI), %	RR (95% CI)
Age			1.06 (1.04–1.08)
Sex			
Male (comparator)	26/270	9.6 (6.1–13.1)	—
Female	38/261	14.6 (10.3–18.8)	1.51 (0.95–2.42)
Marital status			
Single (comparator)	7/105	6.7 (1.9–11.4)	—
Married	25/206	12.1 (7.7–16.6)	1.82 (0.81–4.07)
Divorced	7/45	15.6 (5–26.1)	2.33 (0.87–6.27)
Widowed	18/95	18.9 (11.1–26.8)	2.84 (1.24–6.5)
Cohabiting	7/80	8.8 (2.6–14.9)	1.31 (0.48–3.59)
Professional status			
Student (comparator)	1/18	5.6 (0–16.1)	—
Unemployed	27/140	19.3 (12.8–25.8)	3.47 (0.5–24.03)
Vocational	10/169	5.9 (2.4–9.5)	1.07 (0.14–7.85)
Undergraduate	5/63	7.9 (1.3–14.6)	1.43 (0.18–11.46)
Retired	21/141	14.9 (9–20.8)	2.68 (0.38–18.75)
Comorbidities			
0	1/26	3.8 (0–11.2)	—
1	7/179	3.9 (1.1–6.8)	1.02 (0.13–7.93)
2	17/168	10.1 (5.6–14.7)	2.63 (0.37–18.94)
3	20/102	19.6 (11.9–27.3)	5.10 (0.72–36.25)
4 or more	19/56	33.9 (21.5–46.3)	8.82 (1.25–62.4)

95% CI: 95% confidence interval; RR: relative risk.

tically significant differences were found in the incidence of mixed and hypoactive delirium between the departments.

With regards to associated factors, statistically significant differences were found in the incidence and relative risk of delirium as age increased, for widowhood status and in the presence of four or more comorbidities (Table 3).

After adjusting the model for age in years, marital status, number of comorbidities and hospital unit, the results shown in Table 4 were obtained.

According to this model adjusted for the aforementioned variables, there are statistically significant differences in the risk of delirium according to age, presence of four or more comorbidities and admission to the ICU.

Discussion

The general incidence of delirium found may be considered low for a hospital with tier 4 level of complexity, and is also lower than the incidence previously reported at this

Table 4 – Relative risk of delirium adjusted for age, marital status, number of comorbidities and hospital department.

Variables	ARR (95% CI)
Age	1.07 (1.05–1.09)
Widowed	0.70 (0.44–1.12)
4 or more comorbidities	2.04 (1.31–3.2)
Admitted to the Intensive Care Unit	2.02 (1.22–3.35)

95% CI: 95% confidence interval; ARR: adjusted relative risk.

institution.^{15,22} We consider that both the current low incidence of delirium and its decrease over time are related to the implementation of an institutional policy for interdisciplinary treatment based on multi-component interventions that include environmental-based interventions, permanent review of drug interactions and de-prescribing for patients who are receiving multiple medications. Multi-component interventions in patients who are at risk of delirium have been

shown to be effective in the prevention of delirium,^{23,24} with statistically significant decreases in the incidence of delirium and in the frequency of in-hospital falls and costs of medical care.²³⁻²⁵

Likewise, the low incidence found in patients admitted to the ICU may be related to the aforementioned institutional policy, plus the prescription of central alpha-2 agonists for patients requiring conscious sedation or to be kept under deep sedation,²⁶ and to an open door component in which the visit schedule is free, broad and flexible. As a result, our institution can be considered a friendly hospital for the prevention of delirium.^{24,27}

The finding of a statistically significant relationship between age and the incidence of delirium agrees with the findings of several studies.^{16,17,28} The mixed motor subtype as the most common variant of delirium and the low frequency of the hypoactive variant may be related to the early diagnosis implemented in this study, together with the fact that patients were assessed by liaison psychiatry specialists, and patients whose primary symptoms were attributable to depression were labelled as having delirium, a bias that is often reported in the literature.²⁸⁻³⁰ However, our study was not designed to specifically explore these associations and these should therefore be clarified in subsequent research.

The prospective nature of the follow-up of patients admitted to the selected hospital departments and early identification of those patients who were starting to experience delirium support the validity of the results presented. However, the results are also limited by the fact that other risk factors were not assessed, such as the specific use of high-risk medications, standardised nutritional assessment guidelines, sensory deprivation or acute phase biomarkers.³¹

Funding

This research was performed during the time that the authors were under contract with the Faculty of Medicine of the Pontifical Javeriana University in Bogotá and San Ignacio University Hospital. Project Number FM-CIE-0493-17.

Conflicts of interest

There are no conflicts of interest for this study.

REFERENCES

1. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed. Washington: American Psychiatric Publishing; 2013.
2. NICE Guidelines. Delirium: prevention, diagnosis and management. Natl Collaborating Cent Acute Chronic Cond; 2019.
3. Maldonado JR. Acute brain failure: pathophysiology, diagnosis, management, and sequelae of delirium. *Crit Care Clin*. 2017;33:461-519.
4. Inouye SK. Delirium in hospitalized older patients: Recognition and risk factors. *J Geriatr Psychiatry Neurol*. 1998;11:118-25.
5. Schenning KJ, Deiner SG. Postoperative delirium in the geriatric patient. *Anesthesiol Clin*. 2015;33:505-16.
6. Evensem S. Delirium motor subtypes and prognosis in hospitalized geriatric patients – A prospective observational study. *J Psychosom Res*. 2019;122:24-8.
7. Maldonado JR. Acute brain failure: pathophysiology, diagnosis, management, and sequelae of delirium. *Crit Care Clin*. 2017;33:461-519.
8. Morandi A, Pandharipande P, Trabucchi M, Rozzini R, Mistraretti G, Trompeo AC, et al. Understanding international differences in terminology for delirium and other types of acute brain dysfunction in critically ill patients. *Intens Care Med*. 2008;34:1907-15.
9. Salluh JIF, Wang H, Schneider EB, Nagaraja N, Yenokyan G, Damluji A, et al. Outcome of delirium in critically ill patients: Systematic review and meta-analysis. *BMJ*. 2015;350:h2538.
10. Gutiérrez SJ, Posada-Guzmán A, Osorio-Tabares E, Jaramillo J, Cuartas S, Corredor- Marín M. Delirium en pacientes con fractura de cadera. *Hospital Universitario San Jorge, Pereira-2005. Rev Médica Risaralda*. 2006;12:4.
11. Flórez P, Velásquez J. Frecuencia y factores de riesgo del delirium en población geriátrica de la unidad de cuidados intensivos (UCI) de la Clínica Reina Sofía. *Rev Medica Sanitas*. 2009;12:66-73.
12. Rincón HG, Granados M, Unutzer J, Gomez M, Duran R, Badiel M, et al. Prevalence, detection and treatment of anxiety, depression, and delirium in the adult critical care unit. *Psychosomatics*. 2001;42:391-6.
13. Sánchez JC, González MI, Gutiérrez JC. Delirium in patients over 60 years of age in a tertiary level public hospital in the City of Pereira (Colombia): Under-diagnosed and under-recorded. *Rev Colomb Psiquiatr*. 2013;42:191-7.
14. Bernal DR, Ospina SO, Bernal JC. Delirium: incidencia y características clínicas y epidemiológicas en un hospital universitario. *Rev Colomb Psiquiatr*. 2009;38:471-87.
15. Gómez-Restrepo C, Murillo Mosquera AX, García Amador JM. Incidencia de delirium en pacientes hospitalizados en los servicios de medicina interna, cuidados intensivos, ortopedia y cirugía general del hospital universitario san Ignacio de Bogotá durante un mes típico. Bogotá: Repositorio biblioteca Pontificia Universidad Javeriana; 2012.
16. Vasilevskis EE, Han JH, Hughes CG, Ely EW. Epidemiology and risk factors for delirium across hospital settings. *Best Pract Res Clin Anaesthesiol*. 2012;26:277-87.
17. Krewulak KD, Stelfox HT, Leigh JP, Ely EW, Fiest KM. Incidence and prevalence of delirium subtypes in an adult ICU: a systematic review and meta-analysis. *Crit Care Med*. 2018;46:2029-35.
18. Siddiqi N, Harrison JK, Clegg A, Teale EA, Young J, Taylor J, et al. Interventions for preventing delirium in hospitalised non-ICU patients. *Cochrane Database System Rev*. 2016;3:CD005563.
19. Toro AC, Escobar LM, Franco JG, Díaz-Gómez JL, Muñoz JF, Molina F, et al. Versión en español del método para la evaluación de la confusión en cuidados intensivos, estudio piloto de validación. *Med Intensiva*. 2010;34:14-21.
20. Rojas-Gambasica JA, Valencia-Moreno A, Nieto-Estrada VH, Méndez-Osorio P, Molano-Franco D, Jiménez-Quimbaya AT, et al. Transcultural and linguistic adaptation of the Richmond Agitation-Sedation Scale to Spanish. *Rev Colomb Anesthesiol*. 2016;44:216-21.
21. Sessler CN, Gosnell MS, Grap MJ, Brophy GM, O'Neal PV, Keane KA, et al. The Richmond Agitation-Sedation Scale: validity and reliability in adult intensive care unit patients. *Am J Respir Crit Care Med*. 2002;166:1338-44.
22. Maldonado JR. Pathoetiological model of delirium: a comprehensive understanding of the neurobiology of

- delirium and an evidence-based approach to prevention and treatment. *Crit Care Clin.* 2008;24:789-856.
23. Hshieh TT, Yue J, Oh E, Puelle M, Dowall S, Trivison T, et al. Effectiveness of multicomponent nonpharmacological delirium interventions: a meta-analysis. *JAMA Intern Med.* 2015;175:512-20.
 24. Rosa RG, Tonietto TF, Da Silva DB, Gutierrez FA, Ascoli A, Madeira LC, et al. Effectiveness and safety of an extended ICU visitation model for delirium prevention: A before and after study. *Crit Care Med.* 2017;45:1660-7.
 25. O'Keeffe S, Lavan J. Predicting delirium in elderly patients: development and validation of a risk stratification model. *Age Ageing.* 1996;25:317-21.
 26. Pandharipande PP, Sanders RD, Girard TD, McGrane S, Thompson JL, Shintani AK, et al. Effect of dexmedetomidine versus lorazepam on outcome in patients with sepsis: an a priori-designed analysis of the MENDS randomized controlled trial. *Crit Care.* 2010;14:R38.
 27. O'Connell H, Kennelly S, Cullen W, Meagher D. Managing delirium in everyday practice: Towards cognitive-friendly hospitals. *Adv Psychiatr Treat.* 2014;20:380-9.
 28. O'Sullivan R, Inouye S, Meagher D. Delirium, and depression: Inter-relationship and clinical overlap in elderly people. *Lancet Psychiatry.* 2014;1:303-11.
 29. Devlin JW, Brummel NE, Al-Qadheeb NS. Optimising the recognition of delirium in the intensive care unit. *Best Pract Res Clin Anaesthesiol.* 2012;26:385-93.
 30. Farrell KR, Ganzini L. Misdiagnosing delirium as depression in medically ill elderly patients. *Arch Intern Med.* 1995;155:2459-64.
 31. Lindroth H, Bratzke L, Purvis S, Brown R, Coburn M, Mrkobrada M, et al. Systematic review of prediction models for delirium in the older adult inpatient. *BMJ Open.* 2018;8:e019223.