

Conclusions: Arg16Gly and Gln27Glu polymorphisms in the gene encoding Beta-2 Adrenergic Receptor (ADRB2) may be risk factors for reduced bone mineral density in females.

P03 -IDENTIFICATION OF NEUROPATHIC PAIN COMPONENT IN PATIENTS OF VARIOUS AGE WITH KNEE OSTEOARTHRITIS

V.V. Povoroznyuk, U.I. Pryimych

D.F. Chebotarev Institute of Gerontology NAMS Ukraine, Kyiv, Ukraine.

Introduction: Osteoarthritis-induced pain is a result of nociceptor stimulation, associated with local tissue damage and inflammation. Recent data suggest the presence of neuropathic pain symptoms in patients with osteoarthritis.

Objectives: The aim of this study was to estimate the structure of pain syndrome, reveal the presence of neuropathic pain (NP) component, symptoms and signs of NP in patients suffering from knee osteoarthritis.

Methods: We've examined 44 patients with knee osteoarthritis of the II-III stages by the Kallgren-Lawrence scale aged 47-85 years (average age 66.1 ± 1.5 years). To assess the NP component, we used screening scales painDETECT, LANSS, DN4 questionnaires. To assess intensity of pain, visual analogue scale (VAS) was used. Besides WOMAC and EuroQol-5D questionnaire were applied. For statistical analysis of results, ANOVA, correlation and regression analysis was applied.

Results: 4.6% of patients with knee osteoarthritis examined by painDETECT were likely to have the NP component. LANSS scale: 25% were probably to have NP. DN4 scale: 31.2% probably had NP. Moderate to significant correlations were found between intensity of pain by VAS data and Neuropathic Pain Scales (painDETECT, LANSS, DN4) data ($p < 0.05$). It was established that higher results of screening by painDETECT and DN4 positively correlate with a disturbance of physical function tested by WOMAC ($p < 0.05$). PainDETECT data have moderate to significant correlations with EuroQol-5D questionnaire ($p < 0.01$). Verbal descriptors as pins and needles, tingling, numbness and allodynia, pain from light touch which are revealed by 3 screening scales can significantly contribute to the likely neuropathic component in patients with knee osteoarthritis ($p < 0.05$). Burning pain ($p < 0.01$), pins and needles ($p < 0.05$) can be associated with a more severe pain in patients with knee osteoarthritis.

Conclusions: Thus, in patients with osteoarthritis the pain syndrome may reveal NP features. Identification of these would promote a targeted treatment strategy.

P04. EVALUATION OF BONE MINERAL DENSITY IN ADOLESCENTES WITH ANOREXIA NERVOSA

Patrícia Gouveia, Ricardo Teixeira, Rui Brito, Rosa Castro, Inês Amorim

Medicina Nuclear.

Introduction: One of Anorexia Nervosa's (AN) most frequent complications is the loss of bone mass, which represents a serious problem in adolescence, as it may reduce the peak bone mass attained, leading to osteopenia and osteoporosis.

Objectives: The objective of our study was to evaluate the prevalence of the reduction of Bone Mineral Density (BMD) in adolescents with AN, and its relation to anthropometric data (weight, height and BMI).

Methods: 66 adolescents with AN, referred to the Santo António Hospital Nuclear Medicine service between 2008 and 2015, were

retrospectively studied. A Hologic QDR-4500 densitometer was used to calculate BMD, Z-score and BMC. The correlation between variables was determined by linear regression analysis. The significance level was established at $p \leq 0.05$.

Results: The studied population presented average values of 0.86 ± 0.11 g/cm² of BMD, -0.67 ± 1.21 of Z-score and 26.82 ± 15.05 g of BMC. 14% of patients presented with bone mineralisation values below the variation expected for the age (Z-score < -2), it being that 40% of patients had a reduction of bone mass, defined, in this study, by a Z-score < -1 . Further, there was a significant correlation between the anthropometric data and bone mineral density, more notorious in relation to weight and to BMC.

Conclusions: The results showed that, in an adolescent population with AN, more than half of the individuals show no significant changes in BMD. These results are similar to those of previous studies and demonstrate that, in the disease's initial stages, BMD is relatively maintained. This study stresses the importance of early diagnosis of AN in adolescents, so as to find strategies to prevent loss of bone mass in this age group and impede the reduction in peak bone mass expected, which could leave to future severe complications.

P05. OSTEOPOROSIS IN NEUROLOGICAL DISORDERS

Vladyslav Povoroznyuk, Maryna Bystrytska, Iryna Karaban, Nina Karasevych

Institute of Gerontology NAMS Ukraine, Kyiv, Ukraine.

Introduction: Neurological disorders, notably stroke, Parkinson's disease and spinal cord injury are associated with increased falls risk and fracture risk.

Objectives: The aim of the research is to define the bone mineral density in patients, with neurological disorders.

Methods: We examined 48 men and women with Parkinson's disease (duration of Parkinson's disease was at list 5 years and all were on L-dopa treatment), 52 stroke patients (men and women), 21 patients with chronic spinal cord injury (AIS score A, B) and healthy people of appropriate age.

Results: BMD of patients with Parkinson's disease was significantly lower compared with BMD of healthy persons, both men and women (BMD lumbar spine women = 1.00 ± 0.15 vs 1.11 ± 0.15 g/cm², $p < 0.05$, the difference of BMD was about 15%, and BMD lumbar spine men = 1.16 ± 0.24 vs 1.29 ± 0.24 g/cm², $p < 0.05$). BMD stroke patients was also significantly lower compared with BMD of healthy persons: (BMD lumbar spine = 1.13 ± 0.20 vs 1.28 ± 0.46 g/cm², $p < 0.05$, BMD total hip = 0.97 ± 0.18 vs 1.09 ± 0.25 g/cm², $p < 0.05$). In women the difference was significant at the level of the total body and distal forearm. In men, the difference was significant only in the group of the patients with moderate and severe paresis. BMD SCI patients was dramatically low (BMD lumbar spine = 1.14 ± 0.19 vs 1.24 ± 0.26 g/cm², $p < 0.05$, BMD total hip = 0.82 ± 0.28 vs 1.07 ± 0.27 g/cm², $p < 0.05$), the difference of BMD was about 23%.

Conclusions: BMD in men and women with neurological disorders (Parkinson's disease, stroke and spinal cord injury) was significantly lower than in healthy persons of the same age.

P06. EVIDENCE OF ISOFLAVONES USE IN OSTEOPOROSIS PREVENTION

Ângela Lee Chin, Ana Filipa Vicente, Olena Berezovska Lourenço

Medicina Geral e Familiar.

Osteoporosis is characterized by decreased bone mass and microarchitectural deterioration of bone, increasing fracture risk.