



Scientific communications of 68th Conference of the Catalan Society of Neuropsychology

Comunicaciones científicas de la 68ª Jornada de la Societat Catalana de Neuropsicologia

The effects of distractors in unilateral spatial neglect: Evidence from an eye-tracking device

Alberto García-Molina^{1,2,3,*}, Rebeca Lauren Emerson^{1,2,3},
Jaume López Carballo^{1,2,3}, Stefan J. Lentelink⁴, Juan
García Fernández^{1,2,3}, Celeste Aparicio-López^{1,2,3}, Rocío
Sánchez-Carrión^{1,2,3}, Pablo Rodríguez Rajo^{1,2,3}, Antonia
Enseñat-Cantalops^{1,2,3}, Joan Deus Yela^{1,2,3}

¹ Institut Guttmann, Institut Universitari de Neurorehabilitació adscrit a la UAB, Badalona, Barcelona, Spain

² Universitat Autònoma de Barcelona, Bellaterra (Cerdanyola del Vallès), Spain

³ Fundació Institut d'Investigació en Ciències de la Salut Germans Trias i Pujol, Badalona, Barcelona, Spain

⁴ Twente University, Enschede, The Netherlands

Introduction: Disrupted visual search is a common symptom in unilateral spatial neglect (USN).

Objective: To examine the influence of distractors on visual behaviour in patients with USN when confronted with a vertically moving visual stimuli task using the approach of monitoring eye movements with an eye-tracking device.

Methods: Thirty-one right hemisphere stroke patients with USN were recruited. They performed a task in which participants were asked to click on the target stimuli that appeared from the top of the screen and moved vertically downwards. The task had two conditions: with and without distractors. We assessed performance in two ways, by scoring correct answers and by the percentage of time spent searching for targets measured over five vertical columns, i.e., (1) leftmost, (2) left, (3) central, (4) right, (5) rightmost.

Results: Participants scored a higher number of correct answers when there were no distractors on column 1 [$t(30)=2.619$; 95% CI: 0.003 to 0.029; $p=0.014$; Cohen's $d=0.3$] and column 2 [$t(30)=3.363$; 95% CI: 0.013 to 0.053; $p=0.002$; Cohen's $d=0.33$]. When distractors were present participants scored a higher number of correct answers on column 4 [$t(30)=-1.430$; 95% CI: -0.093 to -0.025; $p=0.001$; Cohen's $d=0.45$].

With respect to search time, participants spent more time searching for stimuli on column 1 when there were no distractors [$t(30)=2.723$; 95% CI: 0.006 to 0.042; $p=0.011$; Cohen's $d=0.23$]. When there were distractors more time was spent searching column 4 [$t(30)=-3.367$; 95% CI: -0.101 to -0.024; $p=0.002$; Cohen's $d=0.37$].

Conclusion: Distractors influence patients' performance and search time, limiting left visual search and reducing the number of correct left answers.

Keywords: Stroke; Unilateral spatial neglect; Eye-tracker; Distractors

References

- Adair, J. C., & Barrett, A. M. (2008). Spatial neglect: Clinical and neuroscience review: A wealth of information on the poverty of spatial attention. *Annals of the New York Academy of Sciences*, 1142(127), 21–43. doi:10.1196/annals.1444.008.
- Gainotti, G., De Luca, L., Figliozzi, F., & Doricchi, F. (2009). The influence of distracters, stimulus duration and hemianopia on first saccade in patients with unilateral neglect. *Cortex*, 45(4), 506–516. doi:10.1016/j.cortex.2007.12.008.

- Husain, M., Mannan, S., Hodgson, T., Wojciulik, E., Driver, J., & Kennard, C. (2001). Impaired spatial working memory across saccades contributes to abnormal search in parietal neglect. *Brain*, 124(5), 941–952. doi:10.1093/brain/124.5.941.
- Trepagnier, C. (2002). Tracking gaze of patients with visuospatial neglect. *Topics in Stroke Rehabilitation*, 8(4), 79–88. doi:10.1310/RH1W-Y1Y9-4Y8H-E933.

Registering visual search within unilateral spatial neglect using an eye-tracking device

Alberto García-Molina^{1,2,3,*}, Rebeca Lauren Emerson^{1,2,3}, Jaume López Carballo^{1,2,3}, Stefan J. Lentelink⁴, Juan García Fernández^{1,2,3}, Celeste Aparicio-López^{1,2,3}, Rocío Sánchez-Carrión^{1,2,3}, Pablo Rodríguez Rajo^{1,2,3}, Antonia Enseñat-Cantallops^{1,2,3}, Joan Deus Yela^{1,2,3}

¹ Institut Guttmann, Institut Universitari de Neurorehabilitació adscrit a la UAB, Badalona, Barcelona, Spain

² Universitat Autònoma de Barcelona, Bellaterra (Cerdanyola del Vallès), Spain

³ Fundació Institut d'Investigació en Ciències de la Salut Germans Trias i Pujol, Badalona, Barcelona, Spain

⁴ Twente University, Enschede, The Netherlands

Introduction: Patients with Unilateral Spatial Neglect (USN) display a biased exploratory visual search towards the contralesional hemifield.

Objective: To examine eye movements in patients with USN when confronted with a vertically moving stimuli task while monitoring visual search with an eye-tracking device.

Methods: Thirty-one right hemisphere stroke patients with USN were recruited. They performed a task in which participants were asked to click on the target stimuli that appeared from the top of the screen and moved vertically downwards. The task had two conditions: with and without distractors. The eye-tracking device registered ocular movements on both conditions. Based on the horizontal (X-axis; range: 0-1) and vertical coordinates (Y-axis; range: 1-0) of the screen we calculated the average gaze-position for each condition.

Results: The average gaze-position [X;Y] was [0.54;0.48] in the non-distractors condition and [0.59; 0.44] in the distractors condition. We found significant differences in the average gaze-position in the X-axis between the two conditions [0.54(0.11) vs. 0.59(0.12); $t(30) = -2.766$; 95% CI: -0.085 to -0.012 ; $p = 0.010$; Cohen's $d = 0.42$]. With respects to average gaze-position for axis Y there were also significant differences [0.48 (0.13) vs. 0.44 (0.14); $t(30) = 2.446$; 95% CI: 0.006 to 0.071; $p = 0.021$; Cohen's $d = 0.28$].

Conclusion: Distractors negatively influence visual search. They are able to modify the average gaze-position, both horizontally and vertically.

Keywords: Stroke; Unilateral spatial neglect; Eye-tracker; Visual search

References

- Behrmann, M. (1997). Impaired visual search in patients with unilateral neglect: An oculographic analysis.
- Gainotti, G., De Luca, L., Figliozzi, F., & Doricchi, F. (2009). The influence of distracters, stimulus duration and hemianopia on first saccade in patients with unilateral neglect. *Cortex*, 45(4), 506–516. doi:10.1016/j.cortex.2007.12.008.
- Plummer, P., Dunai, J., & Morris, M. E. (2006). Understanding the effects of moving visual stimuli on unilateral neglect following stroke. *Brain and Cognition*, 60(2), 156–165. doi:10.1016/j.bandc.2005.11.001.
- Trepagnier, C. (2002). Tracking gaze of patients with visuospatial neglect. *Topics in Stroke Rehabilitation*, 8(4), 79–88. doi:10.1310/RH1W-Y1Y9-4Y8H-E933.

Cognitive functions, coping strategies and psychological distress in patients with resected cancer receiving chemotherapy

Caterina Calderón Garrido^{1,*}, Alberto Carmona-Bayonas², Sara Fernández Arrojo³, Eva Martínez de Castro⁴, Alejandra Rodríguez Capote⁵, Pilar de la Morena², María Angeles Vicente², María de las Nieves Gómez Camacho⁵, Ana Fernández², Paula Jiménez-Fonseca³

¹ Facultad de Psicología, Universidad de Barcelona, Spain

² Hospital Universitario Morales Meseguer de Murcia, Spain

³ Hospital Universitario Central de Asturias, Oviedo, Spain

⁴ Hospital Universitario Marqués de Valdecilla, Santander, Spain

⁵ Hospital Universitario de Canarias, Tenerife, Spain

Background: The loss of cognitive functions is a symptom that impairs the quality of life and one of the most feared by cancer patients.

Objectives: The aim of this study is to analyze the cognitive functioning in cancer patients initiating chemotherapy and to study the relationship between coping strategies, and psychological distress.

Methods: NEOcoping is a prospective, multicenter and observational study. The project involves 23 Spanish hospitals and 34 researchers. The study has a website to collect clinical data and questionnaires for doctors and patients that are filled out before starting adjuvant chemotherapy and at the end of the treatment. The applied tests were: Mini-Mental Adjustment to Cancer (MAC), EORTC QLQ-C30 (cognitive function scale), and Brief Symptom Inventory (BSI-18).

Results: One hundred and ninety five patients' data (mean 58 years, and 60% female) were recruited. The primary tumor localization was mainly colon (41%) and breast (34%). The results indicate that patients perceive that their physical condition or treatment interferes with their cognitive functions (concentration, memory, etc.) have coping strategies more anxious characterized by a constant concern for health type, fear of the spread or recurrence of the disease that leads to frequent reaffirmation search ($t = 2.699$, $p = .008$). The cognitive function problems were

negatively related with somatization ($t=4.344$, $p<.001$), depression ($t=4.280$, $p<.001$) and anxiety ($t=4.777$, $p<.001$).

Conclusion: The perceived loss of cognitive abilities, uncertainty and loss of control of the cancer survivor patient may hinder adaptation to their situation, the loss of quality of life and greater emotional distress.

Keywords: Cognitive functions; Anxious strategies; Quality of life and psychological distress

Disclosure: This research has been funded by a grant from Sociedad Española de Oncología Médica (SEOM).

References

- Crist, J. V., & Grunfeld, E. A. (2013). Factors reported to influence fear of recurrence in cancer patients: A systematic review. *Psycho-Oncology*, 22, 978–986. <http://dx.doi.org/10.1002/pon.3114>.
- Mullaney, T., Olausson, K., Sharp, L., Zackrisson, B., Edvardsson, D., Nyholm, T. (2016). The influence of a department's psychosocial climate and treatment environment on cancer patients' anxiety during radiotherapy. *European Journal of Oncology Nursing*, 20, 113–118. <http://dx.doi.org/10.1016/j.ejon.2015.06.009>.

Social cognition in early phases of psychosis: A case-control study of gender-related differences

Guillem Navarra^{1,*}, Sol Fernández-Gonzalo^{1,2}, Marc Turón^{2,3}, Esther Pousa⁴, Diego Palao^{4,5}, Mercè Jodar⁶

¹ Clinical and Health Psychology Department, Universitat Autònoma de Barcelona (UAB), Bellaterra, Barcelona, Spain

² Research Department, Innovation and Investigation Institute Parc Taulí (I3PT), University Institute, UAB, Sabadell, Barcelona, Spain

³ CIBERES, Instituto de Salud Carlos III, Madrid, Spain

⁴ Mental Health Department, Parc Taulí Hospital, University Hospital-UAB, Sabadell, Barcelona, Spain

⁵ Department of Psychiatry and Forensic Medicine, Universitat Autònoma de Barcelona, Bellaterra, Barcelona, Spain

⁶ Neurology Department, ParcTaulí Hospital, University Hospital-UAB, Sabadell, Barcelona, Spain

Background: Deficits across multiple Social Cognition (SC) domains have been found in schizophrenia (Savla, Vella, Armstrong, Penn, & Twamley, 2012). However, gender-related differences in specific aspects of SC such as Theory of Mind (ToM) (Abu-Akel & Bo, 2013) and Emotional Processing (EP) (Erol, Putgul, Kosger, & Ersoy, 2013; Scholten, Aleman, Montagne, & Kahn, 2005) are almost unexplored.

Objective: To study possible gender-related differences in Affective (AToM) and Cognitive ToM (CToM) and EP skills in patients with recent diagnosis of psychosis and healthy subjects.

Methods: 39 patients (19 males, 20 females) with schizophrenia/schizoaffective disorder and 39 healthy subjects (19 males, 20 females), matched by age and years of

education, were recruited. The Reading the Mind in the Eyes Test (RMET) was used to evaluate AToM. First and Second-order False-Believe tasks (FB_1o and FB_2o) were used to evaluate CToM. The Pictures of Facial Affect (POFA) was used to evaluate EP.

Results: Patients' performance was significantly impaired in RMET ($p<0.001$), FB_2o ($p=0.002$) and POFA ($p<0.001$), but not in FB_1o. No significant differences were found between male and female patients in any tests. However, significant gender-related differences were observed in RMET ($p=0.004$) and POFA fear ($p=0.004$) in healthy subjects.

Conclusions: Gender-related differences were only found in healthy subjects, with females outperforming males in AToM and specific aspects of EP (fear recognition). Thus, these results suggest that SC performance in patients could be attributable to the illness itself, rather than gender. Further studies focused in the identification of the specific illness characteristics related to the SC difficulties in psychosis, and in the relation between EP and AToM in healthy subjects will be discussed.

Keywords: Psychosis; Gender differences; Social cognition; Theory of mind; Emotional processing

References

- Abu-Akel, A., & Bo, S. (2013). Superior mentalizing abilities of female patients with schizophrenia. *Psychiatry Research*, 210, 794–799.
- Erol, A., Putgul, G., Kosger, F., & Ersoy, B. (2013). Facial emotion recognition in schizophrenia: The impact of gender. *Psychiatry Investigation*, 10, 69–74.
- Savla, G., Vella, L., Armstrong, C., Penn, D., & Twamley, E. (2012). Deficits in domains of social cognition in schizophrenia: A meta-analysis of the empirical evidence. *Schizophrenia Bulletin*, 39(5), 979–992.
- Scholten, M., Aleman, A., Montagne, B., & Kahn, R. (2005). Schizophrenia and processing of facial emotions: Sex matters. *Schizophrenia Research*, 78, 61–67.

Decision making, obesity and binge eating. Taking the risk toward future interventions in eating disorders

J. Ottino^{1,3,8,*}, M. Garolera^{2,4,5}, I. Marqués-Iturria^{1,3}, P. González-Tartière¹, I. García-García^{1,3,6}, X. Caldú^{1,3,8}, X. Prats¹, M.J. Sender-Palacios^{4,7}, M. Vernet-Vernet^{4,7}, M.A. Jurado^{1,2,3,8}

¹ Department of Clinical Psychology and Psychobiology, University of Barcelona, Spain

² Consolidated Research Group in Neuropsychology, University of Barcelona, Spain

³ Institute of Neurosciences, University of Barcelona, Spain

⁴ Brain, Cognition and Behavior Clinical Research Group, Consorci Sanitari de Terrassa, Spain

⁵ Unit of Neuropsychology, Consorci Sanitari de Terrassa, Spain

⁶ Max Plank Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

⁷ CAP Terrassa Nord, Consorci Sanitari de Terrassa, Spain

⁸ Institut de Recerca Pediàtrica Hospital Sant Joan de Déu, Spain

Introduction: Binge eating (BE) is a disrupted eating pattern characterized by an excessive, rapid and compulsive food intake (Danner, Ouwehand, Van Haastert, Hornsvelt, & De Ridder, 2012; Voon, 2015). This may lead to overweight and obesity, plus some other psychological and medical conditions (Voon, 2015). Impairment in decision making (DM) was suggested as an underlying factor in disrupted eating behavior in both obesity and BE (Billingsley-Marshall, 2013; Danner et al., 2012; Davis, Patte, Curtis, & Reid, 2010; Manasse et al., 2014, 2015). Despite this, some studies did not find differences in DM between obese with and without BE (Danner et al., 2012; Manasse et al., 2014; Svaldi, Brand, & Tuschen-Caffier, 2010).

Objective: The aim of this study is to clarify the role that decision making has over eating behavior in order to suggest new frameworks of intervention.

Methods: DM was assessed using the Iowa's Gambling Test (IGT) on a healthy sample of obese adult women with ($n = 16$) and without BE ($n = 16$). BE group was formed according to BITE symptoms subscale (≥ 10). Parametric (t -test) and non-parametric tests (Mann-Whitney's U) were performed using IBM SPSS Statistics (v.21.0). Effect size (r) was calculated for every significant difference ($p < 0.05$).

Results: There were no significant differences in demographic variables. BE group performed worse in almost every IGT variables. BE group chose bad decks most (A-B) ($p = .009$, $r = .45$) over than good decks (C-D) ($p = .009$, $r = .45$), lost more money ($p = .022$, $r = .40$) and behaved more risky than non-BE group overall ($p = .012$, $r = .47$), showing difficulties in learning from negative feedback and hypersensitivity to positive reinforcement.

Conclusion: Impaired decision making may lead to unhealthy choices and disrupted eating behaviors. Besides psychological state and personality traits, executive function training might be a good aim for future interventions in eating disorders.

Keywords: Obesity; Binge eating; Decision making; IOWA; Executive functions

Grants: Ottino, J: Ayudas predoctorales de formación en Docencia e Investigación (ADR); Jurado, M.A.: PSI2013-48045-C2-1-P Obesidad y sobrepeso: Correlatos neuronales a lo largo de la vida. Caracterización de redes neuronales (Ministerio de Economía y Competitividad).

References

- Billingsley-Marshall, R. L., Basso, M. R., Lund, B. C., Hernandez, E. R., Johnson, C. L., Drevets, W. C., ... Yates, W. R. (2013). Executive function in eating disorders: The role of state anxiety. *International Journal of Eating Disorders*, 46(4), 316–321. doi:10.1002/eat.22086.
- Danner, U. N., Ouwehand, C., Van Haastert, N. L., Hornsvelt, H., & De Ridder, D. T. D. (2012). Decision-making impairments in women with binge eating disorder in comparison with obese and normal weight women. *European Eating Disorders Review*, 20(1). doi:10.1002/erv.1098.
- Davis, C., Patte, K., Curtis, C., & Reid, C. (2010). Immediate pleasures and future consequences. A neuropsychological study of binge eating and obesity. *Appetite*, 54(1), 208–213. doi:10.1016/j.appet.2009.11.002.
- Duchesne, M., Mattos, P., Carlos, J., Silvia, A., Freitas, R. De, Coutinho, G., ... Coutinho, W. (2010). Assessment of executive functions in obese individuals with binge eating disorder. *Revista Brasileira de Psiquiatria*, 32, 381–389.
- Manasse, S. M., Forman, E. M., Ruocco, A. C., Butryn, M. L., Juarascio, A. S., & Fitzpatrick, K. K. (2015). Do executive functioning deficits underpin binge eating disorder? A comparison of overweight women with and without binge eating pathology. *The International Journal of Eating Disorders*, 48(6), 677–683. doi:10.1002/eat.22383.
- Manasse, S. M., Juarascio, A. S., Forman, E. M., Berner, L. A., Butryn, M. L., & Ruocco, A. C. (2014). Executive functioning in overweight individuals with and without loss-of-control eating. *European Eating Disorders Review: The Journal of the Eating Disorders Association*, 22(5), 373–377. doi:10.1002/erv.2304.
- Svaldi, J., Brand, M., & Tuschen-Caffier, B. (2010). Decision-making impairments in women with binge eating disorder. *Appetite*, 54(1), 84–92. doi:10.1016/j.appet.2009.09.010.
- Voon, V. (2015). Cognitive biases in binge eating disorder: The hijacking of decision making. *CNS Spectrums*, 20(6), 566–573. doi:10.1017/S1092852915000681.

Estudi sobre el Rendiment Executiu dels pacients amb Trastorn de la Conducta Alimentària

Marta Molins^{1,2,*}, Anna Sintes¹, Meritxell Pacheco², Alba Castells^{1,3}, Jordina Tor⁴, Laia Portolés^{1,2}, Eduard Serrano¹, Montserrat Dolz¹

¹ Servei de Psiquiatria i Psicologia infanto-juvenil, Hospital Sant Joan de Déu, Esplugues de Llobregat, Barcelona, Spain

² Universitat Ramon Llull, Spain

³ Universitat de Barcelona, Spain

⁴ Fundació Sant Joan de Déu, Spain

Introduction: Eating disorders (ED) are increasingly affecting teenagers. There are cognitive deficits especially with the executive functions, highlighting the difficulties in central coherence and cognitive flexibility (Duchesne et al., 2004; Fagundo et al., 2012; Roberts, Tchanturia, & Treasure, 2010, 2013; Stuss et al., 2007; Tchanturia, Campbell, Morris, & Treasure, 2005). There is a growing interest in knowing whether they are predisposing factors or consequences of the disease, especially in child and adolescent population.

Method: The present study investigates neuropsychological changes associated with weight improvement, the differences between groups evaluated by the Body Mass Index (BMI) and the correlations between cognitive and psychopathological variables. The research includes 51 patients with eating disorders, admitted at the Hospital Sant Joan de Déu, and makes use of a neuropsychological battery, and self-report questionnaires. Subsequently, 13 patients who had achieved the established clinical improvement were included in the study.

Results: The results indicate that there are no differences in most cognitive domains in the longitudinal study ($p > 0.05$), although there are significant differences in cognitive flexibility, evaluated by the Trail Making Test B, getting better scores in the retest. There are differences by groups made by BMI in cognitive flexibility and resistance to interference ($p < 0.05$ and $p < 0.01$) being those who weigh less the ones who perform better. Moreover, there are significant correlations between executive functions and a wide range of psychopathological variables (BMI, Impulse Regulation, Drive for Thinness, Body Dissatisfaction, Depression, Anxiety, etc.).

Conclusions: These results suggest that the cognitive difficulties are a predisposing factor and, therefore, do not remit when the weight improves.

Keywords: Eating disorders; Neuropsychology; Executive functions; Cognitive flexibility; Central coherence

References

- Duchesne, M., Mattos, P., Fontenelle, L. F., Veiga, H., Rizo, L., & Appolinario, J. C. (2004). Neuropsychology of eating disorders: A systematic review of the literature. *Revista Brasileira de Psiquiatria*, 26(2), 107–117.
- Fagundo, A. B., de la Torre, R., Jiménez-Murcia, S., Agüera, Z., Granero, R., Tárrega, S., ... Fernández-Aranda, F. (2012). Executive functions profile in extreme eating/weight conditions: From anorexia nervosa to obesity. *PLoS ONE*, 7(8), 1–9.
- Roberts, M. E., Tchanturia, K., & Treasure, J. L. (2010). Exploring the neurocognitive signature of poor set-shifting in anorexia and bulimia nervosa. *Journal of Psychiatric Research*, 44(14), 964–970.
- Roberts, M. E., Tchanturia, K., & Treasure, J. L. (2013). Is attention to detail a similarly strong candidate endophenotype for anorexia nervosa and bulimia nervosa? *The World Journal of Biological Psychiatry*, 14(6), 452–463.
- Stuss, D. T., Robertson, I. H., Craik, F. I. M., Levine, B., Alexander, M. P., Black, S., ... Winocur, G. (2007). Cognitive rehabilitation in the elderly: A randomized trial to evaluate a new protocol. *Journal of the International Neuropsychological Society: JINS*, 13(1), 120–131.
- Tchanturia, K., Campbell, I. C., Morris, R., & Treasure, J. (2005). Neuropsychological studies in anorexia nervosa. *The International Journal of Eating Disorders*, 37(Suppl(April)), S72–S76; discussion S87–9.
- ³ Department of Clinical Sciences, School of Medicine, University of Barcelona, Barcelona, Spain
- ⁴ Mental Health Department, Corporació Sanitària Parc Taulí, Sabadell, Spain
- ⁵ Department of Psychiatry and Forensic Medicine, Universitat Autònoma de Barcelona, Barcelona, Spain
- ⁶ Department of Psychobiology and Methodology in Health Sciences, Universitat Autònoma de Barcelona, Barcelona, Spain

Introduction: Emotion regulation may be thought of as the capacity of prefrontal cortex regions to regulate activity in limbic regions (Ochsner, Silvers, & Buhle, 2012), but this frontolimbic connectivity may differ as a function of the emotion regulation strategy at use. In this study we aimed at differentiate the corticolimbic functional connectivity correlates of two emotion regulation strategies: cognitive reappraisal and suppression (Gross & John, 2003).

Methods: 48 healthy controls completed the Emotion Regulation Questionnaire (ERQ) (Gross & John, 2003) and underwent a resting-state functional magnetic resonance imaging acquisition. Functional connectivity (FC) of basolateral and centromedial amygdala (BLA/CMA) (Baur, Hänggi, Langer, & Jäncke, 2013) with a set of regions related to emotion regulation (prefrontal cortex, insula and cingulate cortex) was estimated with a seed-based approach. Scores in the suppression and reappraisal factors of the ERQ were correlated with FC maps.

Results: We found that FC between the left BLA and the left insula, and between the right BLA and the dorsomedial prefrontal cortex (dmPFC), was inversely correlated with reappraisal scores. On the other hand, FC between the right ventrolateral prefrontal cortex (vlPFC) and the left BLA correlated inversely with suppression scores, while FC between the dmPFC and the right BLA was directly correlated with suppression capacity. Finally, FC between the left CMA and the dmPFC was inversely correlated with suppression scores.

Conclusion: In conclusion, BLA connectivity appears to be more related to emotion regulation strategies than CMA connectivity, with a specific pattern of associations for reappraisal (lower left BLA-left insula and right BLA-dmPFC coupling) and for suppression (lower left BLA-right vlPFC connectivity and higher right BLA-dmPFC coupling) strategies.

Keywords: Functional connectivity; Emotion regulation; Amygdala; Cognitive reappraisal; Suppression

References

- Baur, V., Hänggi, J., Langer, N., & Jäncke, L. (2013). Resting-state functional and structural connectivity within an insula-amygdala route specifically index state and trait anxiety. *Biological Psychiatry*, 73(1):85–92.
- Gross, J. J., & John, O. P. (2003) Individual differences in two emotion regulation processes: Implications for affect, relationships, and well-being. *Journal of Personality and Social Psychology*, 85(2):348–362.
- Ochsner, K. N., Silvers, J. A., & Buhle, J. T. (2012). Functional imaging studies of emotion regulation: A synthetic

Emotion regulation strategies and frontolimbic functional connectivity

Maria Picó-Pérez^{1,3,*}, Pino Alonso^{1,2,3}, Narcís Cardoner^{1,4,5}, Marta Subirà^{1,2,3}, Clara López-Solà^{1,3,4}, Cinto Segalàs^{1,2}, Ignacio Martínez-Zalacain¹, Eva Real^{1,2}, Rosa Hernández-Ribas^{1,2,3}, José M. Menchón^{1,2,3}, Carles Soriano-Mas^{1,2,6}

¹ Department of Psychiatry, Bellvitge University Hospital-IDIBELL, Barcelona, Spain

² Carlos III Health Institute, Centro de Investigación Biomédica en Red (CIBERSAM), Barcelona, Spain

review and evolving model of the cognitive control of emotion. *Annals of the New York Academy of Sciences*, 1251(1):E1–E24.

Neurobiological correlates of electroconvulsive therapy effects in treatment-resistant depression: Longitudinal multimodal neuroimaging approach

Marta Cano^{1,2,*}, Narcís Cardoner^{1,3}, Ignacio Martínez-Zalacain¹, Harry Hallock⁴, Jesús Pujol^{5,6}, Mikel Urretavizcaya^{1,2,5}, Esther Via³, Oren Contreras-Rodríguez^{1,5}, José M. Menchón^{1,2,5}, Michael Valenzuela⁴, Carles Soriano-Mas^{1,5,7}

¹ Department of Psychiatry, Bellvitge University Hospital-IDIBELL, Hospitalet de Llobregat, Barcelona, Spain

² Department of Clinical Sciences, School of Medicine, University of Barcelona, Barcelona, Spain

³ Mental Health Department, Parc Taulí Sabadell, Universitat Autònoma de Barcelona, Barcelona, Spain

⁴ Regenerative Neuroscience Group, Brain and Mind Centre, University of Sydney, Sydney, Australia

⁵ Carlos III Health Institute, CIBERSAM, Spain

⁶ MRI Research Unit, Hospital del Mar, Barcelona, Spain

⁷ Department of Psychobiology and Methodology in Health Sciences, Universitat Autònoma de Barcelona, Barcelona, Spain

Background: Neuroplasticity and neuroinflammation have been proposed as two possible mechanisms of action of electroconvulsive therapy (ECT) (Andrade & Bolwig, 2014; Bouckaert et al., 2014, 2016). Although both hypotheses may be supported by medial temporal volume increases (Abbott et al., 2014; Bouckaert et al., 2016; Dukart et al., 2014; Joshi et al., 2016; Nordanskog et al., 2010; Nordanskog, Larsson, Larsson, & Johanson, 2014; Ota et al., 2015; Sartorius et al., 2016; Tendolkar et al., 2013) the specific mechanisms of action still remain unclear.

Objective: To assess changes in (1) regional gray matter (GM) volumes, (2) hippocampal subfields volumes, and (3) hippocampal metabolite concentrations during a bilateral ECT course, and to explore the putative differences between bilateral (BL) ECT vs. right unilateral (RUL) ECT.

Methods: Longitudinal structural magnetic resonance and spectroscopy study. We assessed 12 patients with treatment-resistant depression (TRD) throughout the entire course of a BL ECT protocol and 10 healthy participants at two neuroimaging examinations.

Results: Patients with TRD showed limbic complex and perigenual anterior cingulate cortex volume increases. Hippocampal subfield segmentation revealed volume increases of hippocampal CA2/3, subiculum and CA4/dentate gyrus regions. The limbic complex volume increase was associated with an hippocampal *N*-acetylaspartate concentration decrease and an hippocampal Glutamate+Glutamine concentration increase. Patients with TRD who underwent a RUL ECT protocol, showed a GM volume increase limited to the right hippocampal complex.

Conclusions: Our findings suggest that both forms of neuroplasticity and neuroinflammatory changes may be mechanisms of action of ECT. In this sense, our findings may also suggest the existence of a common neurobiological mechanism accounting for ECT-induced clinical improvement and cognitive side-effects (e.g., memory and executive function impairment (Scott, 2010; Semkovska, Keane, Babalola, & McLoughlin, 2011)).

Acknowledgments: This study was supported in part by the Carlos III Health Institute (PS09/01961 and CIBER-CB06/03/0034), FEDER funds, ‘‘A way to build Europe’’, and by the Agency of University and Research Funding Management of the Catalan Government (AGAUR; 2014SGR1672). M.C. is supported by a grant from the Spanish Ministry for Education, Culture and Sport (FPU13/02141). O.C.-R. is supported by Sara Borrell postdoctoral fellowships from the Carlos III Health Institute (CD14/00246). C.S.-M. is funded by a Miguel Servet contract from the Carlos III Health Institute (CP10/00604).

Keywords: Treatment-resistant depression; Electroconvulsive therapy; Hippocampus; *N*-acetylaspartate; Glutamate+Glutamine

References

- Abbott, C. C., Jones, T., Lemke, N. T., Gallegos, P., McClintock, S. M., Mayer, A. R., ... Calhoun, V. D. (2014). Hippocampal structural and functional changes associated with electroconvulsive therapy response. *Translational Psychiatry*, 4, e483. <http://dx.doi.org/10.1038/tp.2014.124>.
- Andrade, C., & Bolwig, T. (2014). Electroconvulsive therapy, hypertensive surge, blood–brain barrier breach, and amnesia: Exploring the evidence for a connection. *Journal of ECT*, 30, 160–164. <http://dx.doi.org/10.1097/YCT.0000000000000133>.
- Bouckaert, F., De Winter, F. L., Emsell, L., Dols, A., Rhebergen, D., Wampers, M., ... Vandenbulcke, M. (2016). Grey matter volume increase following electroconvulsive therapy in patients with late life depression: A longitudinal MRI study. *Journal of Psychiatry & Neuroscience*, 41, 105–114.
- Bouckaert, F., Sienaert, P., Obbels, J., Dols, A., Vandenbulcke, M., Stek, M., & Bolwig, T. (2014). ECT: Its brain enabling effects: A review of electroconvulsive therapy-induced structural brain plasticity. *Journal of ECT*, 30, 143–151. <http://dx.doi.org/10.1097/YCT.0000000000000129>.
- Dukart, J., Regen, F., Kherif, F., Colla, M., Bajbouj, M., Heuser, I., ... Draganski, B. (2014). Electroconvulsive therapy-induced brain plasticity determines therapeutic outcome in mood disorders. *Proceedings of the National Academy of Science of the United States of America*, 111, 1156–1161. <http://dx.doi.org/10.1073/pnas.1321399111>.
- Joshi, S. H., Espinoza, R. T., Pirnia, T., Shi, J., Wang, Y., Ayers, B., ... Narr, K. L. (2016). Structural plasticity of the hippocampus and amygdala induced by electroconvulsive therapy in major depression. *Biological Psychiatry*, 79, 282–292. <http://dx.doi.org/10.1016/j.biopsych.2015.02.029>.

- Nordanskog, P., Dahlstrand, U., Larsson, M. R., Larsson, E. M., Knutsson, L., & Johanson, A. (2010). Increase in hippocampal volume after electroconvulsive therapy in patients with depression: A volumetric magnetic resonance imaging study. *Journal of ECT*, 26, 62–67. <http://dx.doi.org/10.1097/YCT.0b013e3181a95da8>.
- Nordanskog, P., Larsson, M. R., Larsson, E. M., & Johanson, A. (2014). Hippocampal volume in relation to clinical and cognitive outcome after electroconvulsive therapy in depression. *Acta Psychiatrica Scandinavica*, 129, 303–311. <http://dx.doi.org/10.1111/acps.12150>.
- Ota, M., Noda, T., Sato, N., Okazaki, M., Ishikawa, M., Hattori, K., ... Kunugi, H. (2015). Effect of electroconvulsive therapy on gray matter volume in major depressive disorder. *Journal of Affective Disorders*, 186, 186–191. <http://dx.doi.org/10.1016/j.jad.2015.06.051>.
- Sartorius, A., Demirakca, T., Böhringer, A., Clemm von Hohenberg, C., Aksay, S. S., Bumb, J. M., ... Ende, G. (2016). Electroconvulsive therapy increases temporal gray matter volume and cortical thickness. *European Neuropsychopharmacology*, 26, 506–517. <http://dx.doi.org/10.1016/j.euroneuro.2015.12.036>.
- Scott, A. I. (2010). Electroconvulsive therapy, practice and evidence. *British Journal of Psychiatry*, 196, 171–172. <http://dx.doi.org/10.1192/bjp.bp.109.074054>.
- Semkowska, M., Keane, D., Babalola, O., & McLoughlin, D. M. (2011). Unilateral brief-pulse electroconvulsive therapy and cognition: Effects of electrode placement, stimulus dosage and time. *Journal of Psychiatric Research*, 45, 770–780. <http://dx.doi.org/10.1016/j.jpsychires.2010.11.001>.
- Tendolkar, I., van Beek, M., van Oostrom, I., Mulder, M., Janzing, J., Voshaar, R. O., & van Eijndhoven, P. (2013). Electroconvulsive therapy increases hippocampal and amygdala volume in therapy refractory depression: A longitudinal pilot study. *Psychiatry Research*, 214, 197–203. <http://dx.doi.org/10.1016/j.psychresns.2013.09.004>.

Cognitive changes in Parkinson's disease: A 4 year follow-up study

A. Campabadal^{1,2,3,*}, B. Segura^{1,2}, H.C. Baggio^{1,2}, C. Uribe^{1,2}, A. Abos^{1,2}, A.I. Garcia-Diaz^{1,2}, Y. Compta^{2,4,5}, M.J. Martí^{2,4,5}, F. Valdeoriola^{2,4,5}, C. Junque^{1,2,3,5}

¹ Medical Psychology Unit, Department of Medicine, Faculty of Medicine, University of Barcelona, Barcelona, Catalonia, Spain

² Institute of Neurosciences, University of Barcelona, Barcelona, Catalonia, Spain

³ Institute of Biomedical Research August Pi i Sunyer (IDIBAPS), Barcelona, Catalonia, Spain

⁴ Movement Disorders Unit, Neurology Service, Hospital Clinic de Barcelona, Barcelona, Catalonia, Spain

⁵ Centro de Investigación Biomédica en Red sobre Enfermedades Neurodegenerativas (CIBERNED), Hospital Clinic de Barcelona, Barcelona, Catalonia, Spain

Introduction: Although Parkinson's disease (PD) is a degenerative illness, changes over time in cognition are mild. Longitudinal studies have reported impairment in speed of

processing, memory and verbal fluency (Broeders et al., 2013; Muslimovic, Post, Speelman, De Haan, & Schmand, 2009). Cortical changes related to cognitive decline have been poorly investigated.

Objective: The aim of the study was to investigate differences in the progression of cognitive decline in PD patients compared to healthy controls (HC) and its neuroanatomical correlations.

Method: Forty-four PD patients and 22 HC underwent neuropsychological and magnetic resonance imaging assessment (scan interval $\mu = 45.7$ months). Repeated measures ANOVA were performed from adjusted z-scores (Aarsland, Brønnick, Larsen, Tysnes, & Alves, 2009). Patients were classified as having mild cognitive impairment (MCI) according to recent diagnostic criteria (Litvan et al., 2012). Subcortical segmentations and parcellations were extracted using FreeSurfer software and correlated with neuropsychological percentage of change.

Results: There were significant interactions group per time in semantic fluency ($F = 4.23$; $p = 0.044$), total recall ($F = 4.99$; $p = 0.029$) and delayed recognition from Rey's Auditory-Verbal Learning Test (RAVLT) ($F = 9.33$; $p = 0.003$), Trail Making Test Part A ($F = 5.29$; $p = 0.025$) and B ($F = 5.69$; $p = 0.020$), Symbol Digit Modalities Test ($F = 12.27$; $p < 0.0001$) and Stroop Word-Color ($F = 5.84$; $p = 0.019$). Over 4 years, 29 (66%) patients had PD-MCI from which 16 (36%) were cognitively preserved at baseline and 2 (5%) PD-MCI patients converted to PD dementia. Hippocampal volume correlated with percentage of change in RAVLT total recall ($r = 0.413$; $p = 0.011$).

Conclusions: PD patients had significant decline of speed processing, memory and semantic fluency and an increased rate of MCIs over time. Progressive hippocampal degeneration is related to verbal learning memory changes.

Keywords: Parkinson's disease; Mild cognitive impairment; Neuropsychological tests; Cortical atrophy

Disclosure: Spanish Ministry of Economy and Competitiveness (PSI201341393P), Generalitat de Catalunya (2014SGR 98), Fundació La Marató de TV3 in Spain (20142310), CU was supported by a fellowship from 2014, Spanish Ministry of Economy and Competitiveness (BES2014068173) and cofinanced by the European Social Fund (ESF) and AA was supported by a fellowship from 2016, Departament d'Empresa i Coneixement de la Generalitat de Catalunya, AGAUR (2016FI.B 00360).

References

- Aarsland, D., Brønnick, K., Larsen, J. P., Tysnes, O. B., & Alves, G. (2009). Cognitive impairment in incident, untreated Parkinson disease: The Norwegian ParkWest study. *Neurology*, 72(13), 1121–1126.
- Broeders, M., Velseboer, D. C., de Bie, R., et al. (2013). Cognitive change in newly-diagnosed patients with Parkinson's disease: A 5-year follow-up study. *Journal of the International Neuropsychological Society*, 19(6), 695–708.
- Litvan, I., Goldman, J. G., Tröster, A. I., et al. (2012). Diagnostic criteria for mild cognitive impairment in Parkinson's disease: Movement Disorder Society Task Force guidelines. *Movement Disorders*, 27(3), 349–356.

Muslimovic, D., Post, B., Speelman, J. D., De Haan, R. J., & Schmand, B. (2009). Cognitive decline in Parkinson's disease: A prospective longitudinal study. *Journal of the International Neuropsychological Society*, 15, 426–437.

Metamemory impairment is related to cortical thinning patterns in Parkinson's disease

C. Uribe^{1,2,*}, B. Segura^{1,2}, H.C. Baggio^{1,2},
A. Campabadal^{1,2,3}, A. Abos^{1,2}, A.I. Garcia-Diaz^{1,2},
Y. Compta^{2,4,5}, M.J. Martí^{2,4,5}, F. Valldeoriola^{2,4,5},
C. Junque^{1,2,3,5}

¹ Medical Psychology Unit, Department of Medicine, Faculty of Medicine, University of Barcelona, Barcelona, Catalonia, Spain

² Institute of Neurosciences, University of Barcelona, Barcelona, Catalonia, Spain

³ Institute of Biomedical Research August Pi i Sunyer (IDIBAPS), Barcelona, Catalonia, Spain

⁴ Movement Disorders Unit, Neurology Service, Hospital Clínic de Barcelona, Barcelona, Catalonia, Spain

⁵ Centro de Investigación Biomédica en Red sobre Enfermedades Neurodegenerativas (CIBERNED), Hospital Clínic de Barcelona, Barcelona, Catalonia, Spain

Introduction: Fronto-subcortical alterations have been widely described in Parkinson's disease (PD). Furthermore, frontal dysfunction has been related to metamemory impairment (Janowsky, Shimamura, & Squire, 1989; Smith, Souchay, & Moulin, 2011) which is still poor investigated in PD.

Objective: The aim of the current study was to investigate metamemory in PD patients with different patterns of cortical thinning. We hypothesized that patients with frontal atrophy will lose the normal correlation between real performance and its estimation due to the loss of feedback mechanism.

Method: Rey Auditory-verbal learning test and self-appraisal of the test performance were assessed in a sample of 41 non-demented PD patients and 22 healthy controls (HC). PD patients were classified in three different groups according to cortical thickness patterns identified by cluster analysis (Uribe et al., 2016). There was a first pattern with parietal-temporal atrophy; a second with frontal and occipital atrophy and a third without atrophy. Spearman's correlations were computed with SPSS software.

Results: There were significant correlations between RAVLT delayed recall and subjective performance in HC ($\rho=0.628$; $p=0.002$), all PD sample ($\rho=0.598$; $p<0.0001$) and PD-MCI patients ($\rho=0.699$; $p<0.0001$). Regarding PD subgroups, we obtained significant correlations in non-atrophic subgroup ($\rho=0.595$; $p=0.004$) and frontal and occipital atrophy subgroup ($\rho=0.661$; $p=0.007$). No significant correlations were found in patients with parietal-temporal atrophy ($\rho=0.289$; $p=0.637$).

Conclusions: The impairment of metamemory in PD is associated with the presence of parietal-temporal atrophy.

Keywords: Parkinson's disease, metamemory, magnetic resonance imaging, cortical thickness.

Disclosure: Spanish Ministry of Economy and Competitiveness (PSI201341393P), Generalitat de Catalunya (2014SGR 98), Fundació La Marató de TV3 in Spain (20142310), CU was supported by a fellowship from 2014, Spanish Ministry of Economy and Competitiveness (BES2014068173) and cofinanced by the European Social Fund (ESF) and AA was supported by a fellowship from 2016, Departament d'Empresa i Coneixement de la Generalitat de Catalunya, AGAUR (2016FI.B 00360).

References

- Janowsky, J. S., Shimamura, A. P., & Squire, L. R. (1989). Memory and metamemory: Comparisons between patients with frontal lobe lesions and amnesic patients. *Psychobiology*, 17(1), 3–11.
- Smith, S. J., Souchay, C., & Moulin, C. J. A. (2011). Metamemory and prospective memory in Parkinson's disease. *Neuropsychology*, 25(6):734–740.
- Uribe, C., Segura, B., Baggio, H. C., Abos, A., Martí, M. J., Valldeoriola, F., et al. (2016, Apr). Patterns of cortical thinning in nondemented Parkinson's disease patients. *Movement Disorders*.

Toxoplasmosis' coinfection and HIV+. Neuropsychological and neuroimaging study

Mireia Forner Puntonet*, Jordi Blanch Andreu, Araceli Rousaud Parés, Ricard Navinés de la Cruz, Sílvia Cañizares Alejos

Department of Psychiatry and Clinical Psychology, Clinic Institute of Neurosciences (ICN), Hospital Clínic, Barcelona, Spain

Introduction: Combination antiretroviral therapy (cART) has notably reduced HIV-associated dementia. However, the CHARTER study (Heaton, 2010) found a 47% prevalence of HIV-associated neurocognitive disorders (HAND) in patients under treatment.

HAND are associated with deficits in motor skills, processing speed and executive functions (Woods, Moore, Weber & Grant, 2009). Individuals with HIV+ and coinfections are most impaired on cognitive and psychomotor speed (Levine et al., 2008). Latent toxoplasmosis has been related to memory deterioration in seniors (Mendy, Vieira, Albatineh, & Gasana, 2014).

Different studies suggest an additive and synergistic effect of combined HIV and coinfections in the pathogenesis of cognitive impairment (Ciccarelli et al., 2013).

Objective: Describe the neuropsychological and neuroimaging profile of a VIH+ patient with toxoplasmosis.

Methods: A 73 years old woman, Independent for Basic and Instrumental ADL. VIH+, cART treatment until 2008. Treatment discontinuation until 2012. Jiroveci pneumonia (November 2011), cerebral toxoplasmosis (December 2011). Longitudinal neuropsychological study (February 2015, 2016): WAIS-IV, BNT, Verbal Fluency Test, RAVLT, WMS-III, RCFT, JLO, TMT, BADS, WCST, Stroop.

Results: The neuropsychological profile suggests a deep subcortical and frontal dysfunction with some progression during the last year.

Conclusions: Given the variability in the evolution of HAND, the worst evolution in older individuals and in coinfecting patients, it is recommended to carry out a neuropsychological assessment to study the evolution of patients' state, since often there is not a functional translation of the brain lesions.

A neuropsychological evaluation is compulsory in order to establish an early and specific cognitive rehabilitation.

Keywords: HIV; Toxoplasmosis; Coinfections; Neuropsychological profile

References

- Ciccarelli, N., Fabbiani, M., Grima, P., Falasca, K., Tana, M., Baldonero, E., ... Di Giambenedetto, S. (2013). Comparison of cognitive performance in HIV or HCV mono-infected and HIV-HCV co-infected patients. *Infection*, 41(6), 1103–1109.
- Heaton, R. K., Clifford, D. B., Franklin, D. R., Woods, S. P., Ake, C., Vaida, F., ... Grant, I. (2010). HIV-associated neurocognitive disorders persist in the era of potent antiretroviral therapy: CHARTER Study. *Neurology*, 75(23), 2087–2096.
- Levine, A. J., Hinkin, C. H., Ando, K., Santangelo, G., Martinez, M., Valdes-Sueiras, M., ... Singer, E. J. (2008). An exploratory study of long-term neurocognitive outcomes following recovery from opportunistic brain infections in HIV+ adults. *Journal of Clinical and Experimental Neuropsychology*, 30(7), 836–843.
- Mendy, A., Vieira, E. R., Albatineh, A. N., & Gasana, J. (2015). Immediate rather than delayed memory impairment in older adults with latent toxoplasmosis. *Brain, Behavior, and Immunity*, 45, 36–40.
- Woods, S. P., Moore, D. J., Weber, E., & Grant, I. (2009). Cognitive neuropsychology of HIV-associated neurocognitive disorders. *Neuropsychology Review*, 19(2), 152–168.

Brain structural correlates of subclinical obsessive-compulsive symptoms in healthy children

Maria Suñol^{1,*}, Carles Soriano-Mas², Oren Contreras-Rodríguez², Dídac Macià³, Gerard Martínez³, Ignacio Martínez-Zalacain¹, Marta Subirà², Jesús Pujol⁴, Jordi Sunyer⁵

¹ Department of Psychiatry, Bellvitge Biomedical Research Institute-IDIBELL, L'Hospitalet de Llobregat, Spain

² Department of Psychiatry, Bellvitge Biomedical Research Institute-IDIBELL; CIBERSAM, L'Hospitalet de Llobregat, Spain

³ MRI Unit, Hospital del Mar, Barcelona, Spain

⁴ MRI Unit, Barcelona, Spain; CIBERSAM, Barcelona, Spain

⁵ ISGlobal Centre for Research and Environment Epidemiology, CIBERESP, Barcelona, Spain

Background: Subclinical obsessive-compulsive (OC) symptoms are frequent in healthy individuals and their presence at infancy may predict an adult diagnosis of obsessive-compulsive disorder (OCD). In this study we aimed to study the brain structural correlates of subclinical OC symptoms in healthy children.

Methods: Two hundred and fifty-five children (mean age = 9.7 ± 0.8 years, 49% females) were assessed with the Obsessive-Compulsive Inventory-Child Version (OCI-CV) and underwent a brain structural magnetic resonance imaging examination. Total severity and the scores on the different symptom domains (doubting/checking, hoarding, ordering, obsessing, washing and neutralizing) were correlated with regional gray and white matter volumes following a voxel-based morphometry protocol. Statistical significance was set at $p < 0.05$, corrected for multiple comparisons.

Results: Mean total score on the OCI-CV was 9.17 ± 4.18 . Mean scores greater than 1 were observed in the doubting/checking (mean = 2.3 ± 1.5), hoarding (mean = 2.1 ± 1.3) ordering (mean = 2.3 ± 1.6) and obsessing (mean = 1.5 ± 1.4) dimensions. Total severity was associated with less gray and white matter in the left hippocampus, although at an uncorrected threshold. Doubt/checking symptoms correlated with less gray matter in the left middle frontal gyrus and more white matter in the right insula. Hoarding symptoms correlated with more gray and white matter in the left inferior frontal gyrus, and ordering symptoms correlated with less gray matter in the ventral caudate. Obsessing symptoms correlated with less white matter in the right temporal pole.

Conclusions: Subclinical dimension-specific OC symptoms are associated with distinct alterations in prefrontal, striatal and limbic/paralimbic areas. Such results may shed light on the neurobiological bases of clinical heterogeneity in OCD.

Keywords: Subclinical obsessive-compulsive symptoms; Symptom heterogeneity; Brain morphometry

The Spanish version of constraint-induced aphasia therapy improves aphasia severity, discourse effectiveness, and visual attention: Evidence from a case of acquired childhood conduction aphasia

Julia Vázquez De Sebastian^{1,*}, Isabel Leiva¹, Carmen García-Sánchez², Lourdes Janz³, María José Torres-Prioris⁴, Guadalupe Dávila⁴, Marcelo L. Berthier⁴

¹ Institut de Recerca de l'Hospital de la Santa Creu i Sant Pau, Barcelona, Spain

² Servei de Neurologia de l'Hospital de la Santa Creu i Sant Pau, Barcelona, Spain

³ Servei de Rehabilitació de l'Hospital de la Santa Creu i Sant Pau, Barcelona, Spain

⁴ Unidad de Neurología Cognitiva y Afasia (UNCA), Centro de Investigaciones Médico-Sanitarias (CIMES), Cátedra ARPA de Afasia, IBIMA, Universidad de Málaga, Spain

Introduction: Evidence-based childhood aphasia (CA) treatments are strongly needed. REGIA-Plus (Rehabilitación

Grupul Intensiva de la Afasia-Plus) (Berthier et al., 2014), the Spanish version of Constraint-Induced Aphasia Therapy, has not been used in CA so far.

Objectives: To report beneficial effects of REGIA-Plus in an adolescent with chronic conduction aphasia.

Method: A 14-year-old right-handed boy become aphasic after the cerebral bleeding of an AVM in the left hemisphere that required decompressive craniotomy and radiosurgery. Multiple baseline evaluations (2month apart) in the chronic period 9 months post-onset and 1 post-treatment testing were performed. The patient received REGIA-Plus (30h/10 semi-consecutive days) complemented with a set of expressive and written naming home tasks. The Primary outcome measures were Aphasia Severity (Aphasia Quotient-Western Aphasia Battery; AQ-WAB), Discourse Effectiveness (Words per Minute – WpM, Content Information Units-per minute (CIUpM) and % CIU) and Visual attention (CalCAP T-Scores; SART version). Background cognitive assessment was also performed at baseline.

Results: Two weeks of REGIA-Plus induced significant benefits in comparison to baselines in AQ-WAB (baselines: 81.2 and 81.3; REGIA-PLUS: 87.46), WpM (baselines: 69.76 and 69.47; REGIA-Plus: 84), CIUpM (baselines: 67.80 and 65.05; REGIA-Plus: 80.45) and CIU% (baselines: 65.47% and 70%; REGIA-PLUS: 83.74%), and Visual Attention (SART-CalCAP Choice RT; baseline T-Scores: 15 and 19; REGIA-Plus T-Score: 35).

Conclusions: This preliminary evidence indicates that REGIA-Plus can be used to improve aphasia, discourse effectiveness, and visual attention in acquired CA.

Keywords: Acquired childhood aphasia; Treatment; Constraint-induced aphasia therapy; Spanish-speaking

References

- Berthier, M. L., & Pulvermüller, F. (2011). Neuroscience insights improve neurorehabilitation of poststroke aphasia. *Nature Reviews Neurology*, 7(2), 86–97.
- Berthier, M. L., Green Heredia, C., Juarez Ruiz de Mier, R., Lara, J. P., & Pulvermüller, F. (2014). *REGIA. Rehabilitación Grupal Intensiva de la Afasia*. Madrid: TEA Ediciones.
- Johnson, C. J. (2006). Getting started in evidence-based practice for childhood speech-language disorders. *American Journal of Speech-Language Pathology*, 15(1), 20–35.
- Nicholas, L. E., & Brookshire, R. H. (1993). A system for quantifying the informativeness and efficiency of the connected speech of adults with aphasia. *Journal of Speech, Language, and Hearing Research*, 36(2), 338–350.
- Paquier, P. F., & Van Dongen, H. R. (1996). Review of research on the clinical presentation of acquired childhood aphasia. *Acta Neurologica Scandinavica*, 93, 428–436.
- Pulvermüller, F., Neininger, B., Elbert, T., Mohr, B., Rockstroh, B., Koebbel, P., & Taub, E. (2001). Constraint-induced therapy of chronic aphasia after stroke. *Stroke*, 32, 1621–1626.
- Szaflarski, J. P., Ball, A. L., Vannest, J., Dietz, A. R., Allendorfer, J. B., Martin, A. N., et al. (2015). Constraint-induced aphasia therapy for treatment of chronic post-stroke aphasia: A randomized, blinded, controlled pilot trial. *Medical Science Monitor*, 21, 2861–2869.

Is the MoCA a more valid cognitive screening tool to detect Alzheimer's disease and Mild Cognitive Impairment compared to the MMSE?: A bibliographic review

Sílvia Valldeneu^{1,*}, Amadeu Obach², Ángel Priego²

¹ *Universitat Oberta de Catalunya, Màster en Neuropsicologia, Spain*

² *Hospital Clínic de Barcelona, Spain*

Introduction: MMSE is the most widely used cognitive screening tool but its applicability, specificity and sensitivity compared to MoCA has been recently discussed.

Objectives: To review the results of existing studies which compared MoCA and MMSE for the detection of mild cognitive impairment and Alzheimer's disease.

Methods: Pubmed database was searched using the keywords "mmse, moca, Alzheimer" as a first screening. A second screening was made adding "mild cognitive impairment" keyword to the previous list. A total of 40 articles were obtained, being only 17 able to analyse after applying the exclusion criterion (other languages than English, other systematic reviews or studies not focused on MCI-AD).

Results: Compared to the MMSE, most of these 17 articles analysed agree that the MoCA has psychometric superiority in detecting MCI-AD, AD-nonAD and, specially, HC-MCI. Thanks to the greater difficulty of some items, the incorporation of executive functions and the more complex measures in all neurocognitive aspects, it seems to be more sensitive to detect MCI and early AD.

Highly variable cut-offs were reported for the MMSE and MoCA. There is a general agreement that MoCA is more sensitive, but at the recommended threshold MoCA favours sensitivity at the expense of specificity.

Conclusions: Compared to the MMSE, the MoCA is the most valid and clinically feasible screening tool to identify MCI. It allows a better prevision of the evolution, different levels of severity and determine MCI-AD subtypes. However, the optimal cut off score is open to debate and it would be convenient to do more investigation.

Keywords: Mini Mental State Examination (MMSE); Montreal Cognitive Assessment (MoCA); Alzheimer's disease (AD); Mild Cognitive Impairment (MCI)

References

- Damian, A. M., Jacobson, S. A., Hentz, J. G., Belden, C. M., Shill, H. A., Sabbagh, M. N., et al. (2011). The Montreal Cognitive Assessment and the mini-mental state examination as screening instruments for cognitive impairment: Item analyses and threshold scores. *Dementia and Geriatric Cognitive Disorders*, 31(2), 126–131. doi:10.1159/000323867.
- Freitas, S., Simões, M. R., Alves, L., & Santana, I. (2013). Montreal cognitive assessment: Validation study for mild cognitive impairment and Alzheimer disease. *Alzheimer's Disease and Associated Disorders*, 27(1), 37–43. doi:10.1097/WAD.0b013e3182420bfe.

- Fujiwara, Y., Suzuki, H., Yasunaga, M., Sugiyama, M., Ijuin, M., Sakuma, N., et al. (2010). Brief screening tool for mild cognitive impairment in older Japanese: Validation of the Japanese version of the Montreal Cognitive Assessment. *Geriatrics & Gerontology International*, 10(3), 225–232. doi:10.1111/j.1447-0594.2010.00585.x.
- Horton, D. K., Hynan, L. S., Lacritz, L. H., Rossetti, H. C., Weiner, M. F., & Cullum, C. M. (2015). An Abbreviated Montreal Cognitive Assessment (MoCA) for dementia screening. *The Clinical Neuropsychologist*, 29(4), 413–425. doi:10.1080/13854046.2015.1043349.
- Kaya, Y., Aki, O. E., Can, U. A., Derle, E., Kibaroglu, S., & Barak, A. (2014). Validation of Montreal Cognitive Assessment and discriminant power of Montreal Cognitive Assessment subtests in patients with mild cognitive impairment and Alzheimer dementia in Turkish population. *Journal of Geriatric Psychiatry and Neurology*, 27(2), 103–109.
- Lam, B., Middleton, L. E., Masellis, M., Stuss, D. T., Harry, R. D., Kiss, A., et al. (2013). Criterion and convergent validity of the Montreal Cognitive Assessment with screening and standardized neuropsychological testing. *Journal of the American Geriatric Society*, 61(12), 2181–2185. doi:10.1111/jgs.12541.
- Lee, J. Y., Dong Woo Lee, Cho, S. J., Na, D. L., Hong Jin Jeon, Kim, S. K., et al. (2008). Brief screening for mild cognitive impairment in elderly outpatient clinic: Validation of the Korean version of the Montreal Cognitive Assessment. *Journal of Geriatric Psychiatry and Neurology*, 21(2), 104–110. doi:10.1177/0891988708316855.
- Luis, C. A., Keegan, A. P., & Mullan, M. (2009). Cross validation of the Montreal Cognitive Assessment in community dwelling older adults residing in the Southeastern US. *International Journal of Geriatric Psychiatry*, 24(2), 197–201. doi:10.1002/gps.2101.
- Magierska, J., Magierski, R., Fendler, W., Kłoszewska, I., & Sobów, T. M. (2012). Clinical application of the Polish adaptation of the Montreal Cognitive Assessment (MoCA) test in screening for cognitive impairment. *Neurologia i Neurochirurgia Polska*, 46(2), 130–139.
- Nasreddine, Z. S., Phillips, N. A., Bédirian, V., Charbonneau, S., Whitehead, V., Collin, I., et al. (2005). The Montreal Cognitive Assessment, MoCA: A brief screening tool for mild cognitive impairment. *Journal of the American Geriatrics Society*, 53(4), 695–699.
- Razali, R., Jean-Li, L., Jaffar, A., Ahmad, M., Shah, S. A., Ibrahim, N., et al. (2014). Is the Bahasa Malaysia version of the Montreal Cognitive Assessment (MoCA-BM) a better instrument than the Malay version of the Mini Mental State Examination (M-MMSE) in screening for mild cognitive impairment (MCI) in the elderly? *Comprehensive Psychiatry*, 55(1), 70–75. doi:10.1016/j.comppsy.2013.04.010.
- Roalf, D. R., Moberg, P. J., Xie, S. X., Wolk, D. A., Moelter, S. T., & Arnold, S. E. (2012). Comparative accuracies of two common screening instruments for classification of Alzheimer's disease, mild cognitive impairment and healthy aging. *Alzheimers & Dementia: The Journal of the Alzheimer's Association*, 9(5), 529–537. doi:10.1016/j.jalz.2012.10.001.
- Smith, T., Gildeh, N., & Holmes, C. (2007). The Montreal Cognitive Assessment: Validity and utility in a memory clinic setting. *Canadian Journal of Psychiatry*, 52(5), 329–332.
- Trzepacz, P. T., Hochstetler, H., Wang, S., Walker, B., Saykin, A. J., & Alzheimer's Disease Neuroimaging Initiative (2015). Relationship between the Montreal Cognitive Assessment and Mini-mental State Examination for assessment of mild cognitive impairment in older adults. *BMC Geriatrics*, 15, 107. doi:10.1186/s12877-015-0103-3.
- Tsai, C. F., Lee, W. J., Wang, S. J., Shia, B. C., Nasreddine, Z., & Fuh, J. L. (2012). Psychometrics of the Montreal Cognitive Assessment (MoCA) and its subscales: Validation of the Taiwanese version of the MoCA and an item response theory analysis. *International Psychogeriatrics/IPA*, 24(4), 651–658. doi:10.1017/S1041610211002298.
- Van Steenoven, I., Aarsland, D., & Weintraub, D. (2015). Reply to a letter to the editor "Independent application of MoCA/MMSE conversion" by Armstrong et al. *Movements Disorder: Official Journal of the Movement Disorder Society*, 30(12), 1711. doi:10.1002/mds.26367.
- Wang, C. S., Pai, M. C., Chen, P. L., Hou, N. T., Chien, P. F., & Huang, Y. C. (2013). Montreal Cognitive Assessment and Mini-Mental State Examination performance in patients with mild-to-moderate dementia with Lewy bodies, Alzheimer's disease, and normal participants in Taiwan. *International Psychogeriatrics/IPA*, 25(11), 1839–1848. doi:10.1017/S1041610213001245.

Neural correlates of moral sensitivity as an endophenotype in obsessive-compulsive disorder

Caterina Viver^{1,*}, Marta Subirà^{1,2,3}, Maria Picó-Perez^{1,3}, Pino Alonso^{1,2,3}, Rosa Hernández-Ribas^{1,2,3}, Cinto Segalàs^{1,2}, Eva Real^{1,2}, Jesús Pujol^{2,4}, Ben Harrison⁵, Narcís Cardoner^{1,6,7}, José Manuel Menchón^{1,2,3}, Carles Soriano-Mas^{1,2,8}

¹ Department of Psychiatry, Bellvitge University Hospital-IDIBELL, Barcelona, Spain

² Carlos III Health Institute, Centro de Investigación Biomédica en Red (CIBERSAM), Barcelona, Spain

³ Department of Clinical Sciences, School of Medicine, University of Barcelona, Barcelona, Spain

⁴ Magnetic Resonance Unit, CRC-Hospital del Mar, Barcelona, Spain

⁵ Department of Psychiatry & Melbourne Health, Melbourne Neuropsychiatric Centre, The University of Melbourne, Melbourne, Australia

⁶ Mental Health Department, Corporació Sanitària Parc Taulí, Sabadell, Spain

⁷ Department of Psychiatry and Forensic Medicine, Universitat Autònoma de Barcelona, Barcelona, Spain

⁸ Department of Psychobiology and Methodology in Health Sciences, Universitat Autònoma de Barcelona, Barcelona, Spain

Introduction: OCD patients show a high moral sensitivity, which could be conceptualized in the cognitive biases that underpin OCD symptoms (Salkovskis, 1985). Moral judgments are related with the hyperresponsibility feelings and the

thought-action cognitive process, for which the mere occurrence of intrusions is perceived as the moral equivalent of deliberately acting on them (Rachman, 1997). The aim of this study was to compare the brain activation and connectivity maps obtained from a functional magnetic resonance study (fMRI) during a moral dilemma task between OCD patients, first-degree healthy relatives and healthy controls (HC).

Methods: 18 OCD patients, 19 first-degree healthy relatives and 19 HC were included in the study. The participants underwent an fMRI moral dilemma task and changes in BOLD signal were analyzed among the three groups. Analyses were focused on three predefined regions, previously related with this task in OCD (Harrison et al., 2012): the ventromedial prefrontal cortex (vmPFC), the dorsolateral prefrontal cortex (dlPFC) and the lateral temporal cortex. When significant differences in activation were found, we also performed psychophysiological interaction (PPI) analyses.

Results: OCD patients and healthy relatives showed a hyperactivation in the dlPFC in comparison to HC, while OCD patients alone showed a hyperactivation in the vmPFC. Regarding PPI analyses, OCD patients and healthy relatives showed a diminished vmPFC-neostriatum connectivity in comparison to HC, while OCD patients alone showed a diminished connectivity between the vmPFC and dopaminergic midbrain regions.

Conclusions: Results obtained from our study suggest the hyperactivation in the dlPFC during situations with high moral content as a candidate to an endophenotype in OCD.

Keywords: Moral sensitivity; OCD; Endophenotype; vmPFC; dlPFC

References

- Harrison, B. J., Pujol, J., Soriano-Mas, C., Hernández-Ribas, R., López-Solà, M., Ortiz, H., et al. (2012). Neural correlates of moral sensitivity in Obsessive-Compulsive Disorder. *Archives of General Psychiatry*, 69(7), 741–749.
- Rachman, S. (1997). A cognitive theory of obsessions. *Behaviour Research and Therapy*, 35(9), 793–802.
- Salkovskis, P. M. (1985). Obsessional-compulsive problems: A cognitive-behavioural analysis. *Behaviour Research and Therapy*, 23(5), 571–583.

Additional effect of early neurocognitive rehabilitation on executive cognitive function in subacute stroke patients, in the context of intensive rehabilitation program

L. Costa^{1,*}, S. Fernandez-Gonzalo², N. Ridao³, D. Cánovas⁴, M. Jódar^{1,4}

¹ Clinical and Health Psychology Department, Universitat Autònoma de Barcelona (UAB), Bellaterra, Barcelona, Spain

² Research Department, Innovation and Investigation Institute Parc Taulí (I3PT), Universitary Institute, UAB, Sabadell, Barcelona, Spain

³ Rehabilitation Medicine Department, Parc Taulí Hospital, Universitary Hospital-UAB, Sabadell, Barcelona, Spain

⁴ Neurology Department, Parc Taulí Hospital, Universitary Hospital-UAB, Sabadell, Barcelona, Spain

Objective: To explore the additional effect of early neurocognitive rehabilitation on executive functions (EEFF), processing speed (SP) and functional status in subacute stroke patients included in an intensive rehabilitation program.

Subjects and methods: Sixteen subacute stroke patients included in an intensive 3-week intensive rehabilitation program were randomized into two groups. Both groups received intensive general rehabilitation, including physical, occupational and speech therapy. The training group additionally received neurocognitive rehabilitation during 50 min/day, 5 times/week. Neurocognitive rehabilitation was based on paper-pencil tasks and was carried out by a neuropsychologist during the 3 weeks. Phonetic fluency (FAS) was used to assess EEFF, Symbol Search WAIS-IV-subtest for SP and the Barthel index for functional status.

Results: A significant time effect was observed in all cognitive (FAS: $F=20.36$; $p<0.001$ /Symbol Search: $F=8.78$; $p=0.01$) and functional (Barthel: $F=75.23$; $p<0.001$) measures. Moreover, a significant time by group effect ($F=6.54$; $p=0.023$) was observed in the FAS, favoring the training group. No other significant time by group effects were observed (Symbol Search: $F=1.59$; $p=0.23$ /Barthel: $F=0.08$; $p=0.78$).

Conclusion: Neurocognitive rehabilitation may add a beneficial impact, due to the specific improvement of the executive functions, in subacute stroke patients that received an intensive rehabilitation therapy. Neurocognitive rehabilitation had no specific effect on the functional status of the participants. However, further studies should be carried out to explore the impact of early neurocognitive rehabilitation on other cognitive domains and its relationship with other functional and quality of life outcomes.

Keywords: Neurocognitive rehabilitation; Rehabilitation therapy; Stroke; Cognitive functions

References

- Barker-Collo, S., Feigin, V. L., Parag, V., Lawes, C. M., & Senior, H. (2010). Auckland Stroke Outcomes Study. Part 2: Cognition and functional outcomes 5 years poststroke. *Neurology*, 75, 1608–1616.
- Boiko, E. A., Kulishova, T. V., Shumakher, G. I., & Iusup-khodzhaev, R. V. (2008). The role of physical exercises in the improvement of cognitive functions in patients who survived stroke, in the early rehabilitative period. *Vopr Kurortol Fizioter Lech Fiz Kult*, 6, 9–12.
- Cicerone, K. D., Dahlberg, C., Kahmar, K., Langenbahn, D. M., Malec, J. F., Bergquist, T. F., et al. (2000). Evidence-based cognitive rehabilitation: Recommendations for clinical practice. *Archives of Physical Medicine and Rehabilitation*, 81, 1596–1615.
- Cicerone, K. D., Langenbahn, D. M., Braden, C., et al. (2011). Evidence-based cognitive rehabilitation: Updated review of the literature from 2003 through 2008. *Archives of Physical Medicine and Rehabilitation*, 92(2), 519–530.

- Freeman, J. A., Hobart, J. C., Playford, E. D., Undy, B., & Thompson, A. J. (2005). Evaluating neurorehabilitation: Lessons from routine data collection. *Journal Neurology Neurosurgery and Psychiatry*, 76, 723–728.
- Gillespie, D. C., Bowen, A., Chung, C. S., Cockburn, J. C., Knapp, P., & Pollock, A. (2014). Rehabilitation for post-stroke cognitive impairment: An overview of recommendations arising from systematic reviews of current evidence. *Clinical Rehabilitation*, 1, 9.
- Gorelick, P. B., Scuteri, A., Black, S. E. et al. (2011). Vascular contributions to cognitive impairment and dementia: A statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*, 42, 267–713.
- Gottesman, R. F., & Hillis, A. E. (2003). Predictors and assessment of cognitive dysfunction resulting from ischaemic stroke. *Lancet Neurology*, 9, 895–905.
- Hyndman, D., Pickering, R. M., & Ashburn, A. (2008). The influence of attention deficits on functional recovery post stroke during the first 12 months after discharge from hospital. *Journal of Neurology. Neurosurgery and Psychiatry*, 79, 656–663.
- Kluding, P. M., Tseng, B. Y., Billinger, S. A. (2011). Exercise and executive function in individuals with chronic stroke: A pilot study. *Journal of Neurologic Physical Therapy*, 35, 11–27.
- Knopman, D. S., Roberts, R. O., Geda, Y. E., et al. (2009). Association of prior stroke with cognitive function and cognitive impairment. *Archives of Neurology*, 66, 614–619.
- Kwakkel, G., Kollen, B., Lindeman, E. (2004). Understanding the pattern of functional recovery after stroke: Facts and theories. *Restorative Neurology and Neuroscience*, 22, 281–299.
- Langhorne, P., Coupar, F., & Pollock, A. (2009). Motor recovery after stroke: A systematic review. *Lancet Neurology*, 8(8), 741–754.
- Lesniak, M., Bak, T., Czepiel, W., Seniow, J., Czlonkowska, A. (2008). Frequency and prognostic value of cognitive disorders in stroke patients. *Dementia Geriatrics Cognitive Disorders*, 26, 356–363.
- Loetscher, T., & Lincoln, N. B. (2013). Cognitive rehabilitation for attention deficits following stroke. *Cochrane Database Systematic Review*.
- Mackay, J., & Mensah, G. A. (2004). *The atlas of heart disease and stroke*. Geneva: World Health Organization.
- McDonnell, M. N., Bryan, J., Smith, & A. E., Esterman, A. J. (2011). Assessing cognitive impairment following stroke. *Journal of Clinical Neuropsychology*, 33, 945–953.
- Narasimhalu, K., Ang, S., De Silva, D. A., et al. (2011). The prognostic effects of poststroke cognitive impairments in nondisabled ischemic stroke patients. *Stroke*, 42, 883–888.
- Nys, G. M. S. (2005). *The neuropsychological of acute stroke: Characterisation and prognostic implications*. Tesi doctoral. Nederland: Universiteit Utrecht.
- Pendlebury, S. T., & Rothwell, P. M. (2009). Prevalence, incidence and factors associated with pre-stroke and post-stroke dementia: A systematic review and meta-analysis. *Lancet Neurology*, 8, 1006–1018.
- Pollock, A., Baer, G., Campbell, P., Choo, P. L., Forster, A., Morris, J., et al. (2014). Physical rehabilitation approaches for the recovery of function and mobility following stroke. *Cochrane Database Systematic Review*, 4.
- Rand, D., Eng, J. J., Liu-Ambrose, T., & Tawashy, A. E. (2010). Feasibility of 6-month exercise and recreation program to improve executive functioning and memory of individuals with chronic stroke. *Neurorehabilitation Neural Repair*, 24(8), 722–729.
- Riepe, M. W., Riss, S., Bittner, D., & Huber, R. (2003). Screening for cognitive impairment in patients with acute stroke. *Dementia and Geriatric Cognitive Disorders*, 17(1–2), 49–53.
- Rohling, M. L., Faust, M. E., Beverly, B., & Demakis, G. (2009). Effectiveness of cognitive rehabilitation following acquired brain injury: A meta-analytic re-examination of Cicerone et al.'s (2000, 2005) systematic reviews. *Neuropsychology*, 23(1), 20–39.
- Sameniene, J., Krisciunas, A., & Endzelyté, E. (2008). The evaluation of the rehabilitation effects on cognitive dysfunction and changes in psychomotor reactions in stroke patients. *Medicina (Kaunas)*, 44, 11.
- Stapleton, T., Ashburn, A., & Stack, E. (2001). A pilot study of attention deficits, balance control and falls in the subacute stage following stroke. *Clinical Rehabilitation*, 15(4), 437–444.
- Strong, K., Mathers, C., & Bonita, R. (2007). Preventing stroke: Saving lives around the world. *Lancet Neurology*, 6, 182–187.
- Sun, J. H., Tan, L., & Yu, J. T. (2014). Post-stroke cognitive impairment: Epidemiology, mechanisms and management. *Annals of Translational Medicine*, 2(8), 80.

High-resolution neuromelanin sensitive magnetic resonance as an early marker in Alzheimer's disease

I. Del Cerro^{1,*}, I. Martínez-Zalacain¹, G. Blasco², A. Guinea¹, J. Gascón-Bayarri³, R. Reñé³, V. Soria¹, J.M. Menchón^{1,4}, C. Soriano-Mas^{1,5}

¹ Department of Psychiatry, Bellvitge University Hospital-IDIBELL, Barcelona, Spain

² Department of Radiology, Girona Biomedical Research Institute, Diagnostic Imaging Institute, Girona, Spain

³ Unitat de Diagnòstic i Tractament de Demències, Servei de Neurologia, Hospital Universitari de Bellvitge, L'Hospitalet de Llobregat, Barcelona, Spain

⁴ Department of Clinical Sciences, Bellvitge Campus, Barcelona University, Barcelona, Catalonia, Spain

⁵ CIBER Salud Mental (CIBERSam), Instituto Salud Carlos III (ISCIII), Barcelona, Spain

Introduction: Alzheimer's disease (AD) is the first cause of dementia and supposes a very incapacitating illness with a high prevalence in elderly.

Nowadays, we have some diagnostic biomarkers that point to lesions, such neurofibrillary tangles, on brainstem regions in the early stages of the AD (Ferrer, 2012).

Specifically, we focus on locus coeruleus (LC), which is a noradrenergic nucleus situated at the dorsal pons that exhibits extensive cortical and subcortical projections and

has been implicated in modulation of attention (Aston-Jones & Cohen, 2005; Bouret & Sara, 2005).

Objectives: Studying selected patients at risk for developing AD with neuropsychological and neuroimaging techniques, focusing on involvement of LC and its connections.

Methods: Mild cognitive impairment, late-onset depression patients and healthy controls ($n = 30$ per group) will be assessed through clinical, neuropsychological and neuroimaging techniques.

Neuroimaging protocol will include a high-resolution neuromelanin sensitive magnetic resonance. Neuromelanin is a pigment, present on LC and other nuclei, which allow examine the signal intensity of LC and identify and quantify its volume in early stages of the AD.

LC will be functionally assessed through pupil response and an oddball task, which will supply an arousal measure.

Results: It is an underway project, but we expect to find between-group differences on volume and function of LC.

Conclusions: Identifying LC dysfunction could be therefore considered an early marker of AD. Neuroimaging techniques could in this way be used as a helpful tool for prevention, enabling us an advance and more effective intervention in these patients.

Keywords: Neuromelanin, Alzheimer's disease, neuroimaging, locus coeruleus

Project title: "Epigenetic and environmental factors bracing cognitive impairment and late-onset depression in elderly and early stages of Alzheimer's disease". Funded by: Proyectos integrados de excelencia en los IIS acreditados y Ciber.

References

- Aston-Jones, G., & Cohen, J. D. (2005). An integrative theory of locus coeruleus-norepinephrine function: Adaptive gain and optimal performance. *Annual Review of Neuroscience*, 28, 403–450.
- Bouret, S., & Sara, S. J. (2005). Network reset: A simplified overarching theory of locus coeruleus noradrenaline function. *Trends in Neurosciences*, 28(11), 574–582.
- Ferrer, I. (2012). Defining Alzheimer as a common age-related neurodegenerative process not inevitably leading to dementia. *Progress in Neurobiology*, 97, 38–51.

Alzheimer demented confabulators: More confabulations in episodic memory than in semantic memory

A. Llorente^{1,*}, C. Binschaedler², A. Jarne³

¹ Benito Menni Psychiatric Hospital, Sant Boi de Llobregat, Spain

² Neuropsychological and Neurorehabilitation Unit, Lausanne, Switzerland

³ Department of Personality, Barcelona University, Spain

Introduction: In classical view, confabulations are distortions of memory that appear when the subject is raised by specific questions that require retrieval of episodic or semantic memories. It is still controversial whether

confabulation is specific of a type of information, thus it would appear more often in episodic memory questions than in semantic memory questions.

Objectives: The aim of this study is to define how often confabulations appear in episodic and semantic memory questions and to provide more empirical evidence in a group of mild Alzheimer's dementia confabulators.

Method: Thirty-three subjects with mild Alzheimer's dementia and 37 elderlies without dementia between 65 and 85 years old have participated in this study. We used the Spanish version of Dalla-Barba's confabulation questionnaire and other neuropsychological tests. We divided the patients into two groups, which are "provoked confabulators" and "non-confabulators", using a statistical criterion. Sixty-four percent of the patients were considered "provoked confabulators".

Results: This group produces more confabulations in episodic memory questions than in semantic memory questions (Z de Wilcoxon = -3.60 , $p = .00$), and 2.80 times more than "non-confabulators" group (U de Mann-Whitney = 14 , $p < .01$). However, in this last group, no statistically significant differences were found between episodic memory questions and semantic memory questions (Z de Wilcoxon = 1.02 , $p = .31$).

Conclusions: In conclusion, in mild Alzheimer's dementia, provoked confabulators patients produce more confabulations in episodic memory questions than semantic memory questions. It seems that the confabulations are more specific to episodic memory. These results could invalidate some of the explanatory hypothesis of confabulation.

Keywords: Confabulation; Episodic and semantic memory; Dementia; Neuropsychological assessment

References

- Ghosh, V. E., Moscovitch, M., Melo, B., & Gilboa, A. (2014). Schema representation in patients with ventromedial PFC lesions. *Journal of Neuroscience*, 3(36), 12057–12070.
- La Corte, V., Serra, M., Attali, E., Boissé, M. F., & Dalla Barba, G. (2010). Confabulation in Alzheimer's disease and amnesia: A qualitative account and a new taxonomy. *Journal of the International Neuropsychological Society*, 16(6), 967–974.
- Schneider, A. (2008). *The confabulating mind. How the brain creates reality*. New York: Oxford University Press.

Exploring the neural bases of cognitive reserve in Huntington's disease

Maria Garau-Rolandi^{1,*}, Clara García-Gorro², Anira Escrichs³, Nadia Rodríguez-Dechichá¹, Irene Vaquer¹, Matilde Calopa⁴, Saül Martínez⁵, Jesús Pérez⁵, Jaime Kulisevsky⁵, Esteban Muñoz⁶, Pilar Santa-Cruz⁶, Jesús Ruiz⁷, Celia Mareca⁷, Nuria Caballol⁸, Susana Subirá¹, Ruth de Diego-Balaguer^{3,9}, Estela Cámara³

¹ Fundació Sociosanitària de Barcelona, Hospital Duran i Reynals, Spain

² Cognition and Brain Plasticity Unit, University of Barcelona, Spain

³ *Cognition and Brain Plasticity Unit, IDIBELL (Institut d' Investigació Biomèdica de Bellvitge), Spain*

⁴ *Fundació Bosch i Gimpera, Spain*

⁵ *Hospital Universitari de Bellvitge, Servei de Neurologia, Unitat de trastorns del moviment, Spain*

⁶ *Hospital de la Santa Creu i Sant Pau, Movement Disorders Unit, Spain*

⁷ *IDIBAPS, Hospital Clínic, Spain*

⁸ *Hospital Mare de Deu de la Mercè, Spain*

⁹ *Hospital de Sant Joan Despí Moisès Broggi, Spain*

¹⁰ *ICREA, Universitat de Barcelona, Spain*

Introduction: Although age at onset of Huntington's disease (HD) is mainly determined by the size of the CAG repeat expansion, other factors may play a role. One potential factor is Cognitive Reserve (CR), as it has been shown in other neurodegenerative disorders and aging.

Objectives: The objective of this study is to investigate the effect of CR in age of onset in HD and to examine the neural bases underlying the individual differences in executive dysfunction (ED) that could be due to the effects of CR in HD.

Method: 31 HD patients completed a CR questionnaire and were scanned using functional magnetic resonance imaging. We analyzed the Resting State Executive Control Network (RS-ECN), a novel approach to study the brain areas underlying ED. The strength of connectivity within this network was calculated voxel-wise. The difference between the estimated and the real age of onset (26 symptomatic-HD) was calculated for each patient.

Results: High levels of CR significantly delayed the appearance of the clinical symptoms. Functional connectivity analysis showed that higher CR was associated with lower connectivity strength in the anterior cingulate cortex and with an increase of connectivity in the superior parietal cortex (SPC). Furthermore, higher strength of connectivity in the left SPC was related to better performance in cognitive flexibility (TMT (b-a)) and working memory (backward digits) tasks.

Conclusions: These findings provide converging evidence that CR might act as a protective mechanism for the progression of the disease in the RS-ECN by delaying the onset of symptoms and improving the performance in executive functions.

Keywords: Cognitive reserve; Huntington's disease; Executive function; Individual differences

Decision making impairment: A shared vulnerability in obesity, gambling and substance use disorders?

Nuria Mallorqui-Bague^{1,2,*}, Ana B. Fagundo^{1,2}, Susana Jimenez-Murcia^{1,2,3}, Rafael de la Torre^{2,4}, Rosa M. Baños^{2,5}, Cristina Botella^{2,6}, Felipe F. Casanueva^{2,7}, Ana B. Crujeiras^{2,7}, Jose C. Fernández-García^{2,8}, Jose M. Fernández-Real^{2,9}, Gema Frühbeck^{2,10}, Roser Granero^{2,11}, Amaia Rodríguez^{2,10}, Francisco J. Ortega^{2,9}, Francisco J. Tinahones^{2,8}, Jose M. Menchón^{1,3,12}, Fernando Fernández-Aranda^{1,2,3}

¹ *Department of Psychiatry, University Hospital of Bellvitge-IDIBELL, Barcelona, Spain*

² *CIBER Fisiopatología Obesidad y Nutrición (CIBEROBN), Instituto Salud Carlos III, Spain*

³ *Department of Clinical Sciences, School of Medicine, University of Barcelona, Barcelona, Spain*

⁴ *Human Pharmacology and Clinical Neurosciences Research Group, Neuroscience Research Program, IMIM (Hospital del Mar Medical Research Institute), Barcelona, Spain*

⁵ *Department of Psychological, Personality, Evaluation and Treatment of the University of Valencia, Spain*

⁶ *Department of Basic Psychology, Clinic and Psychobiology of the University Jaume I, Castelló, Spain*

⁷ *Endocrine Division, Complejo Hospitalario U. de Santiago, Santiago de Compostela University, Spain*

⁸ *Department of Endocrinology and Nutrition, Hospital Clínico Universitario Virgen de Victoria, Málaga, Spain*

⁹ *Department of Diabetes, Endocrinology and Nutrition, Institut d'Investigació Biomèdica de Girona (IdIBGi) Hospital Dr Josep Trueta, Girona, Spain*

¹⁰ *Department of Endocrinology and Nutrition, Clínica Universidad de Navarra, University of Navarra, Pamplona, Spain*

¹¹ *Departament de Psicobiologia i Metodologia, Universitat Autònoma de Barcelona, Barcelona, Spain*

¹² *CIBER Salud Mental (CIBERSAM), Instituto Salud Carlos III, Spain*

Introduction: Addictions and behavioural addictions have been associated with decision making impairments. The present study explores decision making profiles of Substance use disorder (SUD), Gambling Disorder (GD) and Obesity (OB) when assessed by the Iowa Gambling Test (IGT) and compares them with a healthy control group (HC).

Methods: For the aims of this study, 591 participants (51.7% females) that were distributed as follow 194 HC, 178 GD, 113 OB, 106 SUD were assessed according to DSM-V criteria, completed a sociodemographic interview and conducted the Iowa Gambling Task.

Results: SUD, GD and OB present impaired decision making when compared with the HC in both the overall task and learning across the task parameters, however no differences are found for the overall performance in the IGT among the clinical groups. Finally, results reveal some specific learning across the task patterns within the three clinical groups: OB individuals maintain negative scores until the third set were learning starts but with a less extend to healthy controls, SUD seems to present an early learning than OB followed by a progressive although slow improvement and GD presents more random choices displaying very few or no learning.

Conclusions: Decision making impairments are present in SUD, GD and OB. Some specific patterns of learning across the task are identified for each of the here studied clinical samples which to some extent show differences in learning patterns of decision making task. These results can help understanding the underlying mechanisms of obesity and addiction behaviors as well as improve current clinical treatments.

Keywords: Obesity; Gambling disorder; Substance use disorder; Addiction; Decision making

References

- Avena, N. M., Rada, P., & Hoebel, B. G. (2009). Sugar and fat bingeing have notable differences in addictive-like behavior. *Journal of Nutrition*, 139(Mar (3)), 623–628. Available from <http://jn.nutrition.org/content/139/3/623.short> Cited 31.08.15.
- Barry, D., Clarke, M., & Petry, N. M. (2009). Obesity and its relationship to addictions: Is overeating a form of addictive behavior? *American Journal on Addictions*, 18(6), 439–451.
- Bechara, A. (1997). Deciding advantageously before knowing the advantageous strategy. *Science* (80-), 275(February (5304)), 1293–1295. Available from <http://www.sciencemag.org/content/275/5304/1293.abstract> Cited 18.07.14.
- Bechara, A., Dolan, S., Denburg, N., Hindes, A., Anderson, S. W., & Nathan, P. E. (2001). Decision-making deficits, linked to a dysfunctional ventromedial prefrontal cortex, revealed in alcohol and stimulant abusers. *Neuropsychologia*, 39(Jan (4)), 376–389. Available from <http://www.ncbi.nlm.nih.gov/pubmed/11164876> Cited 18.01.16.
- Brevers, D., Koritzky, G., Bechara, A., & Noël, X. (2014). Cognitive processes underlying impaired decision-making under uncertainty in gambling disorder. *Addictive Behaviors*, 39(Oct (10)), 1533–1536. Available from <http://www.sciencedirect.com/science/article/pii/S0306460314001919> Cited 03.12.15.
- Brogan, A., Hevey, D., O'Callaghan, G., Yoder, R., & O'Shea, D. (2011). Impaired decision making among morbidly obese adults. *Journal of Psychosomatic Research*, 70(Feb (2)), 189–196. Available from <http://www.sciencedirect.com/science/article/pii/S0022399910003107> Cited 14.12.15.
- Brogan, A., Hevey, D., & Pignatti, R. (2010). Anorexia, bulimia, and obesity: Shared decision making deficits on the Iowa Gambling Task (IGT). *Journal of the International Neuropsychological Society*, 16, 711–715.
- Fagundo, A. B., De La Torre, R., Jiménez-Murcia, S., Agü Era, Z., Granero, R., Tá Rrega, S., et al. (2012). Executive functions profile in extreme eating/weight conditions: From anorexia nervosa to obesity. *PLoS ONE*, 7(8).
- First, M. B., Spitzer, R. L., Gibbon Miriam, & Williams, J. B. (2002). Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Patient Edition (SCID-I/P). New York: Biometrics Research, New York State Psychiatric Institute.
- Gearhardt, A. N., & Corbin, W. R. (2011). The role of food addiction in clinical research. *Current Pharmaceutical Design*, 17(Jan (12)), 1140–1142. Available from <http://www.ncbi.nlm.nih.gov/pubmed/21492090> Cited 26.01.16.
- Gearhardt, A. N., Yokum, S., Orr, P. T., Stice, E., Corbin, W. R., & Brownell, K. D. (2011). Neural correlates of food addiction. *Archives of General Psychiatry*, 68(Aug (8)), 808–816. Available from <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3980851&tool=pmcentrez&rendertype=abstract> Cited 15.12.15.
- Goudriaan, A. E., Oosterlaan, J., de Beurs, E., & van den Brink, W. (2005). Decision making in pathological gambling: A comparison between pathological gamblers, alcohol dependents, persons with Tourette syndrome, and normal controls. *Brain Research. Cognitive Brain Research*, 23(Apr (1)), 137–151. Available from <http://www.sciencedirect.com/science/article/pii/S0926641005000236> Cited 20.01.16.
- Grant, J. E., Derbyshire, K., Leppink, E., & Chamberlain, S. R. (2015). Obesity and gambling: Neurocognitive and clinical associations. *Acta Psychiatrica Scandinavica*, 131(May (5)), 379–386. Available from <http://www.ncbi.nlm.nih.gov/pubmed/25346399> Cited 26.01.16.
- Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6, 65–70.
- Krmpotich, T., Mikulich-Gilbertson, S., Sakai, J., Thompson, L., Banich, M. T., & Tanabe, J. (2015). Impaired decision-making, higher impulsivity, and drug severity in substance dependence and pathological gambling. *Journal of Addiction Medicine*, 9(January (4)), 273–280. Available from <http://www.ncbi.nlm.nih.gov/pubmed/25918968> Cited 04.12.15.
- Smith, E., Hay, P., Campbell, L., & Trollor, J. N. (2011). A review of the association between obesity and cognitive function across the lifespan: Implications for novel approaches to prevention and treatment. *Obesity Reviews*, 12(September (9)), 740–755. Available from <http://www.ncbi.nlm.nih.gov/pubmed/21991597> Cited 01.10.15.
- Tanabe, J., Krmpotich, T., Mikulich, S., Sakai, J., Thompson, L., Reynolds, J., et al. (2015). Decision-making, impulsivity, and drug severity in co-occurring substance dependence and pathological gambling. *Drug and Alcohol Dependence*, 146(Jan (146)), e115–e116. Available from <https://www.infona.pl/resource/bwmeta1.element.elsevier-576f0ef7-22d4-3c8d-9a2d-7ce444a46218> Cited 14.12.15.
- Verdejo-García, A., Rivas-Pérez, C., Vilar-López, R., & Pérez-García, M. (2007). Strategic self-regulation, decision-making and emotion processing in poly-substance abusers in their first year of abstinence. *Drug and Alcohol Dependence*, 86(Jan (2–3)), 139–146. Available from <http://www.ncbi.nlm.nih.gov/pubmed/16806737> Cited 18.01.16.
- Villarejo, C., Jiménez-Murcia, S., Álvarez-Moya, E., Granero, R., Penelo, E., Treasure, J., et al. (2014). Loss of control over eating: A description of the eating disorder/obesity spectrum in women. *PubMed – NCBI. European Eating Disorders Review*, 22(1), 25–31. Available from <http://www.ncbi.nlm.nih.gov/pubmed/24338827> Cited 21.01.16.
- Volkow, N. D., & Baler, R. D. (2015). NOW vs LATER brain circuits: Implications for obesity and addiction. *Trends in Neurosciences*, 38, 345–352.
- Volkow, N. D., Wang, G.-J., Tomasi, D., & Baler, R. D. (2013). Pro v con reviews: Is food addictive? Obesity and addiction: Neurobiological overlaps. *Obesity Reviews*, 14, 2–18.

Wiehler, A., & Peters, J. (2015). Reward-based decision making in pathological gambling: The roles of risk and delay. *Neuroscience Research*, 90(Jan), 3–14. Available from <http://www.sciencedirect.com/science/article/pii/S0168010214002193> Cited 18.01.16.

Wu, M., Brockmeyer, T., Hartmann, M., Skunde, M., Herzog, W., & Friederich, H.-C. (2015). Reward-related decision making in eating and weight disorders: A systematic review and meta-analysis of the evidence from neuropsychological studies. *Neuroscience & Biobehavioral Reviews*, 61(Dec), 177–196. Available from <http://www.sciencedirect.com/science/article/pii/S0149763415302207> Cited 21.12.15.

Enduring changes in neuropsychological functioning in anorexia nervosa patients in full remission

Trevor Steward^{1,2}, Gemma Mestre-Bach^{1,2}, Zaida Agüera^{1,2}, Roser Granero^{2,3}, Isabel Sánchez-Díaz^{1,2}, Nadine Riesco^{1,2}, Susana Jiménez-Murcia^{1,2}, Fernando Fernández-Aranda^{1,2,*}

¹ Eating Disorders Unit, Department of Psychiatry, University Hospital Bellvitge-IDIBELL, Barcelona, Spain

² CIBEROBN, Instituto Salud Carlos III, Barcelona, Spain

³ Departament de Psicobiologia i Metodologia de les Ciències de la Salut, Universitat Autònoma de Barcelona, Barcelona, Spain

Introduction and objectives: Deficits in neuropsychological functioning have been repeatedly identified in patients with anorexia nervosa (AN). However, little is known on how cognitive functioning in AN patients evolves in response to treatment.

Method: AN patients ($n = 42$) completed a comprehensive set of neurocognitive (decision-making and selective attention) measures before admission (T0) to a 3-month day-hospital treatment program and at a 1-year follow-up (T1), and were compared to healthy controls ($n = 52$). BMI and eating disorder symptomatology were used as outcome variables.

Results: AN patients presented poorer performance on the Iowa Gambling Task (IGT) and the Stroop test at T0, compared to healthy controls ($p < .001$). AN patients in full remission ($n = 31$; 73.9%) at T1 significantly improved performance on the IGT ($p = 0.019$), and scores were not statistically different from HC scores ($p > 0.05$). Non-recovered patients and patients with partial remission ($n = 11$; 26.1%) did not improve IGT scores ($p > 0.05$). No changes were found in Stroop test scores in AN patients at T1 ($p > 0.05$).

Conclusions: AN patients in full remission showed marked improvements in decision making one year after beginning treatment, although significant impairments in other cognitive domains were still present.

Keywords: Neuropsychology; Eating and feeding disorders; Cognitive impairments; Longitudinal studies; Cognitive behavioral therapy; Choice behavior

Acknowledgment: Research partially supported by FIS PI14/00290.

Clinical and cognitive profiles in children with autism spectrum disorders: The experience of an specific educational center

J. Miralbell^{1,2,*}, E. Pradell², N. Farrés¹, B. Alcàcer¹, C. Castelló¹, T. Vaimberg¹, E. Arias-Pujol^{1,3}, M. Mestres¹

¹ Educative and therapeutic Center Carrilet S.C.C.L., Barcelona, Spain

² Clinical Psychology and Psychobiology Department, University of Barcelona, Spain

³ Psychology Faculty Blanquerna, Ramon Llull University, Barcelona, Spain

Introduction: Cognitive abilities in children with autism spectrum disorders (ASD) is one of the key factors related to school achievement, severity of clinical symptoms and adaptative level (Goldberg et al., 2006).

Objective: To investigate the clinical and medical profile of children with ASD according to their performance on cognitive tests.

Methods: 39 children with diagnosis of ASD attending C.E.E. Carrilet (Barcelona) were included. Cognitive performance was measured using both the WISC-IV or Weschler non-verbal (WNV) scales based on children's verbal abilities. Autism symptoms were assessed using the Autism Diagnostic Observation Schedule (ADOS-2). Medical comorbid pathologies and actual drug use were collected from their clinical history. Children were divided into groups based on the cognitive tests administrated and their clinical profiles were compared.

Results: Cognitive performance in children with ASD attending a specific educational center is heterogeneous, ranging from intellectual disability to talent. Three ASD profiles with different clinical characteristics were derived: (a) children assessed by WISC-IV: severe ASD symptoms, fluent language and no medical comorbidities nor medication; (b) children assessed with WNV: low-moderate ASD symptoms, no fluent language and no medical comorbidities; (c) children not assessed: moderate ASD symptoms, no language, and medical comorbidities and psychiatric medication.

Conclusion: Global and multi-disciplinary assessment in specific educational centers attending children with ASD is essential for planning individualized educative and therapeutic programs (Carrilet C.E.E, 2013; Viloca et al., 2012). Evaluations should include clinical, cognitive and medical aspects.

Keywords: Autism spectrum disorders; Cognitive assessment; Medical comorbidities; Clinical symptoms

References

- Goldberg M. (2006). Are the majority of children with autism mentally retarded? *Focus on Autism and Other Developmental Disabilities*, 21, 66–83.
- Centre educatiu i terapèutic Carrilet. (2013). *Comprensió i abordaje educativo y terapéutico del TEA*. Barcelona: Horsori Editorial.
- Viloca L. (2012). *El niño autista: detección, evolución y tratamiento*. Col·leccions Carrilet.