

Mortality Predictive Factors of a Clinical Cohort of Elderly Patients

R.J. Regal-Ramos,^a M.A. Salinero-Fort,^b and A.J. Cruz-Jentoft^c

Objective. To study the association between the main variables collected in the comprehensive geriatric assessment (CGA) and mortality, in a clinical cohort of elderly people referred from primary care, following standardised criteria, to a geriatric unit.

Design. Retrospective cohort study.

Setting. Outpatient department of a geriatric unit of a hospital in Madrid, Spain.

Participants. A total of 140 patients older than 65 years were followed up for 70 months.

Main measurements. We collected demographic, clinical, functional, and social variables during the CGA carried out by a multidisciplinary team. After 70 months we measured this cohort survival and we analysed the predictive factors for mortality using Cox hazard ratio analysis.

Results. Sixty three patients died after the 70 months of the study, and the survival median was 37 months. In the univariate analysis, age, male gender, diagnosed cancer, COPD, the Katz and Lawton indices, and the Mini-Mental State Examination (MMSE) score of 35 items, were significantly associated with mortality. In the multivariate analysis we found, as predictive factors for mortality: MMSE-35 (HR=0.965; 95% CI, 0.934-0.998; $P=0.037$); male gender (HR=2.75; 95% CI, 1.6-4.74; $P=0.001$); Katz score (HR=1.22; 95% CI, 1.04-1.43; $P=0.017$); Lawton score (HR=0.93; 95% CI, 0.82-1.07; $P=0.30$).

Conclusion. Cognitive impairment is a mortality predictive factor (HR=0.65), for each point less in the MMSE-35, we observed an increase in mortality risk of 3.5% (1-HR) at 70 months, after adjustment for Katz and Lawton index and gender.

Key words: Elderly. Predictive value of tests. Primary health care. Survival analysis.

Objetivo. Estudiar la asociación entre las principales variables recogidas en la valoración geriátrica exhaustiva (VGE) y la mortalidad en una cohorte clínica de ancianos remitidos desde atención primaria, según criterios protocolizados, a una unidad de geriatría.

Diseño. Estudio de cohorte clínica histórica.

Emplazamiento. Consulta de la unidad de valoración geriátrica de un hospital de Madrid.

Participantes. Un total de 140 pacientes > 65 años seguidos durante 70 meses.

Mediciones principales. Se recogieron variables demográficas, clínicas, funcionales y sociales durante la VGE realizada por un equipo multidisciplinario. Al cabo de 70 meses se valoró la supervivencia de esta cohorte y se analizaron los factores pronósticos de mortalidad mediante la técnica de riesgos proporcionales de Cox.

Resultados. El 45% ($n = 63$) de los pacientes había fallecido tras los 70 meses del estudio, con una mediana de seguimiento de 37 meses. En el análisis univariable, la edad, el sexo masculino, el diagnóstico de neoplasia o enfermedad pulmonar obstructiva crónica, los índices de Katz y de Lawton y el Mini-Examen Cognoscitivo (MEC-35) se asociaron significativamente con mortalidad. En el análisis multivariable se mantuvieron como variables pronósticas de mortalidad: el MEC-35 (*hazard ratio* [HR] = 0,965; intervalo de confianza [IC] del 95%, 0,934-0,998; $p = 0,037$), el sexo masculino (HR = 2,75; IC del 95%, 1,6-4,74; $p = 0,001$), el índice de Katz (HR = 1,22; IC del 95%, 1,04-1,43; $p = 0,017$) y el índice de Lawton (HR = 0,93; IC del 95%, 0,82-1,07; $p = 0,30$).

Conclusión. El deterioro cognitivo es un factor predictor de mortalidad (HR = 0,965), de manera que por cada punto de disminución del MEC-35 el riesgo de mortalidad a los 70 meses, ajustado por el sexo y los índices de Katz y Lawton, aumentará un 3,5% (1 - HR).

Palabras clave: Ancianos. Valor predictivo de tests. Atención primaria. Análisis de supervivencia.

Spanish version available at
www.atencionprimaria.com/94.758

A commentary follow
this article (pág. 487)

^aMar Báltico Health Centre, Área 4, IMSALUD, Madrid, Spain.
^bPrimary Care Training and Research Unit, Área 4, IMSALUD, Madrid, Spain.
^cRamón y Cajal Hospital, Área 4, IMSALUD, Madrid, Spain.

Correspondence:
M.A. Salinero Fort.
Unidad de Formación e Investigación de Atención Primaria.
Área 4. IMSALUD.
José Arcones Gil, 11. 28017 Madrid, España.
E-mail:
msalinero.gapm04@salud.madrid.org; masalinero@hotmail.com

Manuscript received March 23, 2004.
Manuscript accepted for publication January 21, 2005.

Introduction

The progressive increase in the population >65 years old¹ in our country is, from a public health perspective, a problem of great concern which requires social and health measures to be adopted. These measures must include screening for incapacities and medical, functional, psychiatric and social problems, with the objective of designing a therapeutic and care plan, as well as long term follow-up. This clinical-preventive approach was the idea of the first British geriatricians² and has been given many names: multidisciplinary geriatric assessment, integral or exhaustive geriatric assessment and comprehensive geriatric assessment (CGA), this latter one being the most prestigious. Five parameters are studied³: organic health or disease, mental function, functional situation, social support, and economic situation.

In short and medium term studies carried out on non-institutionalised elderly people in our country, the relationships between some of the assessments included in the CGA (Katz index and cognitive deterioration) and mortality have been highlighted.^{4,5} However, we have not found any studies where the relationship between these assessments, and others contained in the CGA, and the reduction in survival have been evaluated for more than 5 years. This deficiency should be rectified, since elderly patients take up 50% of the professional time of the primary care doctor,⁶ and it is the most suitable field for prevention and early diagnosis.


The objective of this study is to identify and prioritise the aspects of CGA, carried out in a hospital geriatric assessment unit, which may better predict the mortality of our elderly at 70 months, with the purpose of incorporating those aspects, which should enable the most vulnerable patients and those who might benefit more from health interventions to be detected, into the routine work of family doctors.

Patients and Method

A retrospective observational study of cohorts (historical cohort) has been carried out, which included all the elderly patients who were sent to the Geriatric Assessment Unit of the Ramon and Cajal Hospital during the year 1994, if they complied with the following established criteria: the first 3 and any of the other 9 of the points in Table 1.

The progress of all the patients was observed from 1994 to October 1999 (cut-off point for convenience), by using the hospital clinical records. Besides the time of survival in months, at the beginning of follow-up the following qualitative variables were collected using the CGA: gender, cardiac disease, blood pressure, diabetes mellitus, chronic obstructive pulmonary disease (COPD), diseases of the locomotor system which might involve a limitation for the patient, neurological diseases, auditory and ophthalmic diseases which might involve a limitation, neopla-

Material and methods



General Scheme of the Study

Historical cohort made up of 140 elderly patients with morbidity, referred to geriatrics department during 1 year (1994).

TABLE 1 Inclusion Criteria in the Clinical Cohort of Elderly People

Older than 65 years
Resident in Area IV of Madrid
Capacity to be able to come to the hospital by some means, to carry out the assessment
The study or management of the patient makes it difficult or impossible with the means available in primary care
Physical or mental deterioration of unknown cause after the initial assessment
Multiple chronic disease or invalidity, when the interactions between the diseases and capacity would be seriously difficult to manage
Use of more 5 medications, without being able to reduce this number in any previous attempt
Moderate/severe or progressive malnutrition, of unknown cause or where initial intervention has been unsuccessful
Incapacity to carry out the normal functions of the arms and/or legs, when they are considered rehabilitable
Urine or faecal incontinence, when easily treatable causes are ruled out (urinary infection, faecal impaction...)
Repeated falls (at least 3 in the last 6 months, or with serious consequences
Social problem which habitually interferes with the health workers

sias, number of medications consumed at the time of being given the geriatric assessment, depression (diagnosed according to DSM IV criteria), living alone, and use of social resources. Likewise, the quantitative variables collected at the beginning of the study were: age, Yesavage scale (this has the detection of depression as its aim and is specifically designed for the elderly patient: in our study we have used his short version of 15 questions) and the Mini Mental State Examination with 35 items (MMSE-35). The Katz and Lawton index variables, which are qualitative ordinals, were categorised as quantitative (for the Katz index: A=1, B=2, C=3, D=4, E=5, F=6, G=7, and for the Lawton: total independence, 8 points, decreasing to 0 points depending on the number of functional limitations), although, for the comparison of the mean results between deaths and survivors non-parametric tests were used (Mann-Whitney U test).

At the end of the follow-up period the data from the final outcome (morbidity and mortality) of these patients were collected by reviewing the primary care clinical histories, computerised records of the hospital and, lastly, by telephone calls to the patients or their families. This data was entered into a database where the variables which could identify the patient were separated, complying with the data protection guidelines.

Statistical Methods

A descriptive analysis of each of the study variables was carried out, summarising the qualitative nature using absolute and relative frequency tables, and the quantitative nature using the mean $\pm$ standard deviation.

For the comparisons, the χ^2 test was used for the qualitative variables and the Student *t* test for the quantitative ones.

With the aim of finding out the independent predictive variables of mortality after 70 months of follow-up, a Cox regression model was constructed, for which the hazard ratio assumption is previously verified using the visual observation of the Kaplan-Meier survival curves and log minus log graphs in the stratified models for each covariable. To select the variables included in the model, a bivariate analysis with all the variables collected in the CGA was performed, making time of survival the dependent variable and the rest covariables. The variables with a level of significance $\leq 20\%$ (male, diabetes mellitus, COPD, neoplasias, Katz, and Lawton indices) are included in the Cox maximum hazard ratio model, where we take the MMSE-35 as the main variable and the rest as control variables. The absence of collinearity of the variables which make up the maximum model is checked by applying a macro designed for this effect in those variables which may have correlation coefficients between them of <0.60 .

Using the backward inference method (maximum verisimilitude), controlled manually, the variables with a worse statistical significance are suppressed, provided that, owing to the beta coefficient of the main predictive variable not varying by more than 10%, they may not be considered as confounding. The best possible definitive model is finally obtained. All the data was analysed with the Windows SPSS 10.0 statistics package.

Results

In the almost 6 years of follow-up 63 patients died, 51 continued living, and 26 were lost at some time in the study (not able to contact them or their families and/or new records of their outcome were not noted in hospital or in their health centre). The median survival was 57

months and in month 70 the accumulated survival was 45.4%.

From the basal data of the population studied (Table 2) a high frequency of illness emerges, particularly in processes such as depression, cardiac disease, diabetes, and neurological disturbances. At the end of the follow-up period (Table 3) there were significant differences in mortality in male patients, diabetics, those with a Katz index of $> A$ (at least 1 limitation for activities basic for normal daily living [ABDL]) and Lawton ≤ 2 (dependency in at least 6 activities instrumental for daily living [AIDL]). On the other hand, the proportion of patients with an age greater than the mean (78 years) as well as those with a MMSE-35 score <22 had a higher tendency to die, but did not reach statistical significance ($P=.07$ and $P=.08$, respectively). The rest of the basic characteristics and outcomes of the 140 patients are described in Tables 3 and 4, respectively. It should be pointed out that patients who died had a significantly higher mean age

TABLE 2 Characteristics of the Patients Referred to the Geriatric Assessment Clinic*

No. of patients	140
Age, years, mean $\pm$ SD	78.4 $\pm$ 6.7
Male gender, %	31.2
Diseases, %	
Cardiac disease	42.5
High blood pressure	54.7
Diabetes	17.7
COPD	19.2
Osteo-articular diseases	39.2
Neurological diseases	29.5
Depression	39.3
Severe auditory and/or vision limitation	33
Neoplasias	10.5
Number of medications, mean $\pm$ SD	3.7 $\pm$ 2.0
Social situation and use of services, %	
Lives alone	16.5
Uses domiciliary resources	10.5
Uses non-domiciliary resources	8.6
Functional situations	
Dependent for any ABDL, %	45.3
Katz Index, mean $\pm$ SD	2.0 $\pm$ 1.9
Dependent for >6 AIDL, %	54.1
Lawton Index, mean $\pm$ SD	3.2 $\pm$ 3.0
Cognitive mini-examination, mean $\pm$ SD	21.7 $\pm$ 9.1
MMSE-35 $<22/35$, %	50.9

*SD indicates standard deviation; COPD, chronic obstructive pulmonary disease; AIDL, activities instrumental for daily living; ABDL, activities basic for daily living; MMSE-35, 35 item mini-examination.

TABLE 3
Differences Between Survivors and Deaths at 70 Months of Follow-up*

Characteristics	Alive	Dead	P
Age, years, mean	79.9	77.1	.01
Male gender, %	19.5%	42.9%	.00
Diseases, %			
Cardiac disease	42.1%	42.9%	NS
High blood pressure	57.1%	52.4%	NS
Diabetes	11.7%	23.8%	.05
COPD	13%	25.4%	.06
Osteo-articular diseases	40.3%	38.1%	NS
Neurological diseases	27.3%	31.7%	NS
Depression	39%	39.7%	NS
Severe auditory and/or vision limitation	31.2%	34.9%	NS
Neoplasias	6.7%	14.3%	NS
Number of medications, mean	3.7	3.6	NS
Social situation and use of services			
Lives alone	17.1%	15.9%	NS
Uses domiciliary resources	13.2%	7.9%	NS
Uses non-domiciliary resources	9.4%	7.9%	NS
Functional situations			
Dependent for any ABDL	35.1%	55.6%	.01
Katz Index, mean	1.6	2.4	.00
Dependent for >6 AIDL	41.6%	66.7%	.00
Lawton Index, mean	3.9	2.2	.00
Cognitive mini-examination, mean	23	20.2	.08
MMSE-35<22/35	43.1%	58.7%	.06

*COPD: chronic obstructive pulmonary disease; NS: non-significant; ABDL: activities basic for daily living; AIDL: activities instrumental for daily living; MMSE-35: 35 item mini-examination.

value and Katz index, and a significantly lower Lawton index than those who lived.

The non adjusted hazard ratio (HR) and the significance of the variables considered in the study are set out in Table 5. The neoplasias (HR=2.7), gender (HR=2.2), and COPD (HR=1.8) are highlighted as the variables with a stronger association with mortality. In the multivariate analysis (adjusted HR), the neoplasia and COPD variables lost their statistical significance and left the definitive model, which remained composed of the MMSE-35 variables, gender and the Katz index as significant variables, and the Lawton index as a confounding factor (as can be seen in Table 5). From looking at this table it is seen that, as regards the Katz index, for each decrease or loss of one activity basic for daily living, the death rate after 70 months of follow-up increases by 21.8%. Likewise, with the Lawton index, for each increase of one activity instrumental for daily living, the death rate decreases by 6.5%, which is the same as, for each loss of one activity instrumental for daily living, the death rate increases by 6.5% af-

ter 70 months of follow-up. Finally, the effect of an increase of one unit of the MMSE-35 decreases mortality by 3.5%, that is to say, a reduction of one unit of the MMSE-35 (a situation more common than an increase) implies an increase of 3.5% (1-0.965) in the mortality at 70 months, adjusted for gender and activities basic and instrumental for daily living.

We have noted a good correlation ($r=-0.625$; $P=.001$) between the scores obtained in the Katz and Lawton indices (the negative correlation coefficient is due to both ordinal scales being categorised in reverse order). Given that the correlation is not perfect, we preferred to include the assessment of dependence for the AIDL (Lawton) although there should already be dependence for the ABDL (Katz), with the aim of measuring complementary dimensions of a same problem. In fact, of the 22 patients who had total independence for AIDL, 9 had one limitation less for the ABVD, such as mild degrees of urinary incontinence or difficulty in getting in or out of the bath without help. On the other hand, both indices did not behave as collinear in the multivariate analysis and, besides, the Lawton behaves as a confounding variable, therefore it is obliged to remain in the final Cox regression model.

As a verification of our findings, Figure 1 shows the probability of survival of the patients during the 70 months of the study, and Figure 2 the survival stratified according to the MMSE-35 score (according to the following categories: score ≤ 16 , between 17 and 21, and ≥ 22), with a level of significance associated to the logarithmic ranges test of .0004.

Discussion

The high prevalence of illnesses such as cardiac disease, diabetes, depression, or neurological complaints can be explained by this being a very select population. On the other hand, the low proportion of elderly with oncology problems, which made up part of the cohort, may be due to the elderly who had cancer mainly being sent to palliative care units or were followed-up by the hospital oncology department. By this same reasoning, it has to be thought that the elderly with cancer in our cohort are those with a better short term prognosis, and this could explain its lack of association in the adjusted model.

There are discrepancies in the literature for⁷⁻¹⁷ and against^{18,19} the association between the severity of cognitive deterioration and survival. With this study we support the presence of this relationship in our area. As regards the tools for cognitive assessment, Abizanda⁴ used cut off points in the MMSE, Baldereschi et al¹² the diagnostic criteria of dementia from the DSM-III-R and other authors, such as Freid et al,²⁰ evaluation scales different from MMSE. We have preferred to take MMSE-35, given its high sensitivity and specificity and for being valid for the

TABLE 4 Non-Adjusted Hazard Ratio of the Variables Recorded From the Comprehensive Geriatric Assessment*

	Beta	HR [‡]	HR 95% CI		P
			Lower	Higher	
Age (quantitative)	0.038	1.039	1.003	1.076	.035
Male gender	0.823	2.277	1.380	3.757	.336
Hypertension	-0.243	0.784	0.478	1.286	.336
Cardiac disease	0.035	1.036	0.629	1.707	.889
Diabetes	0.463	1.588	0.889	2.838	.118
COPD	0.610	1.841	1.043	3.250	.035
Osteo-articular diseases	-0.108	0.897	0.539	1.493	.67
Neurological diseases	0.317	1.373	0.806	2.336	.243
Depression	0.024	1.024	0.618	1.697	.926
Yesavage (quantitative)	0.008	1.008	0.948	1.072	.797
Severe auditory and/or vision limitation	0.210	1.234	0.734	2.073	.427
Neoplasias	0.998	2.713	1.326	5.549	.006
Number of medications (quantitative)	-0.018	0.982	0.864	1.115	.778
Lives alone	-0.298	0.742	0.377	1.461	.389
Uses domiciliary resources	-0.439	0.645	0.259	1.608	.347
Uses non-domiciliary resources	-0.214	0.807	0.324	2.013	.646
Katz Index (quantitative)	0.252	1.287	1.134	1.1460	.000
Lawton index (quantitative)	-0.200	0.819	0.745	0.900	.000
MMSE-35 (quantitative)	-0.046	0.955	0.930	0.981	.001

*HR indicates hazard ratio; COPD, chronic obstructive pulmonary disease.

‡For the quantitative variables (age, number of medications, Yesavage, Katz and Lawton indices, MMSE-35), the HR corresponds to the increase of one unit.

††The HR must be interpreted as the times that it is quicker to die in the group exposed to the study phenomenon compared to the non-exposed group. Likewise, an interpretation similar to that of the relative risk is acceptable.

geriatric population in our environment.²¹ Likewise, we have categorised the MMSE-35 as a discrete quantitative variable with the objective of not losing information and quantifying the effect the small variations in scores on the scales had on mortality.

TABLE 5 HR Adjusted to the Variables Which Make up the Definitive Model*

	Beta	HR	HR 95% IC		P
			Lower	Higher	
MMSE-35 (quantitative)	-0.35	0.965	0.934	0.998	.037
Male gender	1.013	2.753	1.600	4.737	.000
Katz index (quantitative)	0.197	1.218†	1.036	1.432	.017
Lawton index (quantitative)	-0.67	0.935‡	0.818	1.070	.329

*HR indicates hazard ratio.

†Must be interpreted as: each decrease or loss of one activity basic for daily living increases the death rate by 21.8% after 70 months follow-up.

‡Must be interpreted as: each increase of one activity instrumental for daily living decreases the death rate by 6.5% or, which is the same as, for each loss of one activity instrumental to daily living increases the death rate by 6.5% after 70 months of follow-up.

As regards the marked significance of physical dependence on the mortality, our results agree with those obtained by Gambassi et al,¹⁸ Narain et al,²² and other authors.^{4,20,23} To be male has been shown to be an independent predictive factor, with a mortality risk of approximately twice that of women, in different studies,^{13,20,22,24} and as was obtained in ours (HR=2.75).

Although age may appear as a potential confounding variable on being associated with the risk of death and with the worsening of the MMSE-35 score,^{7,12,18,24} it has not behaved as such, as has also happened in other studies.^{22,25,26} However, other authors found it necessary to adjust for age.²⁷ Despite our sample having been consecutive patients attending the Geriatric Assessment Unit, we believe that it can be considered as a non-biased clinical cohort since, on the one hand, it has been made up of the whole population of patients sent to this unit and, on the other hand, the period of inclusion in the study had been 1 year, therefore it has avoided the seasonal nature of certain diseases. For this reason, it should be treated as a non-representative sample of the group of elderly people who live in

the area, even though those who present with morbidity and morbidity have to be sent to the Geriatric Assessment Unit according to the criteria in Table 1.

In conclusion, it is observed that, after adjusting for gender, the ABDL (Katz index) and the AIDL (Lawton index), for each one point increase in the MMSE-35 it will decrease the probability of dying by 3.5% (HR=0.965) or, said another way, given that it is going to be difficult for an elderly person to improve his/her MMSE-35 score with time, each point decrease in the MMSE-35, will increase the risk of mortality at 70 months by 3.5%.

To avoid inaccuracies in the determination of the MMSE-35 it is recommended, as was done in our study, that the assessor who carries out the MMSE-35 receives previous training and the person is not chan-




FIGURE 1 Kaplan-Meier survival graph.




FIGURE 2 Kaplan-Meier survival graph according to the MMSE-35 values (categorised in: ≤16; 17-21, and ≥22 points). Logarithmic ranges: 15; $P=.0004$.

ged at any time. However, we are conscious of the possibility that in normal clinical practice a decrease of one point of the MMSE-35 may be due test-retest variability or to variations occurred due to a change of the assessor, more

Discussion Key points



What Is Known About the Subject

- There are no studies referenced in the major literature data bases, where the medium or long term total or partial efficacy of the comprehensive geriatric assessment is evaluated.
- Male gender is an independent predictive factor of mortality (RR=2) in the elderly population.
- There are discrepancies in the literature on the association between cognitive deterioration and mortality.

What This Study Contributes

- After male gender, the deterioration in the activities basic for daily living is the most significant predictive factor in the elderly population.
- Minimal variations in the MMSE-35 predict a significant increase in mortality, adjusted for gender and in activities basic and instrumental for daily living.
- Periodic cognitive assessment of the elderly population with morbidity seen in primary care can be a first line tool to identify the more vulnerable patients.

than true changes in cognitive function. Therefore, it would be more interesting to find out the effect on the mortality of changes of 3, 4, or 5 points in the MMSE-35. To calculate it only requires multiplying the MMSE-35 beta coefficient by the appropriate number (3, 4, or 5) and exponentiate it. The result of this simple operation would tell us that a decrease of 3 points increases the risk of mortality by 10% (HR=0.90), of 4 points by 13% (HR=0.87), and 5 points up to 16% (HR=0.84).

Therefore, small or medium variations in the MMSE-5 predict a significant increase in mortality. For this reason we believe that the cognitive assessment of elderly patients with morbidity seen in primary care can constitute a tool of the first order to identify the most vulnerable elderly people and can at the same time help us to plan our clinics, the needs of referral and resources and, in short, to improve the quality of care for our patients.

However, other studies with similar characteristics are necessary in the field of primary care which would give external validity to our findings and to determine what

would be the most appropriate intervals to carry out the cognitive assessment of our elderly.

References

1. La tercera edad en España: aspectos cuantitativos. Proyecciones de la población española de 60 años para el período 1986-2010. Madrid: Ministerio de Asuntos Sociales; 1991.
2. Warren MW. Care of chronic sick. A case for treating the chronic sick in blocks in a general hospital. *BMJ*. 1943;2:823-4.
3. Williams ME. Geriatric assessment. *Ann Intern Med*. 1986;104:720-21.
4. Stuck AE, Siu AL, Wieland GD, Adams J, Rubenstein LZ. Comprehensive geriatric assessment: a meta-analysis of controlled trials. *Lancet*. 1993;342:1032-6.
5. Cortés A, Méndez-Bonito E, Koutsourais R, Utrilla J, Macías M, Casado MA, et al. ¿Cuál es la prevalencia de ancianos de alto riesgo en atención primaria? *Aten Primaria*. 1996;18:327-30.
6. Ostbye T, Steenhuis R, Wolfson C, Walton R, Hill G. Predictors of five-year mortality in older Canadians: the Canadian Study of Health and Aging. *J Am Geriatric Soc*. 1999;47:1249-54.
7. Walsh JS, Welch HG, Larson EB. Survival of outpatients with Alzheimer-type dementia. *Ann Intern Med*. 1990;113:429-34.
8. Perls TT, Morris JN, Ooi WL, Lipsitz LA. The relationship between age, gender and cognitive performance in the very old: the effect of selective survival. *J Am Geriatr Soc*. 1993;41:1193-01.
9. Ueki A, Shinjo H, Shimode H, Nakajima T, Morita Y. Factors associated with mortality in patients with early-onset Alzheimer's disease: a five-year longitudinal study. *Int J Geriatr Psychiatry*. 2001;16:810-5.
10. Brodaty H, McGilchrist C, Harris L, Peters KE. Time until institutionalization and death in patients with dementia. Role of caregiver training and risk factors. *Arch Neurol*. 1993;50:643-50.
11. Claus JJ, van Gool WA, Teunisse S, Walstra GJ, Kwa VI, Hijdra A, et al. Predicting survival in patients with early Alzheimer's disease. *Dement Geriatr Cogn Disord*. 1998;9:284-93.
12. Baldereschi M, dDiCarlo A, Maggi S, Grigoletto F, Scarlato G, Amaducci L, et al. Dementia is a major predictor of death among the Italian elderly. ILSA Working Group. Italian Longitudinal Study on Aging. *Neurology*. 1999;52:709-13.
13. Desmond DW, Moroney JT, Sano M, Stern Y. Mortality in patients with dementia after ischemic stroke. *Neurology*. 2002;59:537-43.
14. Hassing LB, Johansson B, Berg S, Nilsson SE, Pedersen NL, Hofer SM, et al. Terminal decline and markers of cerebral and cardiovascular disease: findings from a longitudinal study of the oldest old. *J Gerontol B Psychol Sci Soc Sci*. 2002;57:268-76.
15. Kelman HR, Thomas C, Kennedy GJ, Cheng J. Cognitive impairment and mortality in older community residents. *Am J Public Health*. 1994;84:1255-60.
16. Bassuk SS, Wypij D, Berkman LF. Cognitive impairment and mortality in the community-dwelling elderly. *Am J Epidemiol*. 2000;151:676-88.
17. Gambassi G, Landi F, Lapane KL, Sgadari A, Mor V, Bernabei R. Predictors of mortality in patients with Alzheimer's disease living in nursing homes. *J Neurol Neurosurg Psychiatry*. 1999;67:59-65.
18. Bracco L, Gallato R, Grigoletto F, Lippi A, Lepore V, Bino G, et al. Factors affecting course and survival in Alzheimer's disease. A 9-year longitudinal study. *Arch Neurol*. 1994;51:1213-9.
19. Abizanda P, Luengo C, López-Torres J, Sánchez P, Romero L, Fernández C. Predictores de mortalidad, deterioro funcional e ingreso hospitalario en una muestra de ancianos residentes en la comunidad. *Rev Esp Geriatr Gerontol*. 1998;33:219-25.
20. Freid LP, Kronmal RA, Newman AB, Bild DE, Mittmark MB, Polak JF, et al. Risk factors for 5-year mortality in older adults. The cardiovascular health study. *JAMA*. 1998;279:585-92.
21. Lobo A, Saz P, Marcos G, Díaz JL, de la Cámara C, Ventura T, et al. Revalidación y normalización del Mini-Examen Cognoscitivo (primera versión en castellano del Mini-Mental Status Examination) en la población general geriátrica. *Med Clin (Barc)*. 1999;112:767-74.
22. Narain P, Rubenstein LZ, Wieland GD, Rosbrook B, Strome LS, Pietruszka, et al. Predictors of immediate and 6-month outcomes in hospitalized elderly patients. *J Am Geriatr Soc*. 1988;36:775-83.
23. Incalzi AR, Capparella O, Gemma A, Porcedda P, Raccis G, Sommella L, et al. A simple method of recognizing geriatric patients at risk for death and disability. *J Am Geriatr Soc*. 1992;40:34-8.
24. Palomo L, Gervas J. Mortalidad a los 2 años en pacientes crónicos confinados en el domicilio. *Aten Primaria*. 2000;25:106-14.
25. Maxson PJ, Berg WS, Mc Clearn G. Multidimensional patterns of aging in 70-years-old: survival differences. *J Aging Health*. 1996;3:320-33.
26. Zanonchi M, Ponzetto M, Neirotti M, Maero B, Francisetti F, Spada S, et al. Predictive variables of mortality in elderly patients after hospitalization. Significance of multidimensional evaluation. *Recent Prog Med*. 2001;92:184-8.
27. Bowen JD, Malter AD, Sheppard L, Kukull WA, McCormick WC, Teri L, et al. Predictors of mortality in patients diagnosed with probable Alzheimer's disease. *Neurology*. 1996;47:433-9.

COMMENTARY

Impact of Functional and Cognitive State on the Mortality of Elderly People Living in the Community

M.A. Benítez-Rosario

Member of the Working Group for Care of the Elderly of the Spanish Society of Family and Community Medicine and the Canary Islands Society of Family and Community Medicine, Palliative Care Unit, La Candelaria University Hospital, Tenerife, Spain

Primary care is the level of health care which a greater number of elderly people attend. Its function is to carry out preventive activities to achieve a healthy old age, the early detection of diseases and incapacities, to be able to reverse or stop their progression, and care at the end of life which favours a dignified death.¹ The availability of valid tools which enable the detection, monitoring and prediction of the evolution of their state of health is essential to be able to carry out these functions.

The work of Regal et al² in this issue of primary care is particularly interesting in this context, which is the search for tools or diagnostic tests which give information on what may be taking place in an elderly person, depending on their basal condition. The tool evaluated is the comprehensive geriatric assessment, considering the predictive power of mortality in its different components. Their results show that the functional situation of the elderly person and the errors in the Mini-Mental State Examination (MMSE) are the main independent predictors of mortality at 70 months of follow-up.

The functional situation of the elderly is a major determining factor of their capacity to live in the community, their quality of life, their recovery capacity and their mortality in the short term when they are admitted to hospital with serious illnesses. A basic physical situation, expressed as the absence of incapacity for the performance of activities for daily living, is a marker of the “strength” of the patient to confront external problems. And this effect is independent of age. On the other hand, the functional incapacity appears to behave as an indicator of a situation of general debility which, independent of the morbidity, is a predictor of mortality, in the medium term, in people who live in the community. The results of Regal et al² corroborate previous observations originating from another longitudinal study³ carried out in our country and with a cohort of elderly people with a different state of health. The repeated observation confirms that it is not a casual finding. The consequences of these observations are obvious: it must be

Key Points

- The functional situation of elderly people who live in the community is a predictor of mortality in the medium short-medium term.
- Primary care must strengthen its role in promoting a healthy old age and the early detection of incapacities to try and reverse them.
- The significance of cognitive deterioration detected by the MMSE in the mortality in the elderly should be made clear in future studies.

acted on, to delay, where possible, the appearance of incapacities. And for this reason, the activity of primary care, as a promoter of a healthy old age and the early detection and intervention of functional disturbances, is fundamental.

The studies available, however, do not offer a prediction of mortality in the short term, as this is also important in the care of incapacitated elderly people. The diagnostic and therapeutic decisions when faced with illnesses must be based on the knowledge of the possibilities of one result or another depending on the functional situation of the elderly person. Other investigations, centred on patients with deterioration of two or more basic functions of daily living, are required to find out the impact of these deficiencies in the mortality at 3, 6, or 12 months. Also, in these studies it should be evaluated whether the predictive capacity of mortality in the very short term of a functional problem continues to be independent of the type of illness. The results of Regal et al² agree with those obtained by other authors^{4,5} as regards the influence of cognitive deterioration and, in particular, Alzheimer-type dementia

(ATD) on the mortality in the short-medium term in elderly people. In the study by Regal et al², however, it highlights the predictive capacity independent of the MMSE scores when these represent “losses” as regards the normal score. The interpretation of these findings pose some difficulty. The robustness of the prediction contrasts with the limitations of the tool and the interpretation of their results. The influence of gender, cultural level, and training for re-examining on the diagnostic performance of the MMSE and its variations are known. These factors would change the predictive capacity, in the same person, only by obtaining more or less points in the MMSE. It would be more logical to consider that the MMSE scores are really predictors as regards pointing out particular grades of severity in cognitive deterioration, and ATD, which are clearly associated with mortality in the medium to long term of the elderly. The results obtained by Regal et al,² however, lead to the interesting observation in that minimal cognitive changes, expressed as subtle losses of points in the MMSE normal range, can already have a predictive value of mortality at 70 months follow-up. It could be that the short neurophysical examination which the MMSE provides, may have the capacity to detect variations, considered up until now within normal, which show up a situation of general debility which determines a poor prognosis. It is known that the central nervous system is particularly sensitive in the elderly and in its function the repercussions of inadequate function of other organs are expressed. The MMSE could then behave as a “major” variable in which the influences of debilitating diseases are expressed in the mortality. The results should

also help when considering the interpretations of what is the change and the normality in the MMSE scores.

The investigation by Regal et al² validates the previous opinions on the importance of the functional diagnosis in the elderly and the early diagnosis of dementia in primary care. However, it should be remembered that there is no suitable evidence available which guarantees the generalised use of the comprehensive geriatric assessment in primary care.

References

1. Benítez del Rosario MA. Atención al anciano. In: Martín Zurro A, Cano Pérez JF, editors. Atención primaria. Concepto, organización y práctica clínica. Madrid: Elsevier; 2003. p. 1642-70.
2. Regal Ramos RJ, Cruz Jentoft AJ, Salinero Fort MA. Factores predictores de mortalidad de una cohorte clínica de pacientes ancianos. *Aten Primaria*. 2005;36(9):480-6.
3. Benítez-Rosario MA, Hernández-Estevez P, Aguirre-Jaime A, González-Freire G, Asensio-Fraile A. Functional status and mortality in community-dwelling older people. *J Am Geriatr Soc*. 2001;49:1009-10.
4. Larson EB, Shadlen MF, Wang L, McCormick WC, Bowen JD, Teri L, et al. Survival after initial diagnosis of Alzheimer disease. *Ann Intern Med*. 2004;140:501-9.
5. Porock D, Parker Oliver D, Zweig S, Rantz M, Mehr D, Madsen R, et al. Predicting death in the nursing home: development and validation of the 6-month minimum data set mortality risk index. *J Gerontol*. 2005;60:491-8.
6. Gorroñoigoitia Iturbe A, Martín Lesende I. Valoración geriátrica global en atención primaria. In: Grupo de Trabajo de Atención al Mayor de la semFYC. Atención a las personas mayores. Barcelona: semFYC; 2004. p. 45-52.