

Case report

Endovascular approach for non-thrombotic pulmonary embolism due to metal coil migration after treatment of pelvic varicose vein



Luis Fernando García^a, Nicole Moreno^a, Francisco Arroyo^a, Camilo Bautista^b,
Luis Felipe Cabrera-Vargas^{b,c}, Ivan David Lozada-Martinez^{c,d,*}

^a Department of Vascular and Endovascular Surgery, Colsubsidio Clinic – Hospital Militar Central, Universidad Militar Nueva Granada, Bogotá, Colombia

^b Vascular and Endovascular Surgery Fellowship, Department of Vascular and Endovascular Surgery, Hospital Militar Central, Universidad Militar Nueva Granada, Bogotá, Colombia

^c Medical and Surgical Research Center, Future Surgeons Chapter, Colombian Surgery Association, Cartagena, Colombia

^d Grupo Prometheus y Biomedicina Aplicada a las Ciencias Clínicas, School of Medicine, Universidad de Cartagena, Cartagena, Colombia

ARTICLE INFO

Article history:

Received 11 January 2022

Accepted 1 February 2022

Available online 4 March 2022

Keywords:

Pulmonary embolism

Endovascular procedures

Varicose veins

Foreign-body migration

ABSTRACT

Non-thrombotic pulmonary embolism is a partial or complete occlusion of the pulmonary vasculature by miscellaneous types of materials that includes neoplasm, infective agents, fat, amniotic fluid, gases and foreign materials. The literature reporting the incidence of coil migration after pelvic embolization and concomitant acute pulmonary embolism is scarce. We report a case of coil migration after pelvic embolization and symptomatic acute pulmonary embolism treated with endovascular extraction and standard anticoagulation.

© 2022 Sociedad Española de Cirugía Cardiovascular y Endovascular. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Abordaje endovascular para la embolia pulmonar no trombótica debido a migración de coil después del tratamiento de las várices pélvicas

RESUMEN

La embolia pulmonar no trombótica es una oclusión parcial o completa de la vasculatura pulmonar por diversos tipos de materiales que incluyen neoplasias, agentes infecciosos, grasa, líquido amniótico, gases y materiales extraños. La literatura que ha reportado de la incidencia de la migración de coil después de la embolización pélvica con embolia pulmonar aguda concomitante, es escasa. Presentamos un caso de migración de coil tras embolización pélvica y embolia pulmonar aguda sintomática tratada con extracción endovascular y anticoagulación estándar.

© 2022 Sociedad Española de Cirugía Cardiovascular y Endovascular. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Palabras clave:

Embolia pulmonar

Procedimientos endovasculares

Venas varicosas

Migración de cuerpo extraño

Non-thrombotic pulmonary embolism (NTPE) is a partial or complete occlusion of the pulmonary vasculature by miscellaneous types of materials that includes neoplasm, infective agents, fat, amniotic fluid, gases and foreign materials. NTPE can cause local and systemic inflammation causing endothelial and pulmonary parenchymal injury. This entity may present as completely asymptomatic, with nonspecific symptoms or even cause sudden death. The literature reporting the incidence of coil migration after pelvic embolization and concomitant acute pulmonary embolism is scarce. Most cases present as incidental radiological findings in asymptomatic patients.¹

We report a case of coil migration after pelvic embolization and symptomatic acute pulmonary embolism treated with

endovascular extraction and standard anticoagulation. The patient has agreed to the publication of images and details related to his case.

A 53-year-old female with congestive pelvic syndrome due to pelvic varicose vein underwent pelvic veins embolization using 2 coils (20 mm × 50 cm) (Concerto; Medtronic Inc., Minneapolis, USA) for the right gonadal vein. She presented to the emergency department 30 days later with mild chest discomfort. Scout image from computed tomography (CT) pulmonary angiogram showed completely displaced coil nest in the chest. A 12-Fr sheath was placed in the right common femoral vein and pulmonary angiogram was performed, demonstrating occluded right lower lobe segmental arteries (Figs. 1 and 2). A 5-Fr multipurpose catheter supported by a 0.025 guidewire was advanced into the main pulmonary artery. The coil nest was snared with a 25-mm gooseneck snare (Fig. 3) and retracted into the inferior vena cava. The coil nest was then pulled down into the left external iliac vein and removed through the

* Corresponding author.

E-mail address: ilozadam@unicartagena.edu.co (I.D. Lozada-Martinez).




Fig. 1. Angiogram showed completely displaced coil nest in the chest.




Fig. 2. Pulmonary angiogram, demonstrate occluded right lower lobe segmental arteries.




Fig. 3. The coil nest was snared with a 25-mm gooseneck snare.




Fig. 4. The coil nest was then pulled down into the left external iliac vein.



Fig. 5. Final coil nest extraction.

sheath in two extractions procedures repeating the same sequence (Figs. 4 and 5). The patient was discharged home the next day on a therapeutic dose of enoxaparin. She remained asymptomatic at 3-month clinical follow-up visit.

Complications after pelvic embolization are rare.² Coil migration can occur after pelvic varicose vein embolization and cause a NTPE. The majority of reported cases in the literature are asymptomatic and no need for additional endovascular or open treatment. The chest X-ray may show a linear density at an unusual anatomic location, but CT is the gold standard to confirm the position of the foreign body.^{1,2} The pattern presented in chest CT will

depend on the type of embolization. Disperse radiopaque/metallic small densities, centrilobular ground glass opacities, and micronodules/miliary nodules, as well as fibrotic/consolidative opacities, may indicate NTPE.¹ If the patient develops persistent symptoms without improvement after medical treatment with systemic anticoagulation like the case reported by Shashi et al.,² endovascular coil extraction must be the first line approach. In 2018, Leon et al.¹ reported a case of symptomatic coil NTPE treated only with anticoagulation that had symptoms improvement during the clinical follow up. A case report published by D'Amato et al.³ in 2016, presented a patient with an incidental radiological finding of a nodular image of metal density in the left hemithorax. Chest CT scan showed a foreign body consistent with a coil in the left lower lobe segmental branch. The patient was asymptomatic, so she was treated conservatively with clinical and image follow up.³ Cases of coil migration have been described in men as well. In 1993 Moriel et al.⁴ published a case report of a man with an asymptomatic pulmonary migration of a coil after a pelvic embolization due to the treatment of erectile dysfunction.⁴ There are no reported cases of patients with coil NTPE that needed an open surgical approach.

In conclusion, the non-thrombotic coils pulmonary embolism after the endovascular treatment of congestive pelvic syndrome is a rare complication. However, it should be considered as a differential diagnosis in patients with acute onset dyspnea after pelvic varicose vein embolization. Large studies are needed to define the real incidence the best treatment approach to this entity. This brief report contributes to the world literature on the presentation, management and outcome of this complication, which is a rare occurrence.

Ethical considerations

The ethics committee of the Hospital Militar Central approved this case report. Constant of the use of medical data is obtained verbally and in written form from the patient.

Financial disclosures

None.

Conflict of interest

The authors declare that they have no conflicts of interest.

Acknowledgments

None.

References

1. León FX, Alba B, Arribas A, García R, García I, Quezada CA. A rare case of pulmonary embolism due to endovascular coil embolization and review of literature. *Int J Respir Pulm Med*. 2018;5:094.
2. Shashi KK, Chaudry G, Alomari A, Chewning R. Massive coil nest migration: endovascular retrieval. *J Vasc Interv Radiol*. 2019;30:1610–1.
3. D'Amato R, Figueira Gonçalves JM, Palmero Tejera JM. Embolismo pulmonar por migración de coil metálico tras tratamiento de varices pélvicas. *Arch Bronconeumol*. 2017;53:72.
4. Moriel EZ, Mehringer CM, Schwartz M, Rajfer J. Pulmonary migration of coils inserted for treatment of erectile dysfunction caused by venous leakage. *J Urol*. 1993;149 Pt 2:1316–8.



BIOMED



unidix

Especialistas en cirugía cardiovascular

desde 1977 al cuidado de tu salud



91 803 28 02



info@biomed.es