



## Image of the month

## Juvenile polyp

## Pólipo juvenil



Sara Sofia S. Rodrigues<sup>a,\*</sup>, Daniela Ester Ribeiro<sup>b</sup>, Miguel Costa<sup>a</sup>, Marta Tavares<sup>c</sup>

<sup>a</sup> Pediatric Department, Centro Hospitalar de Entre Douro e Vouga, Santa Maria da Feira, Portugal

<sup>b</sup> Pediatric Department, Centro Hospitalar de Baixo Vouga

<sup>c</sup> Pediatric Department, Centro Materno Infantil do Norte, Porto, Portugal



Fig. 1



Fig. 2

A 5-year-old boy presented with an 8-months history of painless rectal bleeding. Parents reported an occasional retractable mass in the anal region (Fig. 1). Colonoscopy showed a 2.2 cm pedunculated polyp in the distal rectum (Fig. 2). Polypectomy was performed. Histological analysis revealed a juvenile polyp without dysplasia. The child was discharged with no recurrent episode of rectal bleeding during follow-up, therefore, not needing further investigation.

Juvenile polyps are benign hamartomas, that typically occur between the ages of 2 and 10 years. Patients usually present with painless rectal bleeding and polyps located in the rectum may present as prolapsed tissue.

\* Corresponding author.

E-mail address: [sara.rodrigues1906@gmail.com](mailto:sara.rodrigues1906@gmail.com) (S.S.S. Rodrigues).

<https://doi.org/10.1016/j.ciresp.2023.08.005>

0009-739X/© 2023 AEC. Published by Elsevier España, S.L.U. All rights reserved.