

- Cancer. 2013;133:1164–72. <http://dx.doi.org/10.1002/ijc.28102>. PMID: 23404769.
3. Palefsky JM, Holly EA, Efird JT, da Costa M, Jay N, Berry J, et al. Anal intraepithelial neoplasia in the highly active antiretroviral therapy era among HIV-positive men who have sex with men. *AIDS*. 2005;19:1407–14. <http://dx.doi.org/10.1097/01.aids.0000181012.62385.4a>. PMID: 16103772.
 4. De Pokomandy A, Rouleau D, Ghattas G, Vézina S, Côté P, Macleod J, et al., HIPVIRG Study Group. Prevalence, clearance, and incidence of anal human papillomavirus infection in HIV-infected men: The HIPVIRG cohort study. *J Infect Dis*. 2009;199:965–73. <http://dx.doi.org/10.1086/597207>. PMID: 19239366.
 5. Sendagorta E, Herranz P, Guadalajara H, Zamora FX. Detección precoz de la neoplasia intraepitelial anal en pacientes de alto riesgo. *Actas Dermosifiliogr*. 2011;102:757–65. <http://dx.doi.org/10.1016/j.ad.2011.01.005>. PMID: 21764027.
 6. Fernández Serrano JL, Castro Santiago MJ, Fernández Morillo A, Bazan Hinojo C, Casado Maestre MD, Sánchez Ramírez M, et al. Lesiones intraepiteliales anales: opciones de ‘screening’ y tratamiento. *Cir Andal*. 2013;24:12–7.
 7. Cranston RD, Baker JR, Liu Y, Wang L, Elishaev E, Ho KS. Topical application of trichloroacetic acid is efficacious for the treatment of internal anal high-grade squamous intraepithelial lesions in HIV-positive men. *Sex Transm Dis*. 2014;41:420–6. <http://dx.doi.org/10.1097/OLQ.0000000000000145>. PMID: 24922100.
 8. Palefsky JM, Lee JY, Jay N, Goldstone SE, Darragh TM, Dunlevy HA, et al., ANCHOR Investigators Group. Treatment of anal high-grade squamous intraepithelial lesions to prevent anal cancer. *N Engl J Med*. 2022;386:2273–82. <http://dx.doi.org/10.1056/NEJMoa221048>. PMID: 35704479; PMCID: PMC9717677.
- Javier Gómez Sánchez*, Marisol Zurita Saavedra,
Rocio Forneiro Pérez y Benito Mirón Pozo
- Cirugía General y del Aparato Digestivo, Hospital Clínico San Cecilio, Granada, España*
- * Autor para correspondencia.
Correo electrónico: javiergomezsanchez.jg@gmail.com
(J. Gómez Sánchez).
- <https://doi.org/10.1016/j.cireng.2023.09.004>
0009-739X/
© 2023 AEC. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Laparoscopic and robotic distal pancreatectomy: Outcomes and the need for patient-centred evaluation



Pancreatectomía distal laparoscópica y robótica: resultados y necesidad de una evaluación centrada en el paciente

To the Editor,

Recently, we read with great interest the work by Alvarez et al., “Laparoscopic and robotic distal pancreatectomy: the choice and the future”. We congratulate the authors for conducting a prospective study at tertiary hospitals, which analysed the results of distal pancreatectomies with laparoscopic and robotic approaches and presented the outcomes and complications of these procedures, helping us understand the safety and efficacy of each. Information pertaining to the conversion of surgery, from robotic to laparoscopic or from laparoscopy to laparotomy, has also been elucidated. The similarity of the patients involved in the study in both groups in terms of their age, sex, BMI and ASA physical status classification reduces bias, at the same time enhancing the reliability and generalisability of the study.¹

Though postoperative pancreatic fistula was assessed as a complication in this study, new-onset diabetes (which has incidence rates varying up to 50 percent after distal pancreatectomy) can be assessed by comparing long-term outcomes in future studies.^{2,3} Though the study’s primary focus was on clinical outcomes, a more comprehensive understanding of

the patients’ total well-being would have been possible by assessing subjective patient-oriented outcomes, such as postoperative pain level, which gives us insight to how the patient feels after DP. Any surgical intervention’s overall success depends on a number of important factors, including functional recovery, dietary modifications, and long-term effects on the patient’s daily activities and lifestyle choices. A more patient-centred view of efficacy can be provided by using patient-reported outcome measures, such as satisfaction levels and functional recovery. Lack of data pertaining to other short-term outcomes, including ileus, respiratory failure, pulmonary embolism, acute coronary syndrome, stroke, acute renal failure, urinary tract infection and sepsis, is yet another shortcoming.⁴

RDP, in comparison to traditional DP techniques for treating disorders of the pancreas body and tail, appears to have clinical and oncological advantages. However, it may involve a lengthier operation and learning curve according to a meta-analysis, in contrast to the shorter learning curve mentioned by Alvarez et al.⁵ It is noteworthy that the study’s stated non-significant difference in the length of the two

operations is, in fact, significant in the multicenter analysis conducted by Lof et al. with a much larger sample size that found a similar difference in procedure durations.⁶

In conclusion, encouraging further research in the field of minimally invasive procedures, exploring novel surgical procedures, and providing long-term follow-up while addressing the limitations mentioned in the study can pave the way for improved outcomes, enhancing patient care.

Statement and declaration

The authors have no relevant financial or non-financial interests to disclose.

REFERENCES

1. Espin Alvarez F, García-Domingo MI, Cremades Pérez M, et al. Laparoscopic and robotic distal pancreatectomy: the choice and the future. *Cir Esp (Engl Ed)*. 2023;101(11):765-71. <http://dx.doi.org/10.1016/j.cireng.2023.04.017>.
2. Tariq M, Jajja MR, Maxwell. et al. Diabetes development after distal pancreatectomy: results of a 10 year series. *HPB*. 2020;22(7):1034-41. <http://dx.doi.org/10.1016/j.hpb.2019.10.2440>.
3. De Bruijn KM, van Eijck CH. New-onset diabetes after distal pancreatectomy: a systematic review. *Ann Surg*. 2015;261(5):854-61. <http://dx.doi.org/10.1097/SLA.0000000000000819>.

4. Ota M, Asakuma M, Taniguchi K, et al. Short-term outcomes of laparoscopic and open distal pancreatectomy using propensity score analysis: a real-world retrospective cohort study. *Ann Surg*. 2023;278(4):e805-11. <http://dx.doi.org/10.1097/SLA.0000000000005758>.
5. Lyu Y, Cheng Y, Wang B, Zhao S, Chen L. Comparison of 3 minimally invasive methods versus open distal pancreatectomy: a systematic review and network meta-analysis. *Surg Laparosc Endosc Percutan Tech*. 2020;31(1):104-12. <http://dx.doi.org/10.1097/SLE.0000000000000846>. Published 2020 Sep 2.
6. Lof S, van der Heijde N, Abuawwad M, et al. Robotic versus laparoscopic distal pancreatectomy: multicentre analysis. *Br J Surg*. 2021;108(2):188-95. <http://dx.doi.org/10.1093/bjs/znaa039>.

Adwaith Krishna Surendran^a, adwaithkrishnas600@gmail.com, Srisanjith Girish^a, Chellappa Vijayakumar^b

^aMedical Student, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Puducherry, India

^bDepartment of Surgery, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Puducherry, India

*Corresponding author. (A.K. Surendran).

<https://doi.org/10.1016/j.cireng.2024.01.011>
0009-739X/

© 2023 AEC. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Laparoscopic and robotic distal pancreatectomy: the choice and the future



First of all, the authors would like to thank you for your comments and the analysis of the article.

As it highlights, the implementation or incorporation of the robotic platform must follow the surgical principles and must demonstrate its effectiveness, efficiency, and safety, so we believe and so we raise it when analyzing our results.^{1,2}

In response to the questions raised. We have not analyzed the degree of induced diabetes after distal pancreatectomy. Our study assesses short-term outcomes. We consider your comment and keep it in mind for future analysis.

Regarding the perception and satisfaction of the patients, as well as the functional recovery. This analysis is a departure from the main objectives of the study, and would in itself deserve to be presented in a single article. Nevertheless, we consider it elementary and have already

incorporated it into the set of variables to be analyzed in daily clinical practice.

We have carefully reviewed the articles referenced in your letter. We have not found a clear justification in both studies as they have not used a learning curve analysis methodology and both have some biases in this respect.

We believe, and this has been our experience, that the robotic platform shortens the learning curves. As Müller et al. reported, the analysis of learning curves in literature is arbitrary in some cases.³

In the last years, new literature has become available, and robotic pancreatic surgery had a significant expansion. Recently, the internationally validated European guidelines on minimally Invasive Pancreatic Surgery proclaimed that the learning curves for robotic distal pancreatectomy are slightly shorter compared with a laparoscopic approach. Apart from,