



clínica e investigación en ginecología y obstetricia

www.elsevier.es/gine



ORIGINAL ARTICLE

Impact of the COVID-19 pandemic on maternal anxiety during pregnancy: A prevalence study



M. Bermúdez-González*, E. Álvarez-Silvares, J.K. Santa-María-Ortiz,
L. Castro-Vilar, M. Vázquez-Rodríguez

Obstetrics and Gynecology Service, Complejo Hospitalario Universitario de Ourense, Galicia, Spain

Received 21 December 2021; accepted 23 May 2022

Available online 8 June 2022

KEYWORDS

COVID-19;
Low obstetric risk;
Maternal anxiety;
Pandemic;
Pregnancy

Abstract

Aim: To assess the impact of the coronavirus SARS-CoV-2 pandemic on the level of anxiety in low-risk pregnant women.

Material and method: Epidemiological, descriptive, prevalence study. A total of 74 patients who underwent low risk antenatal controls during the state of alarm because of COVID-19, were included. They filled in the Hamilton Anxiety Rating Scale and a specific document about the pandemic. Clinical histories and different variables of clinical interest were reviewed and compiled, respectively.

Results: Mean age was 34.05 years with average amenorrhoea of 28.17 weeks. A total of 77% of the sample presented symptoms and signs compatible with anxiety. Of these, 44.6% and 32.4% presented minor and major anxiety, respectively. Concern over the time of the birth and postpartum and fear of being at greater risk because of possible infection was present in 95.9% and 94.6% of the sample, respectively. A total of 93.2% of the sample was afraid of intrauterine virus transmission; 94.5% admitted fear over the neonatal consequences of infection.

Conclusions: The pregnant women assessed had three times more anxiety during the COVID-19 pandemic. This incidence is independent of most study variables.

© 2022 Published by Elsevier España, S.L.U.

PALABRAS CLAVE

COVID-19;
Bajo riesgo
obstétrico;
Ansiedad materna;
Pandemia;
Embarazo

Impacto de la pandemia por COVID-19 en la ansiedad materna durante el embarazo: Un estudio de prevalencia

Resumen

Objetivo: Evaluar el impacto de la pandemia del coronavirus SARS-CoV-2 en el nivel de ansiedad en mujeres embarazadas de bajo riesgo.

* Corresponding author.

E-mail address: monica.bermudez.gonzalez@sergas.es (M. Bermúdez-González).

Material y método: Estudio epidemiológico, descriptivo, de prevalencia. Se incluyeron un total de 74 pacientes que se sometieron a controles prenatales de bajo riesgo durante el estado de alarma por COVID-19. Completaron la escala de calificación de ansiedad de Hamilton y un documento específico sobre la pandemia. Se revisaron y recopilamos historias clínicas y diferentes variables de interés clínico, respectivamente.

Resultados: La edad promedio fue de 34,05 años con amenorrea promedio de 28,17 semanas. El 77% de la muestra presentó síntomas y signos compatibles con la ansiedad. De estos, el 44,6 y el 32,4% presentaron ansiedad menor y mayor, respectivamente. La preocupación por el momento del parto y el puerperio y el temor de presentar mayor riesgo por una posible infección estuvieron presentes en el 95,9 y 94,6% de la muestra, respectivamente. El 93,2% de la muestra temía una posible transmisión del virus intrauterino; el 94,5% admitió tener miedo a las consecuencias neonatales tras una posible infección.

Conclusiones: Las embarazadas evaluadas tenían tres veces más ansiedad durante la pandemia de COVID-19. Esta incidencia es independiente de la mayoría de las variables de estudio.

© 2022 Publicado por Elsevier España, S.L.U.

Introduction

The current pandemic caused by coronavirus disease (COVID-19) – severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) – has spread quickly across the globe and has led to a public health emergency. From December 31, 2019 to August, 2021, a total of 225,166,539 COVID-19 cases, including 4,636,120 deaths – according to data provided by the European Center for Disease Prevention and Control (ECDC),¹ have been reported.

Pregnant women undergo immunologic and physiologic changes that could make them more susceptible to viral respiratory infections, including COVID-19. Therefore, they are a high-risk population during outbreaks of infectious disease. The impact of COVID-19 infection during pregnancy has been assessed and – although data are limited and confined to case series mainly during the third trimester – infection during pregnancy probably has a clinical presentation and severity that is similar to non-pregnant women. Moreover, some authors do not detect an association with poor maternal or perinatal results,^{2,3} although there is little current evidence, and this is under constant review. To date, there are only a few recorded cases of intrauterine infection caused by vertical transmission in women developing pneumonia because of COVID-19.^{4,5}

Despite available evidence revealing reassuring maternal and fetal outcomes, the current state of the pandemic could lead to the onset of pathologic emotional states and a consequent reduction in the quality of life of pregnant women. The prevalence of maternal anxiety during a non-COVID-19 pregnancy is estimated to be 15%–23% of women according to the series consulted.^{6–9} Antenatal stress and anxiety have been related to the onset of both short and long term adverse maternal, fetal and neonatal events.^{10,11} Anxiety during pregnancy has, therefore, been strongly correlated to anxiety and depression during the postnatal period.^{12,13} Against this backdrop, low birth weight,^{14,15} premature birth¹⁶ and abnormal cognitive and behavioral neurologic

development^{17,18} are among the negative outcomes reported at birth and during childhood.

An increase in premature birth rates, low birth weight and higher child mortality within the scope of other traumatic events, such as the terrorist attacks in New York and Madrid, has also been recorded. The negative effect on reproductive outcomes reported is related to probable post-traumatic stress undergone by the pregnant women exposed to these events.^{19–21}

Therefore, having set out the significant consequences of antenatal anxiety on maternal and child health, the aim of our study will be to determine the impact of the current COVID-19 pandemic on maternal anxiety of the low risk pregnant woman who has not been diagnosed with COVID-19.

Materials and methods

An epidemiological, descriptive, prevalence study was performed. A total of 74 patients was included. The inclusion criteria were those pregnant women who underwent low obstetric risk, antenatal, gynecologic controls in the department of Obstetrics and Gynecology of Ourense Teaching Hospital Complex during the healthcare state of alarm because of coronavirus SARS-CoV-2 between April and May 2020. Pregnant women at high obstetric risk (hypertensive states of pregnancy, diabetes mellitus, BMI > 40, autoimmune disease, twin gestation, epilepsy, clinical hypothyroidism and hyperthyroidism, and poor obstetric history), pregnant women in the first trimester, and those diagnosed with COVID-19 were excluded. Therefore, the sampling method was non-probabilistic for convenience. The selection process is shown in Fig. 1.

For study purposes patients filled in two short questionnaires: first, the Hamilton Anxiety Rating Scale (HARS), a document comprising 14 items and the gold standard for the study of anxiety, validated in Spanish^{22–24} and, second, a questionnaire comprised of 9 specific questions on the current state of the COVID-19 pandemic, with the




Figure 1 Selection process.

Table 1 Study variables.

<i>Pregnant woman variables</i>	
Age	BMI
Number of prior pregnancies	Preconception
Number of prior deliveries	Pregestational hypothyroidism
Number of prior miscarriages	
<i>Gestational variables</i>	
Gain in gestational weight	Fetal weight percentile on ultrasound 3rd trimester
Result of chromosomal disorder screening	Trimester of pregnancy

aim of relating the scores obtained on the previous form with the current pandemic. To assess whether or not anxiety is present the proposal made by Bech was used; by way of guidance, this sets out the following ranges: 0–5 points (no anxiety), 6–14 (less anxiety), 15 or more (more anxiety).²⁵ With the aim of obtaining the study parameters the electronic clinical histories of patients included were also reviewed. These variables are shown in Table 1.

The pregnant women included in the study gave their written informed consent. The study was approved by the institutional ethics committee.

A descriptive analysis where qualitative variables were expressed as frequency and percentage, was initially performed. Continuous variables were expressed as mean \pm standard deviation, median and minimum–maximum.

Non-parametric tests were performed to determine the potential association between the study variables (Chi-squared, Kruskal–Wallis, Mann–Whitney U tests). Correlations were studied to detect a relationship or interaction between the different variables.

Differences with $P < 0.05$ were considered statistically significant for all analyses. Analyses were performed using the software IBM SPSS Statistics for Windows, Version 22.0. Armonk, NY, USA: IBM Corp.

Results

Descriptive analysis of the sample

A total of 74 pregnant women were included in the study. Mean age at the time of the intervention was 34.05 years (minimum age: 19; maximum age: 46). Mean gestational age was 28.17 weeks (minimum gestational age: 15.28; maximum gestational age: 41.42). Most subjects did not present previous deliveries, miscarriages or c-sections (63.5%, 68.9% and 91.9%, respectively). A total of 52.7% of patients were in the third trimester of pregnancy and 47.3% remained in the second trimester. Subjects in the first trimester of pregnancy were not included in the study. Prior history of anxiety was only presented by 4.1% of the sample. Furthermore, only 13.5% of pregnant women were diagnosed with a thyroid disorder in form the subclinical hypothyroidism. A total of 23%, 66.2% and 10.8% had excessive weight gain, normal weight or remained under weight, respectively. Most pregnant women (83.8%) had a low risk result of chromosomal disorder screening compared to 6.8% and 9.5% classified as intermediate and high-risk, respectively. The general characteristics are shown in Table 2.

Table 2 General characteristics.

Study variables	Mean	Standard deviation	Minimum	Maximum
Age	34.05	5.76	19	40
Gestational age	28.17	7.55	15.28	41.42
	Frequency		Valid percentage (%)	
	Yes	No	Yes	No
Deliveries	27	47	36.5	63.5
Miscarriages	23	51	31.1	68.9
C-sections	6	68	8.1	91.9
Prior anxiety	3	71	4.1	95.9
Thyroid abnormality	10	64	13.5	86.5
Weight gain (kg)	Frequency		Valid percentage (%)	
No gain	8		10.8	
Normal	49		66.2	
Excessive	17		23.0	
Chromosomal disorder screening	Frequency		Valid percentage (%)	
Low risk	62		83.7	
Intermediate risk	5		6.8	
High-risk	7		9.5	
HARS	Frequency		Valid percentage (%)	
No anxiety	17		23.0	
Less anxiety	33		44.6	
More anxiety	24		32.4	
Total	74		100.0	

Maternal anxiety and the COVID-19 pandemic

According to results obtained approximately 8 in 10 patients (77%) reported presenting symptoms and signs compatible with anxiety. Of these, 44.6% and 32.4% presented minor and major anxiety, respectively.

Most pregnant women (95.9%) said they were concerned about the time of the birth and postpartum after starting the confinement because of COVID-19. Moreover, a total of 94.6% confessed worry in regard to the possibility of being more at risk because of pregnancy in the event of a possible infection. A total of 93.2% of the sample revealed unease in the event of possible intrauterine transmission of the virus to the baby. And 94.5% confessed being scared about the consequences the baby could present at birth after a possible maternal infection. Moreover, 9 out of every 10 pregnant women (91.4%) were worried about contagiousness during the birth or subsequent stay on the obstetrics ward. To a lesser extent, half of the patients (50.7%) were concerned about the possibility of less healthcare personnel being present at the time of the birth. Conversely, most of the sample (62.9%) had no doubts over starting to breastfeed. Only 37.1% were wary of starting to breastfeed during the current pandemic. These results are shown in Table 3.

No statistically significant differences were observed between most of the different study variables and their relationship with maternal anxiety indices. However, a

statistically significant association was observed between those patients who had not given birth previously and more concern over the time of the birth and postpartum after the healthcare state of alarm began (Exact Fisher test: $P < 0.05$). It was also observed that those patients with diagnostic anxiety scores thought repeatedly about the above issues, which affected their quality of life (Chi-squared test = 7.960, $P < 0.05$).

Discussion

The data obtained in the present study seem to demonstrate an increase in maternal anxiety in the context of the SARS-COV-2 pandemic when the results obtained are compared with those previously published by other authors. Recent publications conclude similar results and also report increased levels of antenatal anxiety.^{26–30}

The prevalence of antenatal anxiety in our sample tripled that reported in scientific literature (15%–21%^{6–9}). This figure is notable among those disseminated by different works. The study by Mappa et al.²⁶ reported half the number of women who attained abnormal anxiety levels; specifically 38.2% of the sample as opposed to 77% obtained in our study. Along the same lines, the work by Wu et al.²⁷ reported 29.6% of pregnant women with symptoms and anxiety and depression. Such discrepancies might be accounted for by the use of different instruments to measure maternal anxiety or be

Table 3 COVID-19 questionnaire.

Questions	Frequency			Valid percentage	
	Yes	No	NA	Yes	No
Given that you are pregnant are you worried about being more at risk in the event of a possible Covid-19 infection?	70	4	0	94.6	5.4
Are you worried about intrauterine Covid-19 transmission to your baby?	68	5	1	93.2	6.8
Have you thought about the consequences your baby could present at birth?	69	4	1	94.5	5.5
Are you more worried about the time of the birth and postpartum after starting the healthcare state of alarm because of Covid-19?	70	3	1	95.9	4.1
Are you scared about your child's Covid-19 contagion during the birth or subsequent stay on the obstetric ward?	64	6	4	91.4	8.6
Are you scared about less healthcare personnel being available at the time of the birth, in case of need?	35	34	5	50.7	49.3
Do you have doubts over the suitability of starting to breastfeed given the Covid-19 pandemic?	26	44	4	37.1	62.9
Do you repeatedly think about the previous questions?	39	34	1	53.4	46.6
Do you believe these thoughts affect your quality of life?	31	41	2	43.1	56.9

NA: no answer.

due to the confinement in Spain – among the strictest in the world – which might increase the perception of level of severity perceived by the population.

This increase can be observed in low obstetric risk pregnant women regardless of specific features studied (age, BMI, existence of preconception anxiety or thyroid pathology). Nor have significant differences been observed in regard to factors such as the number of prior gestations, births and miscarriages, trimester of pregnancy, increase in gestational weight or result of chromosomal disorder screening. Similar results were obtained by Taubman et al.²⁸ when they reported an increase in gestational anxiety regardless of the various sociodemographic features presented. Mappa et al.²⁶ did not reveal differences either according to the variables studied except for the high educational level associated with higher percentages of perinatal anxiety. However, the work by Wu et al.²⁷ reported a higher risk of presenting symptoms of anxiety and depression in those patients giving birth for the first time with a low gestational BMI, aged under 35, employed full time and with an average income.

The most important concerns reported by pregnant women in our study were in more than 9 out of 10 cases: fear of being at risk patients in the event of a possible infection, fear of intrauterine viral transmission, neonatal consequences in the event of a possible maternal infection and infection during the birth or subsequent stay on the obstetric ward. To a lesser extent, up to half the women admitted they were scared about the possibility of less healthcare personnel available at the time of the birth. Other studies have reported similar maternal concerns although it is notable that these were present in lower percentages. Therefore, Taubman et al.²⁸ reported higher anxiety levels in regard to maternal exposure in public

places and transport (87.5% and 70%, respectively); to a lower extent fear of infection of other family members or fetal health, followed by attending gynecologic consultations and maternal infection at the time of the birth. Lower figures were revealed by Mappa et al.²⁶ given that 65% of the sample were afraid of a possible restriction in fetal growth; and to a lesser degree, premature birth and the possibility of fetal structural abnormalities.

The most important limitations of our study are the small sample size and subjects belonging to just one center which does not enable generalization of the data. Furthermore, as this is a prevalence study it is not possible to set out a causal relationship with the onset of possible future maternal and fetal consequences. However, to the best of our knowledge this work is the first to be performed in Spain and to date one of the few to be published in other settings.

To conclude, our work reveals a marked increase in maternal anxiety after onset of the COVID-19 pandemic. This increase was observed regardless of the majority of variables studied; both sociodemographic factors and specific features of the pregnancy.

Ethics approval

This study was performed in line with the principles of the Declaration of Helsinki. Approval was granted by the Galician drug research ethics committee (CEIm-G) (28-05-2020).

Authors' contributions

All authors contributed to the conception and design of the study and performed material preparation, data collection, and analysis.

Ethical responsibilities

Protection of people and animals. The authors declare that the procedures followed were in accordance with the ethical standards of the responsible human experimentation committee and in accordance with the World Medical Association and the Declaration of Helsinki.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors have obtained the informed consent of the patients and/or subjects referred to in the article. This document is in the possession of the corresponding author.

Funding

No grants or other types of support were used to draw up this article.

Conflicts of interest

The authors declare no competing interests.

References

1. ECDC: Situation update worldwide, as of week 36, updated 16 September 2021 [Internet]. Sept 22, 2021. Available at: <https://www.ecdc.europa.eu/en/geographical-distribution-2019-ncov-cases> [Accessed 22.09.21].
2. Yan J, Guo J, Fan C, Juan J, Yu X, Li J, et al. Coronavirus disease 2019 (COVID-19) in pregnant women: a report based on 116 cases. *Am J Obstet Gynecol*. 2020;223:1–48, <http://dx.doi.org/10.1016/j.ajog.2020.04.014>.
3. Elshafeey F, Magdi R, Hindi N, Elshebiny M, Farrag N, Mahdy S, et al. A systematic scoping review of COVID-19 during pregnancy and childbirth. *Int J Gynaecol Obstet*. 2020;150:1–16, <http://dx.doi.org/10.1002/ijgo.13182>.
4. Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet*. 2020;395:809–15, [http://dx.doi.org/10.1016/S0140-6736\(20\)30360-3](http://dx.doi.org/10.1016/S0140-6736(20)30360-3).
5. Schwartz DA. An analysis of 38 pregnant women with COVID-19, their newborn infants, and maternal-fetal transmission of SARS-CoV-2: maternal coronavirus infections and pregnancy outcomes. *Arch Pathol Lab Med*. 2020;144:1–25, <http://dx.doi.org/10.5858/arpa.2020-0901-SA>.
6. Dennis CL, Falah-Hassani K, Shiri R. Prevalence of antenatal and postnatal anxiety: systematic review and meta-analysis. *Br J Psychiatry*. 2017;210:315–23, <http://dx.doi.org/10.1192/bjp.bp.116.187179>.
7. Rubertsson C, Hellström J, Cross M, Sydsjö G. Anxiety in early pregnancy: prevalence and contributing factors. *Arch Womens Ment Health*. 2014;17:221–8, <http://dx.doi.org/10.1007/s00737-013-0409-0>.
8. Goodman JH, Chenausky KL, Freeman MP. Anxiety disorders during pregnancy: a systematic review. *J Clin Psychiatry*. 2014;75:e1153–84, <http://dx.doi.org/10.4088/JCP.14r09035>.
9. National Institute for Health and Care Excellence. Antenatal and Postnatal Mental Health: Clinical Management and Service Guidance (CG192). NICE; 2014.
10. Reissland N, Froggatt S, Reames E, Girkin J. Effects of maternal anxiety and depression on fetal neurodevelopment. *J Affect Disord*. 2018;241:469–74, <http://dx.doi.org/10.1016/j.jad.2018.08.047>.
11. Nowak AL, Anderson CM, Mackos AR, Neiman E, Gillespie SL. Stress during pregnancy and epigenetic modifications to offspring DNA: a systematic review of associations and implications for preterm birth. *J Perinat Neonatal Nurs*. 2020;34:134–45, <http://dx.doi.org/10.1097/JPN.0000000000000471>.
12. Verreault N, DaCosta D, Marchand A, Ireland K, Dritsa M, Khalife S. Rates and risk factors associated with depressive symptoms during pregnancy and with postpartum onset. *J Psychosom Obstet Gynaecol*. 2014;35:84–91, <http://dx.doi.org/10.3109/0167482X.2014.947953>.
13. Milgrom J, Gemmil A, Bilszta JL, Hayes B, Barnett B, Brooks J, et al. Antenatal risk factors for postnatal depression: a large prospective study. *J Affect Disord*. 2008;108:147–57, <http://dx.doi.org/10.1016/j.jad.2007.10.014>.
14. Field T, Diego M, Hernandez-Reif M, Figueroa B, Deeds O, Ascencio A, et al. Comorbid depression and anxiety effects on pregnancy and neonatal outcome. *Infant Behav Dev*. 2010;33:23–9, <http://dx.doi.org/10.1016/j.infbeh.2009.10.004>.
15. Diego MA, Jones NA, Field T, Hernandez-Reif M, Schanberg S, Kuhn C, et al. Maternal psychological distress, prenatal cortisol, and fetal weight. *Psychosom Med*. 2006;68:747–53, <http://dx.doi.org/10.1097/01.psy.0000238212.21598.7b>.
16. Ding XX, Wu YL, Xu SJ, Zhu RP, Jia XM, Zhang SF, et al. Maternal anxiety during pregnancy and adverse birth outcomes: a systematic review and meta-analysis of prospective cohort studies. *J Affect Disord*. 2014;159:103–10, <http://dx.doi.org/10.1016/j.jad.2014.02.027>.
17. Stein A, Pearson RM, Goodman SH, Rapa E, Rahman A, McCallum M, et al. Effects of perinatal mental disorders on the fetus and child. *Lancet*. 2014;384:1800–19, [http://dx.doi.org/10.1016/S0140-6736\(14\)61277-0](http://dx.doi.org/10.1016/S0140-6736(14)61277-0).
18. Talge NM, Neal C, Glover V. Antenatal maternal stress and long-term effects on child neurodevelopment: how and why? *J Child Psychol Psychiatry*. 2007;48:245–61, <http://dx.doi.org/10.1111/j.1469-7610.2006.01714.x>.
19. Sherrieb K, Norris FH. Public health consequences of terrorism on maternal-child health in New York City and Madrid. *J Urban Health*. 2013;90:369–87, <http://dx.doi.org/10.1007/s11524-012-9769-4>.
20. Maslow CB, Caramanica K, Li J, Stellman SD, Brackbill RM. Reproductive outcomes following maternal exposure to the events of September 11, 2001, at the World Trade Center, in New York City. *Am J Public Health*. 2016;106:1796–803, <http://dx.doi.org/10.2105/AJPH.2016.303303>.
21. Eskenazi B, Marcas AR, Catalano R, Bruckner T, Toniolo PG. Low birthweight in New York city and upstate New York following the events of September 11th. *Hum Reprod*. 2007;22:3013–20, <http://dx.doi.org/10.1093/humrep/dem301>.
22. Lobo A, Chamorro L, Luque A, Dal-Ré R, Badiá X, Baró E. Validación de las versiones en español de la Montgomery-Asberg Depression Rating Scale y la Hamilton Anxiety Rating Scale para la evaluación de la depresión y de la ansiedad. *Med Clin (Barc)*. 2002;118:493–9.
23. Shear MK, Vander Bilt JJ, Rucci PP, et al. Reliability and validity of a structured interview guide for the Hamilton Anxiety Rating Scale (SIGH-A). *Depress Anxiety*. 2001;13:166–78.
24. Van Ameringen M. Comorbid anxiety and depression in adults: epidemiology, clinical manifestations, and diagnosis. <https://www.uptodate.com> [accessed 5.03.20].

25. Bech P. *Rating scales for psychopathology, health status and quality of life*. New York: Springer-Verlag Berlin Heidelberg; 1993.
26. Mappa I, Distefano FA, Rizzo G. Effects of coronavirus 19 pandemic on maternal anxiety during pregnancy: a prospective observational study. *J Perinat Med*. 2020;48:545–50, <http://dx.doi.org/10.1515/jpm-2020-0182>.
27. Wu Y, Zhang C, Liu H, Duan C, Li C, Fan J, et al. Perinatal depressive and anxiety symptoms of pregnant women during the coronavirus disease 2019 outbreak in China. *Am J Obstet Gynecol*. 2020;223:240e1–9, <http://dx.doi.org/10.1016/j.ajog.2020.05.009>.
28. Taubman O, Chasson M, Sharkia SA, Weiss E. Distress and anxiety associated with COVID-19 among Jewish and Arab pregnant women in Israel. *J Reprod Infant Psychol*. 2020;38:340–8, <http://dx.doi.org/10.1080/02646838.2020.1786037>.
29. Berthelot N, Lemieux R, Garon-Bissonnette J, Drouin-Maziade C, Martel E, Maziade M. Uptrend in distress and psychiatric symptomatology in pregnant women during the coronavirus disease 2019 pandemic. *Acta Obstet Gynecol Scand*. 2020;99:848–55, <http://dx.doi.org/10.1111/aogs.13925>.
30. Liu X, Chen M, Wang Y, Sun L, Zhang J, Shi Y, et al. Prenatal anxiety and obstetric decisions among pregnant women in Wuhan and Chongqing during the COVID-19 outbreak: a cross-sectional study. *BJOG*. 2020;127:1229–40, <http://dx.doi.org/10.1111/1471-0528.16381>.