

Endocrinología, Diabetes y Nutrición



209 - IDENTIFICATION OF OXYGEN-18 ISOTOPE OF BREATH CARBON DIOXIDE AS A NON-INVASIVE MARKER TO DISTINGUISH TYPE 1 AND TYPE 2 DIABETES

C. Ghosh and M. Pradhan

S. N. Bose National Centre For Basic Sciences. India.

Resumen

Introduction: There is a pressing need to develop a new and an effective strategy for early detection of T1D and to precisely distinguish T1D from type 2 diabetes (T2D). The aim of the present study was to find out the potential link between the erythrocytes carbonic anhydrase (CA) activity and ^{18}O -isotopic exchange of breath CO₂ in T1D and T2D.

Methods: Fasting and post-dose breath and blood samples were collected simultaneously after ingestion of 75-gm normal glucose dissolved in 150-mL water. Blood samples were analysed to measure the CA activity. The breath samples were utilised to measure the carbon dioxide isotopes (\$^{12}C^{16}O^{16}O\$, \$^{13}C^{16}O^{16}O\$ and \$^{12}C\$ and \$^{16}O^{18}O\$) by a laser based high-precision carbon dioxide isotope analyzer.

Results: The CA activities are markedly altered during metabolism of T1D and T2D and this facilitates to oxygen-18 (18 O) isotopic fractionations of breath CO $_2$. In our observations, T1D exhibited considerable depletions of 18 O-isotopes of CO $_2$ whereas T2D manifested isotopic enrichments of 18 O in breath CO $_2$, thus unveiling a missing link of breath 18 O-isotopic fractionations in T1D and T2D. The optimal diagnostic cut-off points were determined to be $?_{DOB}^{18}$ O% = 2.1% and ?CA = 3.15 U/min/mL for screening T1D and T2D individuals.

Conclusions: Our findings suggest the changes in erythrocytes CA activities may be the initial step of altered metabolism of T1D and T2D, and breath ¹⁸O-isotope regulated by the CA activity is a potential diagnostic biomarker that can selectively and precisely distinguish T1D from T2D and thus may open a potential unifying strategy for treating these diseases.