



## 1 - METABOLIC EXPOSURES TO PERIODIC CYCLES OF LIMITED FOOD INTAKE BOOSTS HEALTHSPAN AND DEFINES A MOLECULAR MEMORY OF FASTING

A. Díaz-Ruiz<sup>1,2</sup>, J.L. López-Cánovas<sup>1</sup>, M.D. Gahete<sup>3,4,5</sup>, T. Rhinesmith<sup>2</sup>, L.C.D. Pomatto<sup>2</sup>, N.L. Price<sup>2</sup>, M. Bernier<sup>2</sup> and R. de Cabo<sup>1,2</sup>

<sup>1</sup>Nutritional Interventions Group. Precision Nutrition and Aging. Institute IMDEA Food. Madrid. <sup>2</sup>Experimental Gerontology Section. Translational Gerontology Branch. National Institute on Aging. National Institutes of Health. Baltimore, MD. USA. <sup>3</sup>OncOesity and Metabolism. Instituto Maimónides de Investigación Biomédica de Córdoba (IMIBIC). Córdoba. <sup>4</sup>Departamento de Biología Celular, Fisiología e Inmunología. Universidad de Córdoba. <sup>5</sup>Hospital Universitario Reina Sofía. Córdoba. <sup>6</sup>CIBER Fisiopatología de la Obesidad y Nutrición. CIBERObn. Córdoba.

### Resumen

**Introduction:** Caloric restriction and other forms of fasting, such as cycles of limited food intake, promote healthy aging and prevent age-associated metabolic diseases. Underneath these strategies, identifying biological and nutritional adaptations remains a scientific challenge with social and economic implications.

**Methods:** Middle-aged mice were subjected for 5 months to 4:10 feeding cycles, consisting of 4 days of very low-calorie intake (VLCI) followed by 10 days of *ad libitum* regime. The impact of 4:10 cycles was determined with respect to whole-body physiology, metabolic and bioenergetic status, as well as the dynamic response to stimuli by metabolite profiling. Hepatic plasticity of the spliceosome was also evaluated.

**Results:** Short-term benefits of VLCI included body weight, fat and lean mass loss, and up-modulation of 4 hepatic pathways (purines, lipids/ketone bodies, ascorbate, and redox). Long-term benefits of repeated 4:10 cycles included body weight and fat mass loss, enhanced physical performance, and a long-lasting metabolic memory of fasting. Genes of the hepatic splicing machinery classified into 7 independent clusters of unique expression patterns. Interestingly, metabolic exposures to VLCI primed some clusters to have a memory of fasting. A pairwise comparisons between *ad libitum* vs. fasted state further identified a molecular signature of 64 spliceosomal genes strongly altered, the vast majority downregulated [80%]. Of note, some of them correlated with well-known biomarkers of the organismal bioenergetic and nutritional status.

**Conclusions:** This study supports the concept that organismal, metabolic, and molecular adaptations to periodic cycles of VLCI may be partially driven by a long-lasting nutritional memory, which provide information for the optimization of the use of cycles of limited food intake to promote health.

**Funding:** 2018-T1/BMD-11966 | PID2019-106893RA-I00 | IRP- NIA.