

Tubular ectasia of the rete testis and another benign scrotal lesions

Ectasia tubular de la rete testis y otras lesiones escrotales benignas

Figures 1 to 3 – Patient with a palpable 1-cm nodule in the left hemiscrotum. Ultrasound revealed an extratesticular anechoic lesion compatible with a tunica albuginea cyst (fig. 1). Same patient: cystic and tubular anechoic lesions in the mediastinum testis, in relation to tubular ectasia of the rete testis, which should not be mistaken for a tumor (fig. 2). Spermatocele of the epididymal head, often associated to ectasia of the rete testis (fig. 3).

A.X. Casal Rodríguez*, P. Rodríguez Fernández,
C.M. Rodríguez Paz, A. Tilve Gómez, and M. Otero García

Department of Urology, Hospital Xeral-Cies (CHUVI), Vigo, Spain

*Corresponding author.

E-mail: anton.casal.rodiguez@sergas.es
(A.X. Casal Rodríguez).

Genital reconstruction in a patient with Fournier gangrene

Reconstrucción genital en paciente con gangrena de Fournier



Figure 1 – A 51-year-old patient with a history of chronic alcoholism presenting Fournier gangrene requiring surgical debridement and cleaning of the genital area due to extensive necrosis of the subcutaneous lax tissue and skin.



Figure 2 – Reconstruction of the genital zone using flaps from the remaining scrotal skin. No placement of grafts from other body locations proved necessary.

M.A. Arrabal-Polo^{a,*}, F. Almazán-Fernández^b,
S. Merino-Salas^a, S. Arias-Santiago^b, R. Armijo-Lozano^b,
M. Arrabal-Martín^a, and A. Zuluaga-Gómez^a

^aDepartment of Urology, Hospital Universitario San Cecilio,
Granada, Spain

^bDepartment of Dermatology, Hospital Universitario San Cecilio,
Granada, Spain

*Corresponding author.

E-mail: arrabalp@ono.com (M.A. Arrabal-Polo).