



Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



Review article

Epidemiology, prevention and control of pertussis in Spain: New vaccination strategies for lifelong protection



Juan José González-López^{a,b,*}, Javier Álvarez Aldeán^c, Francisco José Álvarez García^{d,e}, Magda Campins^f, María Garcés-Sánchez^g, Ruth Gil-Prieto^h, Ana M^a Grande-Tejadaⁱ

^a Department of Clinical Microbiology, Hospital Vall d'Hebron, Barcelona, Spain

^b Department of Microbiology and Genetics, Universitat Autònoma de Barcelona, Spain

^c Pediatric Service, Hospital Costa del Sol, Málaga, Spain

^d Pediatrics, Centro de Salud de Llanera, Asturias, Spain

^e Department of Medicine, Universidad de Oviedo, Asturias, Spain

^f Department of Preventive Medicine and Epidemiology, Hospital Vall d'Hebron, Barcelona, Spain

^g Pediatrics, Centro de Salud de Nazaret, Valencia, Spain

^h Department of Medicine and Public Health, Universidad Rey Juan Carlos, Madrid, Spain

ⁱ Pediatrics, Hospital Materno Infantil, Badajoz, Spain

ARTICLE INFO

Article history:

Received 27 January 2021

Accepted 22 April 2021

Available online 18 June 2021

Keywords:

Pertussis

Vaccination

Epidemiology

Prevention

ABSTRACT

Pertussis is a highly contagious, vaccine-preventable respiratory tract infection, with high morbidity and mortality and a particularly severe effect on newborns and infants under 2 months. The first pertussis vaccines were introduced in the 1940s. Since 1980, however, the incidence of cases has risen despite the extensive vaccination programmes and antibiotic adjuvant treatments available. Transition from the use of whole-cell vaccines to acellular vaccines and the antigenic modifications of *Bordetella pertussis* have contributed, among other factors, to a reduction in vaccine-acquired immunity and reemergence of the disease. Today, there are still unmet needs not covered by conventional prevention measures and existing antibiotic treatments. This review aims to update the available data, and to discuss which vaccine strategies might contribute to better disease control and prevention.

© 2021 Sociedad Española de

Enfermedades Infecciosas y Microbiología Clínica. Published by Elsevier España, S.L.U. All rights reserved.

Epidemiología, prevención y control de la tosferina en España: nuevas estrategias de vacunación para la protección a lo largo de la vida

RESUMEN

La tosferina es una infección respiratoria inmunoprevenible altamente contagiosa, con alta morbilidad y mortalidad, y que afecta con especial gravedad a recién nacidos y lactantes menores de 2 meses. Las primeras vacunas comenzaron a emplearse en la década de los 40. Sin embargo, desde 1980 la incidencia de casos ha aumentado a pesar de los amplios programas de vacunación y tratamientos antibióticos adyuvantes disponibles. El cambio del uso de vacunas celulares a vacunas acelulares, y las modificaciones antigénicas de *B. pertussis* han podido contribuir entre otros factores a la disminución de la inmunidad adquirida tras la vacunación y a la reemergencia de la enfermedad. En la actualidad, todavía existen necesidades no cubiertas por las medidas convencionales de prevención y los tratamientos antibióticos existentes. Esta revisión pretende actualizar los datos disponibles y plantear qué estrategias vacunales pueden contribuir a un mejor control y prevención global de la enfermedad.

© 2021 Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Tosferina

Vacunación

Epidemiología

Prevención

* Corresponding author.

E-mail address: jjgonzal@vhebron.net (J.J. González-López).

Rationale

The main aetiological agent of pertussis (also known as whooping cough) is *Bordetella pertussis*, a gram-negative coccobacillus with a strictly human reservoir and high airborne transmissibility. Pertussis is an acute respiratory infection characterized by a violent cough that can last for weeks or months and is seen mostly in the preschool and school-age population. Although adolescents and adults are the main reservoir of the microorganism and play a crucial role in its transmission, it is infants under 4 months who have the highest incidence and most severe forms of the disease, which often requires advanced life support. In the last 20 years, a reemergence of the infection has been observed, making it one of the most prevalent vaccine-preventable diseases. The appearance of *B. pertussis* strains that escape the immunity conferred by the vaccine and the gradual reduction of the protection acquired after vaccination are among the possible causes of this reemergence. These situations justify the need to adapt the current preventive, diagnostic and therapeutic strategies with the primary aim of protecting the most vulnerable population (infants <4 months) and reducing the rising trend in the incidence of pertussis in the general population. This review updates the epidemiological situation of pertussis, the microbiological aspects related to the adaptation of *B. pertussis* and the diagnosis of the disease, and the preventive actions related to existing vaccine strategies that may help to improve its control and prevention.

Epidemiological situation of pertussis

The European Union (EU) network for epidemiological surveillance and control of vaccine-preventable infections estimates that 35 627 cases of pertussis were reported by the participating countries in 2018, with a notification rate of 8.2 cases/100 000 population, similar to that of the previous 4 years but the lowest observed in the same time period.¹ Five countries (Germany, the Netherlands, Norway, Spain and the United Kingdom) accounted for 72% of all notified cases, and 62% of cases were individuals aged over 15 years. Infants under 1 year, who had not completed the primary pertussis vaccination series because of their young age, were the most affected age group with the highest incidence rate (44.4 cases/100 000 population), followed by children aged 10–14 years.

The epidemiological evolution of the infection in adolescents and adults is of particular concern because of their potential to transmit the bacteria to infants. In adults, clinical suspicion of pertussis is low, which may explain the significant underreporting of cases.² The review of the definition of pertussis cases by the EU in 2018 probably helped to identify atypical presentations of the disease in adults, adolescents, and vaccinated individuals, and to clarify aspects of laboratory confirmation.^{3,4}

Most EU states report pertussis cases according to the Community definition, based on extensive passive surveillance systems with national coverage. Even so, the form of notification is heterogeneous. Belgium uses a laboratory-based, voluntary sentinel surveillance system covering the entire population, while France has a hospital-based sentinel surveillance system, which includes only infants under 6 months. There are two surveillance systems in Denmark: one laboratory-based (which includes all age groups) and another in which doctors must report pertussis cases in children under 2 years.

In Spain, pertussis has been a notifiable disease since 1982 and individually notifiable since 1997. Passive epidemiological surveillance systems have significant problems of underdiagnosis, not only in adults, but also in children.² Underreporting of symptomatic cases is considerable in children ≥ 1 year, although the highest percentage of underreporting (50%) has been

identified in infants under 1 year. In the period 1997–2010, the overall reported incidence of hospitalization for pertussis was 1.3 cases/100 000 population in children and adults. Minimum under-reporting ranged from 3.8% to 22.8%, according to the study year.⁵ These data support the need for rapid laboratory diagnostic tests and greater surveillance of family contacts of any age that help to better quantify existing cases. Intensified surveillance may also be useful for a better understanding of pertussis transmission patterns in the family and community. Surveillance and monitoring of pertussis incidence and outbreaks can be improved by standardizing and strengthening regional epidemiological surveillance systems with the use of complementary active systems that allow actual incidence data to be determined and by introducing innovative approaches (such as Big Data analysis, based on real-time tracking of Internet search activities).^{6,7}

Pertussis incidence in Spain

The incidence of pertussis has increased significantly in Spain since 2011. The highest rates are recorded in infants under 1 year, followed by children aged 5–9 years, as observed in Catalonia, Galicia, Madrid and Valencia and in outbreaks in Andalusia, Valencia, and Catalonia.^{8–14}

Despite high vaccination coverage, pertussis maintains its epidemic presentation. Between 1998 and 2018, five epidemic waves of pertussis have been described. Since 2010, there has been a sustained epidemic that retains the cyclical pattern, although at a level higher than in previous years. The incidence has been increasing in infants under 1 year, who continue to be the most affected. Incidence rates (IRs) per 100 000 population were 150.6 and 119.5 cases in 2017 and 2018, respectively, followed by IRs for children aged 1–4 years (56.7 and 39.5 in 2017 and 2018, respectively), 5–9 years (62.3 and 30.3), and 10–14 years (37.1 and 33.0).¹⁵ These data suggest a progressive accumulation of individuals susceptible to infection due to decreased immunity after years of low incidence and highlight the need for new vaccination strategies and more effective vaccines to prevent pertussis in unvaccinated children, in whom the disease may present more severely.¹⁶

The pertussis hospitalization rate follows an increasing epidemic pattern similar to that of the IR in recent years, with the highest rate in children <1 year, especially in infants under 3 months.¹⁷ Vaccination has significantly reduced hospital admissions.¹⁸ The annual rate of hospitalizations in Spain per 100 000 population in the period 1997–2010 was 131.02 cases in infants under 1 year, increasing to 250.13 in the period 2011–2015 and decreasing to 157.69 in 2016–2017. Most cases were recorded in infants ≤ 4 months, with hospitalization rates of 328.80, 670.81, and 385.84 cases/100 000 population in the periods 1997–2010, 2011–2015, and 2016–2017, respectively. In total, 34, 36 and 4 deaths were reported in the three periods, respectively. Following the implementation of maternal vaccination by region, a 20% reduction in the rate of hospitalizations was observed¹⁹ (Fig. 1).

Microbiological aspects of pertussis

Population dynamics and antigenic drift

The resurgence of pertussis has been attributed to several factors, with waning immunity and adaptation of the pathogen being particularly relevant.²⁰ The introduction of acellular vaccines may have contributed to the emergence of new antigenic variants of *B. pertussis* capable of evading the vaccine-induced immune response, and strains that overproduce pertussis toxin (PT) or are deficient in pertactin (PRN), a protein essential for bacterial adhesion to the upper respiratory tract epithelium which is present in




Fig. 1. Annual rate of hospitalization associated with pertussis per 100 000 children according to the year of introduction of maternal vaccination in Spain updated with 2018 data¹⁹.

some vaccines.^{20,21} In Spain, antigen typing of *B. pertussis* strains isolated over the past 30 years has confirmed that after the transition from whole-cell to acellular vaccines, bacterial lineages that produce vaccine antigens different to those contained in current vaccines have emerged, as has been also confirmed in neighbouring countries.^{22,23}

Microbiological diagnosis

The resurgence of pertussis highlights the need for rapid microbiological diagnosis that allows early implementation of adequate antibiotic therapy and the necessary measures to prevent bacterial transmission. This is also essential to identify changes in the disease due to modifications in vaccine effectiveness, waning immunity, the emergence of *B. pertussis* strains that can evade the immunity conferred by the vaccine, and to identify the emergence of other *Bordetella* species that cause clinical symptoms similar to those produced by *B. pertussis*, and for which the vaccines currently employed are not useful.²⁴

Culture is the gold standard for the diagnosis of pertussis, although nucleic acid amplification techniques (NAATs) such as polymerase chain reaction (PCR) or isothermal amplification have become the most widely used diagnostic tools owing to their speed and excellent sensitivity.²⁵ Moreover, NAATs do not depend on the bacterial viability, so bacteria can be detected even in cases of persistent cough lasting more than three weeks or after patients have been treated with antibiotics. In addition, most of the currently available NAATs can detect in the same test not only *B. pertussis* but also *Bordetella* species that cause pertussis-like disease, such as *B. parapertussis*, *B. bronchiseptica* or the recently described *B. holmesii*, which has been associated with major outbreaks and identified in 4.1% of laboratory-confirmed whooping cough cases between 2013 and 2016 in Barcelona.²⁶ As macrolides show lower activity against *B. holmesii* than against other species, and that acellular vaccines do not confer cross-protection against this or other species such as *B. parapertussis*, accurate diagnosis of the pertussis-causing agent is needed in order to establish appropriate antimicrobial therapy, to determine the real incidence of pertussis and to assess the efficacy of the pertussis vaccines currently administered.^{27,28}

Serological diagnosis is particularly indicated in advanced stages of the disease where both NAATs and culture offer less sensitivity. Titration of immunoglobulin Gs (IgGs) generated against PT provides better results, since the toxin is exclusively produced by *B. pertussis* and there is no cross-reactivity with similar antigens present in other microorganisms. However, serological diagnosis is not recommended during the year following vaccination, since the immune response induced by PT included in the vaccine cannot be differentiated from the response to natural infection.^{29,30}

Antimicrobial resistance

Macrolides are the antibiotics recommended for both pertussis treatment and post-exposure chemoprophylaxis. Among them, the most widely used are azithromycin and clarithromycin, because they facilitate therapeutic adherence and eradicate the microorganism as effectively as erythromycin, with a high safety profile. In the case of neonates (under 1 month of age), azithromycin is the recommended antibiotic due to its good tolerance and high safety profile, while erythromycin has been associated with infantile hypertrophic pyloric stenosis.³¹

Over the past 20 years, *B. pertussis* strains that are resistant to macrolides have emerged as a result of mutations in their 23S ribosomal RNA encoding gene. Since the first description in the United States (US) in 1995, sporadic cases have been described in France, Iran and Japan.^{24,32,33} However, the high prevalence of macrolide-resistant *B. pertussis* detected in China since 2013 (>90%) is alarming.^{34,35} Accordingly, macrolide resistance in *B. pertussis* in Spain should be monitored to promptly identify any potential emergence and to establish the necessary measures to prevent its spread.

Prevention and current vaccine types

After the first vaccine authorized in the US in 1914, whole-cell pertussis (wP) vaccines combined with tetanus (T) and diphtheria (D) toxoids (DTwP vaccines) became available from 1948 onwards. Combined vaccines with inactivated poliovirus (IPV), *Haemophilus influenzae* type b (Hib) and hepatitis B (HepB) viral antigens were subsequently developed.

In Spain, inactivated acellular vaccines (DTaP) have been marketed since 1998, and have been progressively replacing DTwP vaccines due to their lower reactogenicity. Depending on the preparation, DTaPs may contain 1–5 *B. pertussis* antigens: PT (present in all), filamentous haemagglutinin (FHA), PRN, and one or two types of fimbrial proteins (FIM-2 and FIM-3). Multiple studies show that none of these factors alone is a determinant of vaccine efficacy, although most authors agree that the fundamental component is the PT that directly induces the generation of protective antibodies in response to immunization.^{36,37}

DTaP vaccines: immunogenicity, efficacy, effectiveness and safety

Differences among the populations analyzed, case definition criteria and the serological tests used make difficult to compare the immunogenicity and efficacy studies of the various DTaP vaccines. In Germany, only PT and PRN were found to be significantly associated with immune protection.³⁸ Two other clinical efficacy trials in Sweden showed that after exposure to *B. pertussis*, the concentration of antibodies against the vaccine antigens was higher than that generated against other bacterial proteins not contained in the vaccine.³⁹ Several clinical studies have estimated efficacies of about 84–85% and 71–78% in the prevention of typical and mild pertussis, respectively, for acellular vaccines formulated with at least three bacterial antigens.^{40,41} Simondon and colleagues estimated the efficacy of a 2-component vaccine at 85%, which led to its approval within the regulatory framework and implementation in vaccination programmes.⁴² A single-component vaccine (PT) is still in use in northern Europe, with an efficacy of 71%.⁴³ However, the comparison of efficacy between wP and aP vaccines or between different aP vaccines has not led to clear conclusions, and there is no evidence to support the superiority of one aP vaccine over another.^{44,45}

Most of the acellular vaccines currently available have shown their efficacy-effectiveness in various clinical trials and are currently used in national immunization programmes. Table 1 shows the acellular vaccines available and marketed in Spain.

In any case, the effectiveness of aP vaccines against pertussis is not optimal, and factors such as waning immunity over time and expansion of clones with antigenic polymorphism could explain why their effectiveness is the lowest within paediatric vaccines. Systemic and local adverse effects are generally less common with aP vaccines, although the rate and intensity of minor local adverse reactions tend to increase with each booster dose of DTaP.^{40,41,46} DTaP vaccines combined with other antigens (DTaP-IPV; DTaP-IPV/Hib; DTaP-IPV/Hib/HepB) have shown seroprotection and efficacy not inferior to non-combination vaccines and a safety profile comparable to that offered by separate administration of any of the components.^{47–51}

Protection throughout life: new vaccination strategies

The reduced effectiveness of acellular vaccines compared to other vaccines and the waning of immunity after the primary vaccination, inherent in both whole-cell and acellular vaccines underline the advisability of designing global prevention strategies that protect throughout life.^{6,52}

Vaccination during pregnancy

Immunization during pregnancy is currently considered the most effective and efficient vaccination strategy to protect infants from severe pertussis and associated mortality in the first months of life before they begin their primary vaccination course.⁵³ This approach protects the newborn (NB) both directly by preventing

infection and transmission from the mother, and indirectly by passive transplacental transfer of antibodies to the foetus, which will protect the baby until the first dose of vaccine is received at 2 months of age. There is a good correlation between maternal post-vaccine IgG₁ concentrations and those detected in the umbilical cord, which has even higher levels. Thus, transplacental antibody transfer can protect the NB if the placenta is healthy.⁵⁴

Early immunization during the second trimester of pregnancy has been associated with higher titres of transferred antibodies against PT and FHA than estimated when vaccination is given from gestational week 25.⁵⁵ Anti-PT IgG avidity to PT in cord blood is higher in mothers vaccinated early in the third trimester than in those vaccinated later.⁵⁶ However, a higher level of scientific evidence is still needed to determine the optimal time for immunization in pregnancy.

Maternal immunization with low-dose acellular pertussis vaccine (Tdap)^{57,58} has been shown to be safe, with no increased risk of maternal or foetal adverse effects (such as prematurity, premature rupture of membranes, intrauterine growth retardation, endometritis, or congenital malformations). In Spain, three acellular vaccines with a low antigenic load of pertussis component (Tdap) (Table 1) are indicated for protection not only during pregnancy, but also as a booster after primary immunization in children, adolescents and adults.

A small increase in the incidence of chorioamnionitis has been reported in vaccinated pregnant women, although this finding has not yet been corroborated by other authors.⁵⁹ The main efficacy/effectiveness studies of vaccination in pregnancy for protection of the infant are shown in Table 2.

One aspect to consider is the potential interference of maternal antibodies with the infant's immune response to primary vaccination (immunological blunting), a phenomenon observed both after natural infection (e.g., measles, polio) and after vaccination (with significant variations between different vaccines and published papers).⁶⁰ Although a lower rate of humoral response to the primary vaccination against pertussis has been described in infants of mothers vaccinated during pregnancy, this effect disappears after administration of the booster dose.^{61–63} The clinical relevance of these data is unknown, as countries with well-established maternal immunization programmes have not observed an increase in pertussis cases in infants or older children. A lower titre of anticapsular antibodies against some vaccine serotypes of *Streptococcus pneumoniae* has also been described, probably due to the effect of the non-toxic mutant of diphtheria toxin (CRM197) used as a carrier protein in the conjugated vaccines. Nevertheless, the percentages of seroprotection in infants of vaccinated and unvaccinated mothers were comparable after the primary vaccination and booster doses, except for serotype 3.⁶⁴

The US was the first country to introduce maternal vaccination in 2011, with a Tdap dose between weeks 27 and 36 of gestation and revaccination in every pregnancy. The estimated vaccine effectiveness in pertussis prevention during the first two months of life of infants was 91.4% (95% confidence interval [CI] 19.5–99.1%), and 69% (95% CI 43.6–82.9%) during the first 12 months of life.⁶⁵ The United Kingdom (UK) incorporated it in 2012, reaching 64% coverage in the first year of implementation and 93% vaccine effectiveness in pertussis prevention during the first two months of life.⁶⁶ According to 2019 data, maternal vaccination is already in place in 18 countries in Europe, although the gestational time at which it is administered differs (Table 3).

In Spain, the Public Health Commission approved the anti-pertussis maternal vaccination programme in 2015,⁶⁷ although it had already been implemented in some autonomous regions (like Catalonia and Asturias) since 2014. The vaccination schedule includes a dose of combined acellular vaccine with reduced diphtheria and pertussis toxoids content (Tdap) between weeks 27–28

Table 1
Composition of pertussis vaccines commercially available in Spain.⁷²

Product (manufacturer)	Type	Composition	Presentation
Boostrix (GSK)	Tdap	Three purified <i>B. pertussis</i> antigens: 8 µg PT, 8 µg FHA, 2.5 µg PRN ≥ 20 IU TT, ≥2 IU DT Adjuvant: aluminium hydroxide and phosphate	0.5 ml pre-filled syringe
Boostrix Polio (GSK)	Tdap	Same composition as Boostrix® + 3 types of IPV	0.5 ml pre-filled syringe
Hexyon (Sanofi Pasteur)	DTaP	Two purified <i>B. pertussis</i> antigens: 25 µg PT, 25 µg FHA, ≥20 IU DT, 12 µg Hib oligosaccharide conjugated with ≥40 IU TT + 10 µg HBsAg + 3 types of IPV Adjuvant: aluminium hydroxide	0.5 ml pre-filled syringe
Infanrix Hexa (GSK)	DTaP	Three purified <i>B. pertussis</i> antigens: 25 µg PT, 25 µg FHA, 8 µg PRN, +≥30 IU DT, 10 µg Hib polysaccharide conjugated with 40 IU TT + 10 µg HBsAg + 3 types of IPV Adjuvant: aluminium hydroxide and phosphate	Lyophilized powder in a vial + 0.5 ml of suspension in a pre-filled syringe
Triaxis (Sanofi Pasteur)	Tdap	Five purified <i>B. pertussis</i> antigens: 2.5 µg PT, 5 µg FHA, 3 µg PRN, 5 µg fimbriae types 2 and 3, ≥20 IU TT, ≥2 IU DT Adjuvant: aluminium phosphate	0.5 ml pre-filled syringe
Vaxelis (MSD)	DTaP	Five purified <i>B. pertussis</i> antigens: 20 µg PT, 20 µg FHA, 3 µg PRN, 5 µg fimbriae types 2 and 3, ≥20 IU DT, 3 µg Hib polysaccharide conjugated with ≥40 IU TT + 10 µg HBsAg + 3 types of IPV Adjuvant: aluminium hydroxide and phosphate	0.5 ml pre-filled syringe

DT, diphtheria toxoid; FHA, filamentous haemagglutinin; HBsAg, hepatitis B surface antigen; Hib, *Haemophilus influenza* type b; IPV, inactivated poliomyelitis virus; PRN, pertactin; PT, pertussis toxoid; TT, tetanus toxoid.

Table 2
Main studies of the effectiveness of pertussis vaccination programmes in pregnant women.

Author	Year of publication	Country	Effectiveness (95% CI)
Amirthalingam et al. ⁶⁶	2014	United Kingdom	91% (83–95%)
Dabrera et al., ⁹⁸	2015	United Kingdom	93% (81–97%)
Baxter et al., ⁶⁵	2017	USA	91.4% (19.5–99.1%)
Winter et al. ⁹⁹	2017	USA	85% (33–98%)
Bellido-Blasco et al. ¹⁰⁰	2017	Spain	90.9% (56.6–98.1%)
Saul et al., ¹⁰¹	2018	Australia	69% (13–89%)
Fernandes et al., ¹⁰²	2019	Brazil	82.6% (60.8–92.3%)
Sancho-Uriarte et al., ¹⁰³	2019	Spain	89% (72–96%)
Romanin et al., ¹⁰⁴	2020	Argentina	80.7% (52.1–92.2%)

Table 3
Recommendations for pertussis vaccination in pregnancy in Europe (data as of November 2019).

Country	Programme (start year)	Gestation time (week)
United Kingdom	2012	16+
Ireland	2012	16+
Austria	2013	13+
Liechtenstein	2013	13+
Swiss	2013	13+
Belgium	2013	24–32
Spain	2014	27–36
Czech Republic	2016	28+
Portugal	2017	20+
Slovenia	2017	24+
Greece	2017	27+
Italy	2017	28+
Poland	2017	28+
Luxembourg	2019	13+
Netherlands	2019	22+
Iceland	2019	28+
Denmark	2019	32+

+, starting from the week indicated.

of pregnancy; if there is risk of prematurity, the Spanish Association of Paediatrics (AEP) also recommends vaccination from week 16, preferably at week 20, after the scheduled 20-week ultrasound.^{68,69} The vaccine should be given in each pregnancy regardless of previous vaccination status. In Spain, current vaccination coverage in pregnant women is around 83%.⁷⁰ The introduction of vaccination in pregnancy resulted in a reduction in the incidence of pertussis hospitalizations in infants, with significantly lower figures in the regions that adopted this strategy earlier¹⁹ (Fig. 1).

Primary vaccination

The World Health Organization (WHO) recommends starting vaccination at 6–8 weeks of life and considers that it is essential not to delay its administration.⁷¹ The interval between primary vaccination doses should be ≥1 month, and the booster dose should be given at least 6 months after the second dose.

In Spain, pertussis vaccination began in 1975, with three doses of whole-cell vaccine before age 6–7 months. Doses at 15–18 months and 4–6 years were introduced in 1996 and 2001, respectively. Acellular vaccines subsequently replaced whole-cell vaccines, especially for the dose given at 18 months of age, due to their lower reactogenicity. Administration of a fifth dose at 4–6 years of age was incorporated into the vaccination schedule in 2005. In 2012, the second booster dose was replaced by the low-dose acellular vaccine (Tdap).

Since 2017, the Interterritorial Council of the Spanish National Health Service (CISNS) has recommended primary vaccination with hexavalent standard-dose vaccines (DTaP) at 2 and 4 months of age, and a booster dose at 11 months in a 2 + 1 schedule that remains in force today.⁶⁹ To complete the vaccine schedule, a fourth dose should be given at 6 years of age with a combined vaccine (DTaP-IPV or Tdap-IPV). The fifth dose should be given at 12–14 years with tetanus and diphtheria vaccine, Td, although the Tdap vaccine has been administered in Asturias since 2017, as recommended by the AEP.^{68,72} Similar vaccine schedules, with slight variations in the number of doses and the time at which they are administered, are used in other European Union countries.⁷³ Since 2020, the Centers for Disease Control (CDC) have authorized the use of Td or Tdap for tetanus, diphtheria, and pertussis vaccination in children > 7 years old who are not vaccinated or who require booster doses.

Table 4

Vaccination coverage during routine immunization programmes with 1 or 3 doses of the DTP vaccine in the World Health Organization (WHO) regions in 2019.⁷⁹

Region	DTP1	DTP3
Africa	81%	74%
America	90%	84%
Eastern Mediterranean	89%	82%
Europe	97%	95%
South-East Asia	94%	91%
Western Pacific	95%	94%

Data are available on the impact of the administration of the first dose of pertussis vaccine in NB at 6 or 8 weeks, but few studies have compared immunogenicity and safety according to the time of administration.⁷⁴ A phase III clinical trial of a monovalent acellular vaccine in NB confirmed satisfactory immunogenicity and safety, suggesting that it could be an alternative to protecting the child against pertussis in cases where the vaccine had not been administered during pregnancy.⁷⁵ Preterm NBs (PNBs) have specific cellular immune responses. In the case of pertussis, no serological surrogate pattern of protection has been established. It has been observed that after the administration of combined vaccines with a 2 + 1 or 3 + 1 schedule, the IgG immune response of PNBs may be somewhat inferior to term NBs (TNBs). Even so, after the booster dose, antibody titres recover to similar levels. PNBs should complete the vaccination schedule based on their chronological age, following the same vaccination guidelines as TNBs.^{68,76,77} The hexavalent vaccines currently available in Spain can be used to vaccinate premature infants, although the potential risk of apnea must be evaluated and respiratory monitoring should be performed for 48–72 h in very premature children (born at ≤ 28 weeks of gestation) and in particular, in infants with a history of respiratory immaturity.⁷⁸

According to 2019 data, 90% of children received at least one dose of DTaP and 85% received three doses⁷⁹ (Table 4). In Spain, in 2018, all autonomous regions except for the Balearic Islands (which did not provide information) reported coverage of more than 95% of the cohort for the first two doses of hexavalent vaccine during the first year. However, for the third dose at 11 months, six communities presented overall coverages slightly below 95.5%, with the lowest being in the Basque Country (92.5%).⁷⁰

Despite the high coverage, the duration of the immunity conferred by primary vaccination (as with natural immunity after infection) is limited, a feature common to all types of vaccines (cellular and acellular).⁸⁰ For each year after the last dose of acellular vaccine, there is an estimated 27–33% increase in the chances of developing pertussis, and that only 10% of infants vaccinated with standard-dose vaccines (DTaP) remain protected 8.5 years after the last dose.^{80,81} This justifies the need for booster doses at preschool, adolescent and adult age.

Preschool-age vaccine booster

Vaccines and vaccine schedules used in primary vaccination can influence acquired immune protection. Consequently, preschool vaccination is essential for reducing the risk of pertussis in an environment where transmissibility conditions are very high. No significant differences have been found so far in the duration of immune protection acquired after a childhood vaccination series with 3 or 5 doses of DTaP.⁸⁰ Nevertheless, published data in Australia showed that removing the booster dose at 18 months in 2003 resulted in an increase in pertussis notification rates in children aged 1–4 years, which translated to an increase in the prevalence of undetectable IgG levels from 1997 to 1998 to 2007.⁸³ Continuous epidemiological surveillance is therefore essential to assess

the impact of changes in vaccine guidelines on the incidence of pertussis in children.

The gradual decline in the protection acquired after the primary vaccination justifies the need for a booster in preschool age. It has been estimated that the odds of acquiring pertussis would increase almost linearly with age by an average of 42% for each year after the last dose a DTaP vaccine. The amount of protection depends on the effectiveness of the DTaP vaccine. An initial DTaP effectiveness of 95% would decrease to 71% after 5 years of last dose of vaccine and 90% effectiveness would decrease to 42% in the same period of time.⁸⁴ In children aged 5–9 years, effectiveness would exceed approximately 75%, and more than 65% of children would remain immune to pertussis 5 years after the last dose of DTaP.⁸⁵

Although low antigen load vaccines have demonstrated adequate antibody responses and lower reactogenicity than standard-dose preparations, the absence of studies evaluating the duration of acquired protection in preschool and school-age children ≥ 4 years prioritizes the use of DTaP vaccines in the booster dose. The estimated age for receiving this dose is 3–7 years. In Spain, the combined standard load vaccine (DTaP-IPV) is recommended whenever possible, and preferably the Tdap-IPV in the corresponding dose at 6 of age.⁶⁹ Official agencies, such as the WHO and the CDC, also recommend the use of standard-dose preparations (DTaP-IPV) versus low-dose preparations (Tdap-IPV) for booster doses, as they offer greater immunogenicity and comparable reactogenicity.^{86,87}

Vaccination of adolescents and adults

Loss of the immunogenicity acquired after vaccination makes adolescents and adults susceptible to re-infection, thus becoming potential sources of pertussis transmission to children and infants with whom they are in contact. The vaccination strategy for adolescents and adults has demonstrated its efficacy, safety and benefit.^{88,89} In the US, Tdap vaccination in adolescents has been effective in reducing the number of pertussis hospitalizations in infants.⁹⁰ The AEP and agencies such as the CDC recommend vaccination with Tdap in adolescence (12–14 years).^{72,91,92} Currently, in Spain, Tdap vaccination at the age of 13 is only funded in Asturias. Most European Union countries include adolescent vaccination in their vaccine schedules.⁷³

The recommendations of the Spanish Ministry of Health for adult vaccination are limited to pregnant women, health professionals working in paediatrics, obstetrics and their respective emergency areas, and professionals working in close contact with cases and in the control of pertussis epidemic outbreaks. These measures are supported by the AEP, which also proposes that child-care providers be included in the programme.^{93,94} In Spain, the approved vaccines for adolescents and adults are Boostrix and Triaxis (Table 1).

Vaccine boosters in adulthood can strengthen the level of immunization of the population and thus reduce bacterial circulation.⁶ The CDC has recently authorized the use of the Tdap vaccine in any adult situation, as a replacement or as a possible alternative to Td in cases where previously only Td was indicated for tetanus wound prophylaxis.⁹⁵ In Spain, the current tetanus vaccination recommendation considers that the administration of 5 doses of Td throughout life is sufficient to protect against tetanus up to the age of 65, but this strategy fails to provide pertussis booster immunization in adults.⁹⁶ In the European Union, several countries recommend a pertussis vaccine booster in adults every 10 years after completing the childhood immunization schedules.⁷³

The proportion of older people is constantly increasing worldwide, especially in Europe, as is the risk of infectious diseases whose severity increases with age. Older people nowadays provide valuable family support by caring for infants and children, so this is

a population group with a particular risk of infection and, consequently, of transmission to their setting. It would therefore be particularly advisable to consider extending the benefits of pertussis vaccination to this risk group.

Cocooning strategy

The cocooning strategy is based on the vaccination of individuals in the child's immediate environment (parents, grandparents, siblings, others who live with the child, etc.) to avoid disease transmission. Despite the potential difficulty implementing this strategy, it protects children whose mothers were not vaccinated during pregnancy. In countries such as Australia, France, Germany and the US, this measure was introduced in 2000. However, since the vaccine coverage achieved is suboptimal in most cases, this strategy is not sufficient in itself to prevent childhood infection and consequently, hospitalizations or deaths.^{71,97}

Working Group recommendations

All data presented reinforce the need for a comprehensive prevention strategy to protect all age groups susceptible to pertussis infection and transmission, that should include the following key points:

- Rapid, sensitive and specific microbiological diagnosis.
- Surveillance of the vaccine antigen variants of *B. pertussis* and susceptibility of *B. pertussis* to macrolides, as well as the emergence of other *Bordetella* species that can produce pertussis-like pathology.
- Vaccination in pregnancy with Tdap as a double measure of protection for mothers and infants under 2 months.
- Compliance with the current 2 + 1 primary vaccination schedule with hexavalent vaccines at 2 and 4 months of age, and booster at 11 months. Vaccination of PNBs (which can be done with hexavalent vaccines) must be completed according to the infants' chronological age, following the same vaccination schedule as TNBs.
- Booster vaccination in preschool-age children as protection against the risk of infection associated with the loss of vaccine immunity achieved after the primary vaccination.
- Vaccination of adolescents to reduce the risk associated with loss of immunity.
- The implementation of complementary measures to the vaccination programme such as the cocooning strategy and the vaccination of adults and healthcare professionals.

Funding

The authors disclose receipt of the following financial support for the research, authorship, and publication of this article: Sanofi Pasteur (Madrid, Spain) supported this work but had no role in the preparation of the manuscript.

Conflict of interests

Juan José González-López has received compensation for consultant services for Sanofi Pasteur, has received research support from Sanofi Pasteur and GlaxoSmithKline (GSK) and has collaborated in educational activities funded by Sanofi Pasteur and Merck Sharp & Dohme (MSD). Javier Álvarez Aldeán has collaborated in educational activities funded by GSK, Pfizer, and Sanofi, has received compensation for services as consultant for Sanofi-Pasteur, GSK, MSD, and Pfizer and has participated as researcher in clinical trials funded by GSK, Novartis, and Sanofi Pasteur.

Francisco José Álvarez García has collaborated in educational activities funded by GSK, MSD, Pfizer, and Sanofi Pasteur and has received compensation for services as consultant for Sanofi-Pasteur, GSK, MSD, and Pfizer. Magda Campins has participated as researcher in vaccine clinical trials funded by GSK, Novartis, Sanofi Pasteur and Novavax, and has received compensation for consultant services for Sanofi Pasteur, GSK, Pfizer, Novartis, and MSD. María Garcés-Sánchez has participated as researcher in vaccine clinical trials from GSK, Novartis, Sanofi Pasteur, and Pfizer, and has received compensation for consultant services for Sanofi Pasteur, GSK, Pfizer, and MSD. Ruth Gil-Prieto has received research grants from Merck, Sanofi Pasteur and Pfizer and has received compensation for services as consultant for Sanofi Pasteur. Ana M^a Grande-Tejada has collaborated in educational activities funded by GSK, MSD, Pfizer, and Sanofi Pasteur and has received compensation for services as consultant for Sanofi Pasteur GSK, MSD, Pfizer, and Novartis.

Acknowledgements

The authors thank Susana Cañón (Medical Statistics Consulting, S.L., Valencia, Spain) for providing medical writing support in accordance with Good Publication Practice (GPP3) guidelines, which was funded by Sanofi Pasteur (Madrid, Spain).

References

1. European Centre for Disease Prevention and Control. Pertussis. In: ECDC. Annual epidemiological report for 2018; 2020. <https://www.ecdc.europa.eu/en/publications-data/pertussis-annual-epidemiological-report-2018>.
2. Solano R, Crespo I, Fernandez MI, Valero C, Alvarez MI, Godoy P, et al. Underdetection and underreporting of pertussis in children attended in primary health care centers: do surveillance systems require improvement? *Am J Infect Control*. 2016; 44:e251–e256.
3. Parliament E. Commission Implementing Decision 2018/945/EU of 22 June 2018 on the communicable diseases and related special health issues to be covered by epidemiological surveillance as well as relevant case definitions; 2018.
4. Domínguez A, Soldevila N, Cayla JA, García-Cenoz M, Ferrus G, Sala-Farre MR, et al. Assessment of clinical symptoms in household contacts of confirmed pertussis cases. *J Infect*. 2017;75:426–32.
5. Fernandez-Cano MI, Armadans Gil L, Martinez Gomez X, Campins Marti M. Incidence of whooping cough in Spain (1997–2010): an underreported disease. *Eur J Pediatr*. 2014;173:721–6.
6. Martinon-Torres F, Heininger U, Thomson A, Wirsing von König CH. Controlling pertussis: how can we do it? A focus on immunization. *Expert Rev Vaccines*. 2018;17:289–97.
7. Gianfredi V, Bragazzi NL, Mahamid M, Bisharat B, Mahroum N, Amital H, et al. Monitoring public interest toward pertussis outbreaks: an extensive Google Trends-based analysis. *Public Health*. 2018;165:9–15.
8. Bruguera S, Rius C, Millet JP, Casals M, Cayla JA, Barcelona Pertussis Working G. Does the economic recession influence the incidence of pertussis in a cosmopolitan European city? *BMC Public Health*. 2019;19:144.
9. Gil-Tomas JJ, Colomina-Rodríguez J, Martínez-Macias O, Borrás-Manez M, Guerrero-Espejo A. Red de Vigilancia Epidemiológica de la Comunidad V. *Enferm Infecc Microbiol Clin*. 2013;31:590–4.
10. Latasa P, García-Comas L, Gil de Miguel A, Barranco MD, Roderio I, et al. Effectiveness of acellular pertussis vaccine and evolution of pertussis incidence in the community of Madrid from 1998 to 2015. *Vaccine*. 2018;36:1643–9.
11. Martínez Lorenzo R, González Pineiro Y, López Vázquez AM, Trigo Daporta M. Increase in the number of diagnosed cases of whooping cough; do we think more in it? *Aten Primaria*. 2017;49:309–11.
12. Miguez Santiyan A, Ferrer Estrems R, Chover Lara JL, Alberola Enguidanos J, Nogueira Coito JM, Salazar Cifre A. Early intervention in pertussis outbreak with high attack rate in cohort of adolescents with complete acellular pertussis vaccination in Valencia, Spain, April to May 2015. *Euro Surveill*. 2015;20.
13. Porras-Povedano M, Roldán-Garrido A, Santacruz-Hamer V. Outbreak of Whooping cough in 2016, Ecija, Seville, Spain. *Rev Esp Salud Publica*. 2017;91.
14. Solano R, Rius C, Simon P, Manzanares-Laya S, Ros M, Toledo D, et al. Evaluation of reported cases of pertussis: epidemiological study in a large city in Spain. *J Med Microbiol*. 2014;63:1688–95.
15. Centro Nacional de Epidemiología. Instituto de Salud Carlos III. Resultados de la vigilancia Epidemiológica de las enfermedades transmisibles. Informe anual. Años 2017–2018. Madrid, 2020.
16. Moraga-Llop FA, Mendoza-Palomar N, Muntaner-Alonso A, Codina-Grau G, Fabregas-Martori A, Campins-Martí M. Pertussis in fully vaccinated infants and children. Are new vaccination strategies required? *Enferm Infecc Microbiol Clin*. 2014;32:236–41.

17. Centro Nacional de Epidemiología. CIBERESP. ISCIII. RENAVE. Situación de la Tos ferina en España, 1998–2016. Análisis preliminar del Impacto de la Vacunación de Tos ferina en embarazadas. Madrid, 25 de abril de 2018.
18. Crespo I, Toledo D, Soldevila N, Jordan I, Solano R, Castilla J, et al. Characteristics of hospitalized cases of pertussis in Catalonia and Navarra. Two regions in the North of Spain. *PLoS One*. 2015; 10:e0139993.
19. Gil-Prieto R, Walter S, San-Roman-Montero J, Marin-García P, Gonzalez-Escalada A, Gil-de-Miguel A. Paediatric hospitalizations due to whooping cough in Spain (1997–2017). *Vaccine*. 2019;37:6342–7.
20. Mooi FR, Van Der Maas NA, De Melker HE. Pertussis resurgence: waning immunity and pathogen adaptation – two sides of the same coin. *Epidemiol Infect*. 2014;142:685–94.
21. Martin SW, Pawloski L, Williams M, Weening K, DeBolt C, Qin X, et al. Pertactin-negative *Bordetella pertussis* strains: evidence for a possible selective advantage. *Clin Infect Dis*. 2015;60:223–7.
22. van Gent M, Heuvelman CJ, van der Heide HG, Hallander HO, Advani A, Guiso N, et al. Analysis of *Bordetella pertussis* clinical isolates circulating in European countries during the period 1998–2012. *Eur J Clin Microbiol Infect Dis*. 2015;34:821–30.
23. Mir-Cros A, Moreno-Mingorance A, Martin-Gomez MT, Codina G, Cornejo-Sanchez T, Rajadell M, et al. Population dynamics and antigenic drift of *Bordetella pertussis* following whole cell vaccine replacement, Barcelona, Spain, 1986–2015. *Emerg Microbes Infect*. 2019;8:1711–20.
24. Kilgore PE, Salim AM, Zervos MJ, Schmitt HJ. Pertussis: microbiology, disease treatment, and prevention. *Clin Microbiol Rev*. 2016;29:449–86.
25. Lee AD, Cassidy PK, Pawloski LC, Tatti KM, Martin MD, Briere EC, et al. Clinical evaluation and validation of laboratory methods for the diagnosis of *Bordetella pertussis* infection: Culture, polymerase chain reaction (PCR) and anti-pertussis toxin IgG serology (IgG-PT). *PLOS ONE*. 2018; 13:e0195979.
26. Mir-Cros A, Codina G, Martin-Gomez MT, Fabrega A, Martinez X, Jane M, et al. Emergence of *Bordetella holmesii* as a causative agent of Whooping Cough, Barcelona, Spain. *Emerg Infect Dis*. 2017;23:1856–9.
27. Pittet LF, Emonet S, Schrenzel J, Siegrist CA, Posfay-Barbe KM. *Bordetella holmesii*: an under-recognised *Bordetella* species. *Lancet Infect Dis*. 2014;14:510–9.
28. Zhang X, Weyrich LS, Lavine JS, Karanikas AT, Harvill ET. Lack of cross-protection against *Bordetella holmesii* after pertussis vaccination. *Emerg Infect Dis*. 2012;18:1771–9.
29. Cherry JD, Tan T, Wirsing von Konig CH, Forsyth KD, Thisyakorn U, et al. Clinical definitions of pertussis: summary of a Global Pertussis Initiative roundtable meeting, February 2011. *Clin Infect Dis*. 2012;54:1756–64.
30. van der Zee A, Schellekens JF, Mooi FR. Laboratory diagnosis of pertussis. *Clin Microbiol Rev*. 2015;28:1005–26.
31. Tiwari T, Murphy TV, Moran J. National Immunization Program CDC. Recommended antimicrobial agents for the treatment and postexposure prophylaxis of pertussis: 2005 CDC Guidelines. *MMWR Recomm Rep*. 2005;54:1–16.
32. Guillot S, Descours G, Gillet Y, Etienne J, Floret D, Guiso N. Macrolide-resistant *Bordetella pertussis* infection in newborn girl, France. *Emerg Infect Dis*. 2012;18:966–8.
33. Yamaguchi T, Kawasaki Y, Katsukawa C, Kawahara R, Kawatsu K. The first report of isolation of macrolide-resistant *Bordetella pertussis* in Japan. *Jpn J Infect Dis*. 2020;73:361–2.
34. Wang Z, Cui Z, Li Y, Hou T, Liu X, Xi Y, et al. High prevalence of erythromycin-resistant *Bordetella pertussis* in Xi'an, China. *Clin Microbiol Infect*. 2014;20:0825–30.
35. Yang Y, Yao K, Ma X, Shi W, Yuan L, Yang Y. Variation in *Bordetella pertussis* susceptibility to erythromycin and virulence-related genotype changes in China (1970–2014). *PLOS ONE*. 2015; 10:e0138941.
36. Gabutti G, Azzari C, Bonanni P, Prato R, Tozzi AE, Zanetti A, et al. Pertussis. *Hum Vaccin Immunother*. 2015;11:108–17.
37. Edwards KM, Decker MD. 44 – Pertussis vaccines. In: Plotkin SA, Orenstein WA, Offit PA, Edwards KM, editors. *Plotkin's Vaccines*. 7th ed. Elsevier; 2018, 711–761.e16.
38. Stehr K, Cherry JD. A comparative efficacy trial in Germany in which infants received either the Lederle/Takeda acellular pertussis component DTP (DTaP) vaccine, the Lederle whole-cell component DTP (DTP) vaccine or DT vaccine. *Dev Biol Stand*. 1997;89:58–62.
39. Cherry JD, Heininger U, Richards DM, Storsaeter J, Gustafsson L, Ljungman M, et al. Antibody response patterns to *Bordetella pertussis* antigens in vaccinated (primed) and unvaccinated (unprimed) young children with pertussis. *Clin Vaccine Immunol*. 2010;17:741–7.
40. Zhang L, Prietsch SO, Axelsson I, Halperin SA. Acellular vaccines for preventing whooping cough in children. *Cochrane Database Syst Rev*. 2012;CD001478.
41. Jefferson T, Rudin M, DiPietrantonj C. Systematic review of the effects of pertussis vaccines in children. *Vaccine*. 2003;21:2003–14.
42. Simondon F, Preziosi MP, Yam A, Kane CT, Chabirand L, Iteman I, et al. A randomized double-blind trial comparing a two-component acellular to a whole-cell pertussis vaccine in Senegal. *Vaccine*. 1997;15:1606–12.
43. Pertussis surveillance in Sweden: Nineteen-year report; 2016. <https://www.folkhalsomyndigheten.se/contentasset/65ed8fdbdab4999bc358fcd9b657e77/pertussis-sweden-nineteen-year-report.pdf>.
44. Esposito S, Stefanelli P, Fry NK, Fedele G, He Q, Paterson P, et al. Pertussis prevention: reasons for resurgence, and differences in the current acellular pertussis vaccines. *Front Immunol*. 2019;10:1344.
45. Harriman K, Healy CM. How important is the type of acellular pertussis vaccine? *Clin Infect Dis*. 2020;70:208–9.
46. Decker MD, Edwards KM, Steinhoff MC, Rennels MB, Pichichero ME, Englund JA, et al. Comparison of 13 acellular pertussis vaccines: adverse reactions. *Pediatrics*. 1995;96:557–66.
47. Johns TL, Hutter GE. New combination vaccines: DTaP-IPV (Kinrix) and DTaP-IPV/Hib (Pentacel). *Ann Pharmacother*. 2010;44:515–23.
48. Lee AW, Jordanov E, Boissard F, Marshall GS. DTaP5-IPV-Hib-HepB, a hexavalent vaccine for infants and toddlers. *Expert Rev Vaccines*. 2017;16:85–92.
49. Obando-Pacheco P, Rivero-Calle I, Raguindin PF, Martinon-Torres F. DTaP5-HBV-IPV-Hib pediatric hexavalent combination vaccine for use in children from 6 weeks through to 4 years of age. *Expert Rev Vaccines*. 2019;18:1115–26.
50. Plotkin SA, Liese J, Madhi SA, Ortiz E. A DTaP-IPV//PRP approximately T vaccine (Pentaxim): a review of 16 years' clinical experience. *Expert Rev Vaccines*. 2011;10:981–1005.
51. Syed YY. DTaP-IPV-HepB-Hib vaccine (Hexyon((R)): an updated review of its use in primary and booster vaccination. *Paediatr Drugs*. 2019;21:397–408.
52. Report from the SAGE Working Group on Pertussis vaccines. *Vaccines*. 26–27 August 2014 meeting, Geneva Switzerland; 2014. https://www.who.int/immunization/sage/meetings/2015/april/1.Pertussis_report_final.pdf
53. Abu-Raya B, Maertens K, Edwards KM, Omer SB, Englund JA, Flanagan KL, et al. Global perspectives on immunization during pregnancy and priorities for future research and development: an international consensus statement. *Front Immunol*. 2020;11:1282.
54. Maertens K, Cabore RN, Huygen K, Hens N, Van Damme P, Leuridan E. Pertussis vaccination during pregnancy in Belgium: results of a prospective controlled cohort study. *Vaccine*. 2016;34:142–50.
55. Eberhardt CS, Blanchard-Rohner G, Lemaitre B, Boukrid M, Combescurie C, Othenin-Girard V, et al. Maternal immunization earlier in pregnancy maximizes antibody transfer and expected infant seropositivity against pertussis. *Clin Infect Dis*. 2016;62:829–36.
56. Abu-Raya B, Giles ML, Kollmann TR, Sadarangani M. The effect of timing of tetanus-diphtheria-acellular pertussis vaccine administration in pregnancy on the avidity of pertussis antibodies. *Front Immunol*. 2019;10:2423.
57. Donegan K, King B, Bryan P. Safety of pertussis vaccination in pregnant women in UK: observational study. *BMJ*. 2014; 349:g4219.
58. Vygen-Bonnet S, Hellenbrand W, Garbe E, von Kries R, Bogdan C, Heininger U, et al. Safety and effectiveness of acellular pertussis vaccination during pregnancy: a systematic review. *BMC Infect Dis*. 2020;20:136.
59. Kharbanda EO, Vazquez-Benitez G, Lipkind HS, Klein NP, Cheetham TC, Naleway AL, et al. Maternal Tdap vaccination: coverage and acute safety outcomes in the vaccine safety datalink, 2007–2013. *Vaccine*. 2016;34:968–73.
60. Jones C, Pollock L, Barnett SM, Battersby A, Kampmann B. The relationship between concentration of specific antibody at birth and subsequent response to primary immunization. *Vaccine*. 2014;32:996–1002.
61. Hardy-Fairbanks AJ, Pan SJ, Decker MD, Johnson DR, Greenberg DP, Kirkland KB, et al. Immune responses in infants whose mothers received Tdap vaccine during pregnancy. *Pediatr Infect Dis J*. 2013;32:1257–60.
62. Ladhani SN, Andrews NJ, Southern J, Jones CE, Amirthalingam G, Waight PA, et al. Antibody responses after primary immunization in infants born to women receiving a pertussis-containing vaccine during pregnancy: single arm observational study with a historical comparator. *Clin Infect Dis*. 2015;61:1637–44.
63. Maertens K, Hoang TT, Nguyen TD, Cabore RN, Duong TH, Huygen K, et al. The effect of maternal pertussis immunization on infant vaccine responses to a booster pertussis-containing vaccine in Vietnam. *Clin Infect Dis*. 2016;63:S197–204.
64. Maertens K, Burbidge P, Van Damme P, Goldblatt D, Leuridan E. Pneumococcal immune response in infants whose mothers received tetanus diphtheria and acellular pertussis vaccination during pregnancy. *Pediatr Infect Dis J*. 2017;36:1186–92.
65. Baxter R, Bartlett J, Fireman B, Lewis E, Klein NP. Effectiveness of vaccination during pregnancy to prevent infant pertussis. *Pediatrics*. 2017;139:e20164091.
66. Amirthalingam G, Andrews N, Campbell H, Ribeiro S, Kara E, Donegan K, et al. Effectiveness of maternal pertussis vaccination in England: an observational study. *Lancet*. 2014;384:1521–8.
67. Grupo de Trabajo tos ferina 2015 de la Ponencia de Programa y Registro de Vacunaciones. Adenda al programa de vacunación frente a tos ferina en España: vacunación en el embarazo. Comisión de Salud Pública del Consejo Interterritorial del Sistema Nacional de Salud. Ministerio de Sanidad. Servicios Sociales e Igualdad. <https://www.mscbs.gob.es/profesionales/saludPublica/prevPromocion/vacunaciones/comoTrabajamos/docs/Adenda.TosFerinaEmbarazo.pdf>.
68. CAV-AEP. Calendario de Vacunaciones de la Asociación Española de Pediatría. Razones y bases de las recomendaciones 2020. [Internet]. Madrid: AEP. 2020. <https://vacunas.aep.org/sites/vacunas.aep.org/files/calvacae2020-razones-y-bases-v.1.3.pdf>.
69. Consejo Interterritorial del Sistema Nacional de Salud. Ministerio de Sanidad. Calendario común de vacunación a lo largo de toda la vida. Calendario recomendado año 2020. <https://www.mscbs.gob.es/profesionales/saludPublica/prevPromocion/vacunaciones/calendario-y-coberturas/docs/CalendarioVacunacion.Todalavida.pdf>.
70. Ministerio de Sanidad. Coberturas de vacunación. Datos estadísticos, 2018. <https://www.mscbs.gob.es/profesionales/saludPublica/prevPromocion/vacunaciones/calendario-y-coberturas/coberturas/home.htm>.
71. World Health Organization. Pertussis vaccines: WHO position paper – September 2015. *Wkly Epidemiol Rec*. 2015;90:433–58.

72. Comité Asesor de Vacunas (CAV-AEP). Tosferina. Manual de vacunas en línea de la AEP [Internet]. Madrid: AEP; ene/2020. <http://vacunasaep.org/documentos/manual/cap-39> [accessed 2.11.20].
73. European Centre for Disease Prevention and Control. Vaccine schedules in all countries of the European Union. <https://vaccine-schedule.ecdc.europa.eu/> [accessed 18.03.21].
74. Foxwell AR, McIntyre P, Quinn H, Roper K, Clements MS. Severe pertussis in infants: estimated impact of first vaccine dose at 6 versus 8 weeks in Australia. *Pediatr Infect Dis J*. 2011;30:161–3.
75. Wood N, Nolan T, Marshall H, Richmond P, Gibbs E, Perrett K, et al. Immunogenicity and safety of monovalent acellular pertussis vaccine at birth: a randomized clinical trial. *JAMA Pediatr*. 2018;172:1045–52.
76. Chiappini E, Petrolini C, Sandini E, Licari A, Pugni L, Mosca FA, et al. Update on vaccination of preterm infants: a systematic review about safety and efficacy/effectiveness Proposal for a position statement by Italian Society of Pediatric Allergy and Immunology jointly with the Italian Society of Neonatology. *Expert Rev Vaccines*. 2019;18:523–45.
77. Grupo de trabajo vacunación en prematuros de la Ponencia de Programa y Registro de Vacunaciones. Comisión de Salud Pública del Consejo Interterritorial del Sistema Nacional de Salud. Ministerio de Sanidad, Consumo y Bienestar Social. Vacunación en prematuros, noviembre 2019. <https://www.mscbs.gob.es/profesionales/saludPublica/prevPromocion/vacunaciones/enfermedades/docs/Vacunacion.Prematuros.pdf>.
78. Comité Asesor de Vacunas (CAV-AEP). Fichas técnicas de las vacunas frente a la tosferina. AEP [Internet]; 2020. <https://vacunasaep.org/profesionales/fichas-tecnicas-vacunas/resultados?diseases=155> [accessed 14.11.20].
79. Chard AN, Gacic-Dobo M, Diallo MS, Sodha SV, Wallace AS. Routine vaccination coverage – worldwide, 2019. *MMWR Morb Mortal Wkly Rep*. 2020;69:1706–10.
80. Chen Z, He Q. Immune persistence after pertussis vaccination. *Hum Vaccin Immunother*. 2017;13:744–56.
81. McGirr A, Fisman DN. Duration of pertussis immunity after DTaP immunization: a meta-analysis. *Pediatrics*. 2015;135:331–43.
82. Schwartz KL, Kwong JC, Deeks SL, Campitelli MA, Jamieson FB, Marchand-Austin A, et al. Effectiveness of pertussis vaccination and duration of immunity. *CMAJ*. 2016;188:E399–406.
83. Pillsbury A, Quinn HE, McIntyre PB. Australian vaccine preventable disease epidemiological review series: pertussis, 2006–2012. *Commun Dis Intell Q Rep*. 2014;38:E179–94.
84. Klein NP, Bartlett J, Rowhani-Rahbar A, Fireman B, Baxter R. Waning protection after fifth dose of acellular pertussis vaccine in children. *N Engl J Med*. 2012;367:1012–9.
85. Domenech de Celles M, Rohani P, King AA. Duration of immunity and effectiveness of diphtheria-tetanus-acellular pertussis vaccines in children. *JAMA Pediatr*. 2019;173:588–94.
86. World Health Organization. Diphtheria vaccine: WHO position paper – August 2017. *Releve epidemiologique hebdomadaire*. 2017;92:417–35.
87. European Center for Disease Prevention and Control. Scientific panel on childhood immunisation schedule: Diphtheria-tetanus-pertussis (DTP) vaccination. [Internet]; 2009. https://www.ecdc.europa.eu/sites/default/files/media/en/publications/Publications/0911.GUL.Scientific_Panel.on.Childhood.Immunisation.DTP.pdf.
88. Xu J, Liu S, Liu Q, Rong R, Tang W, Wang Q, et al. The effectiveness and safety of pertussis booster vaccination for adolescents and adults: a systematic review and meta-analysis. *Medicine (Baltimore)*. 2019;98:e15281.
89. Fernandes EG, Rodrigues CCM, Sartori AMC, De Soarez PC, Novaes HMD. Economic evaluation of adolescents and adults' pertussis vaccination: a systematic review of current strategies. *Hum Vaccin Immunother*. 2019;15:14–27.
90. Baxter R, Bartlett J, Rowhani-Rahbar A, Fireman B, Klein NP. Effectiveness of pertussis vaccines for adolescents and adults: case-control study. *BMJ*. 2013;347:f4249.
91. Álvarez García FJ, Cilleruelo Ortega MJ, Álvarez Aldeán J, Garcés-Sánchez M, García Sánchez N, Garrote Llanos E, et al. Immunisation schedule of the Spanish Association of Paediatrics: 2020 recommendations. *Anales de pediatria (Barcelona, Spain: 2003)*. 2020;92, 52e1–52e10.
92. Centers for Disease, Control and Prevention. Immunization schedules. <https://www.cdc.gov/vaccines/schedules/index.html>.
93. Grupo de trabajo de la Ponencia de Programa y Registro de Vacunaciones. Comisión de Salud Pública del Consejo Interterritorial del Sistema Nacional de Salud. Ministerio de Sanidad SSe. Vacunación en trabajadores sanitarios. 2017. https://www.mscbs.gob.es/profesionales/saludPublica/prevPromocion/vacunaciones/programasDeVacunacion/docs/Vacunacion_sanitarios.pdf.
94. Comité Asesor de Vacunas (CAV-AEP). Vacunaciones del personal sanitario. Manual de vacunas en línea de la AEP (Internet). Madrid: AEP; 2020. <http://vacunasaep.org/documentos/manual/cap-19>.
95. Havers FP, Moro PL, Hunter P, Hariri S, Bernstein H. Use of tetanus toxoid reduced diphtheria toxoid, and acellular pertussis vaccines: updated recommendations of the advisory committee on immunization practices – United States, 2019. *MMWR Morb Mortal Wkly Rep*. 2020;69:77–83.
96. Grupo de trabajo vacunación en población adulta y grupos de riesgo de la Ponencia de Programa y Registro de Vacunaciones. Vacunación en población adulta. Comisión de Salud Pública del Consejo Interterritorial del Sistema Nacional de Salud. Ministerio de Sanidad, Consumo y Bienestar Social. Published date September 2018. https://www.mscbs.gob.es/profesionales/saludPublica/prevPromocion/vacunaciones/programasDeVacunacion/docs/Vacunacion.poblacion_adulta.pdf.
97. Healy CM, Rensch MA, Wootton SH, Castagnini LA. Evaluation of the impact of a pertussis cocooning program on infant pertussis infection. *Pediatr Infect Dis J*. 2015;34:22–6.
98. Dabrera G, Amirthalingam G, Andrews N, Campbell H, Ribeiro S, Kara E, et al. A case-control study to estimate the effectiveness of maternal pertussis vaccination in protecting newborn infants in England and Wales, 2012–2013. *Clin Infect Dis*. 2015;60:333–7.
99. Winter K, Nickell S, Powell M, Harriman K. Effectiveness of prenatal versus postpartum tetanus diphtheria, and acellular pertussis vaccination in preventing infant pertussis. *Clin Infect Dis*. 2017;64:3–8.
100. Bellido-Blasco J, Guiral-Rodrigo S, Miguez-Santiyan A, Salazar-Cifre A, Gonzalez-Moran F. A case-control study to assess the effectiveness of pertussis vaccination during pregnancy on newborns, Valencian community Spain, 1 March 2015 to 29 February 2016. *Euro Surveill*. 2017;22:30545.
101. Saul N, Wang K, Bag S, Baldwin H, Alexander K, Chandra M, et al. Effectiveness of maternal pertussis vaccination in preventing infection and disease in infants: The NSW Public Health Network case-control study. *Vaccine*. 2018;36:1887–92.
102. Fernandes EG, Sato APS, Vaz-de-Lima LRA, Rodrigues M, Leite D, de Brito CA, et al. The effectiveness of maternal pertussis vaccination in protecting newborn infants in Brazil: a case-control study. *Vaccine*. 2019;37:5481–4.
103. Sancho-Uriarte P, Rodriguez SSJ, Sancristobal IG, Agirre NM. Effectiveness of dTpa vaccination during pregnancy in preventing whooping cough in infants under 3 months of age. Bizkaia, Basque Country, Spain. *Heliyon*. 2019;5:e01207.
104. Romanin V, Acosta AM, Juarez MDV, Briere E, Sanchez SM, Cordoba BL, et al. Maternal vaccination in argentina: tetanus diphtheria, and acellular pertussis vaccine effectiveness during pregnancy in preventing pertussis in infants <2 months of age. *Clin Infect Dis*. 2020;70:380–7.