



Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



Editorial

Monitoring of antimicrobial consumption in paediatrics

Monitorización del consumo de antimicrobianos en pediatría



Infectious diseases are the main reason for medical consultation and hospital admission in paediatrics and the first cause of admission to paediatric intensive care units. According to the Antibiotic Resistance and Prescribing in European Children (ARPEC) study, antibiotics are the most prescribed drugs in paediatrics, both within hospitals and primary care.¹ Therefore, the implementation of the Antimicrobial Stewardship Program (ASP) in paediatrics must be a priority for healthcare institutions, especially if we consider that it has been shown, also in paediatrics, a significant impact on the reduction of antibiotic use, and a decrease in healthcare costs and antimicrobial resistance.² Thus, several scientific societies and health organisations have focused their efforts on recognizing paediatrics as a key population group that should be prioritised for the development of the ASPs in order to minimise the inappropriate use of antimicrobials.^{3–11}

Monitoring the amount of antimicrobial use is one of the key strategies in ASP since it allows to control and evaluate the impact of the actions implemented, as well as establishing benchmarking strategies between health centres or between countries or regions.¹² Currently, several units of measurement have been described for monitoring antimicrobials consumption, with the Defined Daily Dose (DDD) being the most widely used.¹³ DDDs have been defined by the World Health Organization (WHO) in collaboration with the Centre for Drug Statistics Methodology in its Anatomical Therapeutic Chemical (WHO)/ATC and correspond to the assumed average maintenance dose per day for a drug used for its primary indication in adults (https://www.whocc.no/atc_ddd_index/). Although there is a broad consensus on the use of DDDs for the definition of quantitative consumption indicators of antimicrobials, they are limited in their application, especially in situations in which a reduced dose from the daily dose administered is used or if there are different dosing recommendations between countries and administration routes (oral, parenteral).

It is especially important to consider that these limitations for the use of DDD are found in the monitoring of consumption in paediatrics, since, by definition, DDD is not an adequate unit of measurement: it is a unit of measurement expressed in grams that do not reflect the Recommended Daily Doses (RDD) or the Prescribed

Daily Dose (PDD). In addition, the great variability of dosing in paediatrics, due to the individualisation of the dose according to the patient's weight and age, means that DDDs do not correctly reflect the real consumption of antimicrobials. Thus, the definition specified that DDDs established by the WHO must not be used for the paediatric or neonatal population.¹⁴ Given this limitation, alternative units of measurement have been suggested. A group of experts established a proposal of 19 possible indicators for monitoring the use of antimicrobials suitable for the paediatric population at the hospital level and clearly stated that the main unit of measurement for antimicrobial consumption in paediatrics should be the Days of Therapy or Days of Treatment (DOT).^{13,15}

However, the calculation of DOT in many organisations indeed presents certain difficulties in obtaining computation, especially due to the lack of the implementation of computerised electronic prescribing or e-prescribing. Faced with this reality, many centres have been forced to suggest initiatives to calculate antimicrobial consumption based on data generated by the consumption in physical units of antimicrobials, complementary to DDD. And it is in this direction that Montecatini et al.^{16,17} published a proposed definition of DDD adapted to paediatrics and neonates (PeDDD). To date, this unit of measurement is an expert proposal, but it has not been sufficiently validated by the international scientific community and, consequently, the authors suggest that it could only be applied in scenarios where other validated units of measurement are impossible to be calculated.

It is in this context that Castagnola et al.¹⁸ conducted a study, published in this issue of *Enfermedades Infecciosas y Microbiología Clínica*, in a tertiary paediatric hospital comparing the results obtained in the monitoring of antimicrobial consumption calculated on the basis DDDs vs. PeDDDs. For this purpose, they estimated the PeDDDD and DDD calculated from the quantity in physical units of each drug consumed. After this analysis, they found that the PeDDDs values were higher than those of the DDDs, but the most remarkable aspect of their results was that the curves between both calculations were parallel and that they obtained identical AUC. Therefore, the calculation for the consumption of antimicrobials using either unit of measurement as an indicator for the exposure to antimicrobials presented no appreciable differences. These results are particularly relevant as they confirm that the monitoring of antimicrobial consumption from PeDDDs should be performed separately for the adult population but in

DOIs of original articles: <https://doi.org/10.1016/j.eimc.2022.06.014>,
<https://doi.org/10.1016/j.eimc.2022.08.008>

<https://doi.org/10.1016/j.eimc.2023.06.007>

0213-005X/© 2023 Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. Published by Elsevier España, S.L.U. All rights reserved.

scenarios where calculating other more appropriate indicators for the paediatric population proves impossible (e.g., DOT), the evaluation of PeDDDs and DDDs would show antimicrobials exposure similarly.

While it is true that, despite attempts to define and use quantitative indicators for antimicrobial use based on units consumed for the paediatric population, in certain circumstances, at the hospital level there is an increasing number of hospitals that have developed tools for obtaining DOTs using relatively automated methods. But in primary care setting it is still more difficult to monitor the consumption with indicators that do not come from the calculation of units dispensed. For this reason, in the primary care setting, the calculation is usually made based on the DDDs per 1000 inhabitants per day (DHD) based on data from the units of containers collected in pharmacies. These indicators can be used for the evaluation of ASPs in antimicrobial consumption or for evaluating the impact of the different social situations that occur, such as the COVID-19 pandemic. At the European level from 2019 to 2020, a substantial decrease in antibiotic consumption in primary care has been observed during the COVID-19 pandemic and in the post-pandemic months.¹⁹ This reduction has been much more evident in the paediatric population in our healthcare environment, as was observed in the study by Perez-Solís et al.,²⁰ also included in this issue of *Enfermedades Infecciosas y Microbiología Clínica*, in which they evaluated the evolution of paediatric antibiotic consumption in primary care in the community of Asturias. In this study, they use data about antibacterial agents for systemic use dispensed for official prescriptions to children under 14 years in primary care. Antibiotic consumption was expressed as defined daily dose (DDD) per 1000 inhabitants per day (DID). The most relevant results were a lower antibiotic use during COVID-19 pandemic, an important decrease in the use of amoxicillin/clavulanic acid and an increased use of amoxicillin and third-generation cephalosporins.

The current challenge is to establish ASP actions to be able to maintain these low levels of antimicrobial consumption and to monitor the impact of other converging situations such as shortages of paediatric pharmaceutical products, the introduction of rapid diagnostic tests in consultations to rule out a bacterial infection in respiratory tract infections or changes in epidemiological profiles in the seasonality of viral infections. It will be necessary to continue developing and implementing technological tools to automate the collection of consumption indicators that allow calculations to be made from units of measurement that more accurately express the real consumption of antimicrobials in the paediatric population in both primary care and hospital care. These indicators will provide ASP teams with relevant and accurate information to design new strategies to improve the use of antimicrobials in the paediatric and neonatal population.

References

- Versporten A, Bielicki J, Drapier N, Sharland M, Goossens H, Calle GM, et al. The worldwide antibiotic resistance and prescribing in European children (ARPEC) pointprevalence survey: developing hospital-quality indicators of antibiotic prescribing for children. *J Antimicrob Chemother.* 2016;71:1106–17.
- Donà D, Barbieri E, Daverio M, Lundin R, Giaquinto C, Zaoutis T, et al. Implementation and impact of pediatric antimicrobial stewardship programs: a systematic scoping review. *Antimicrob Resist Infect Control.* 2020;9:3.
- Policy statement on antimicrobial stewardship by the Society for Healthcare Epidemiology of America (SHEA), the Infectious Diseases Society of America (IDSA), and the Pediatric Infectious Diseases Society (PIDS). *Infect Control Hosp Epidemiol.* 2012;33:322–7.
- Barlam TF, Cosgrove SE, Abbo LM, Macdougall C, Schuetz AN, Septimus EJ, et al. Implementing an antibiotic stewardship program: guidelines by the Infectious Diseases Society of America and the Society for Healthcare Epidemiology of America. *Clin Infect Dis.* 2016;62:e51–77.
- Probst V, Islamovic F, Mirza A. Antimicrobial stewardship program in pediatric medicine. *Pediatr Investig.* 2021;5:229–38.
- Tersigni C, Venturini E, Montagnani C, Chiappini E, de Martino M, Galli L. Antimicrobial stewardship in children: more shadows than lights? *Expert Rev Anti Infect Ther.* 2019;17:871–6.
- Brett A, Bielicki J, Newland JG, Rodrigues F, Schaad UB, Sharland M. Neonatal and pediatric antimicrobial stewardship programs in Europe—defining the research agenda. *Pediatr Infect Dis J.* 2013;32:e456–65.
- Godbout EJ, Pakyz AL, Markley JD, Noda AJ, Stevens MP. Pediatric antimicrobial stewardship: state of the art. *Curr Infect Dis Rep.* 2018;20:39.
- Principi N, Esposito S. Antimicrobial stewardship in paediatrics. *BMC Infect Dis.* 2016;16:1–8.
- Newland JG, Hersh AL. Purpose and design of antimicrobial stewardship programs in pediatrics. *Pediatr Infect Dis J.* 2010;29:862–3.
- McPherson C, Lee BR, Terrill C, Hersh AL, Gerber JS, Kronman MP, et al. Characteristics of pediatric antimicrobial stewardship programs: current status of the sharing antimicrobial reports for pediatric stewardship (SHARPS) collaborative. *Antibiotics.* 2018;7:1–6.
- Centers for Disease Control and Prevention (CDC). Core elements of hospital antibiotic stewardship programs; 2016. <http://www.cdc.gov/getsmart/healthcare/implementation/core-elements.html>.
- Benić MS, Milanić R, Monnier AA, Gyssens IC, Adriaenssens N, Versporten A, et al. Metrics for quantifying antibiotic use in the hospital setting: results from a systematic review and international multidisciplinary consensus procedure. *J Antimicrob Chemother.* 2018;73 Suppl. 6:vi50–8.
- WHO Collaborating Centre for Drug Statistics Methodology. ATC classification index with DDDs, 2021. Oslo, Norway; 2020 [Internet]. Available from: <https://www.who.int/tools/atc-ddd-toolkit/about-ddd>
- Baier J, Höpner J, Haase R, Diexer S, Stareprawo S, Mikolajczyk R, et al. Monitoring antibiotic consumption in pediatrics. How close to reality are days of therapy and recommended daily dose methods? *Pediatr Infect Dis J.* 2022;41:e126–32. <http://dx.doi.org/10.1097/INF.0000000000003446>. PMID: 35027511.
- Montecatine-Alonso E, Gil-Navarro MV, Fernández-Llamazares CM, Fernández-Polo A, Soler-Palacín P, Llorente-Gutiérrez J, et al. Antimicrobial defined daily dose adjusted by weight: a proposal for antibiotic consumption measurement in children. *Enferm Infecc Microbiol Clin.* 2019;37:301–6.
- Villanueva-Bueno C, Montecatine-Alonso E, Jiménez-Parrilla F, Fernández-Llamazares CM, Manrique-Rodríguez S, Zamora-Flores E, et al. Antimicrobial defined daily dose in neonatal population. *Enferm Infecc Microbiol Clin.* 2021;25.S0213-005X(21)00191-9.
- Castagnola E, Lorenzi I, Barabino P, Pistorio A. Antibiotic defined daily dose in pediatrics. A single center study to proof the principle that a specific pediatric definition could be not needed. *Enferm Infecc Microbiol Clin (Engl Ed).* 2023. <http://dx.doi.org/10.1016/j.eimc.2022.08.008>. S2529-993X(23)00027-8.
- Selke Krulichová I, Selke GW, Bennie M, Hajiebrahimi M, Nyberg F, Fürst J, et al. Comparison of drug prescribing before and during the COVID-19 pandemic: a cross-national European study. *Pharmacoepidemiol Drug Saf.* 2022;31:1046–55. <http://dx.doi.org/10.1002/pds.5509>.
- Pérez Solís D, Gómez de Ona C, Nicieza García ML, et al., Consumo de antibióticos en pediatría de atención primaria antes y durante la pandemia de COVID-19, *Enferm Infecc Microbiol Clin*, <https://doi.org/10.1016/j.eimc.2022.06.014>.

Aurora Fernandez-Polo

Servei de Farmàcia, Hospital Infantil, Vall d'Hebron Barcelona
Hospital Campus, Barcelona, Spain

E-mail address: aurora.fernandez@vallhebron.cat