



Enfermedades Infecciosas y Microbiología Clínica

www.elsevier.es/eimc



Original article

The correlation between serum 25-hydroxyvitamin D3 and oxidative balance score in patients with tuberculosis infection



Yan Yong, Li-hong Zhou, Sheng-ya Yang, Xiao-qin Ran, Qing-shan Cai*

Zhejiang Tuberculosis Diagnosis and Treatment Center, Zhejiang Chinese and Western Medicine Integrated Hospital, Hangzhou, Zhejiang, China

ARTICLE INFO

Article history:

Received 11 June 2024

Accepted 16 September 2024

Available online 26 November 2024

Keywords:

Tuberculosis infection

Serum 25-hydroxyvitamin D3

Oxidative balance score

Cross-sectional study

ABSTRACT

Background: Previous study had reported the imbalance of pro- and anti-oxidants in tuberculosis (TB), which was involved in the TB progression. This study aimed to identify the key oxidative stress indicator associated with oxidative balance in patients with TB infection.

Methods: Overall, 339 participants with TB were included in this study. The oxidative balance score (OBS) was composed of scores of dietary and lifestyle factors. Trend regression analysis, generalized additive model, polynomial regression, and threshold effect analysis, were used to examine the relationship between the serum 25-hydroxyvitamin D3 25(OH)D3 and OBS. Moreover, the correlation between serum 25(OH)D3 and OBS components was explored via Pearson analysis, and mediation analysis was used to explore the function of OBS components.

Results: Linear regression analysis showed that among 6 oxidative stress indicators, only serum 25(OH)D3 was independently related to the OBS. A positive association between them was then found (P for trend < 0.05) in a non-linear relationship (all $P < 0.05$). The 25(OH)D3 concentration of 51.9 nmol/L was identified as the key turning point by threshold effect analysis, and their association was only found when it < 51.9 nmol/L. Further, we revealed that 25(OH)D3 mainly correlated with 2 OBS components including BMI and riboflavin. Moreover, BMI and riboflavin were found to mediate the association of serum 25(OH)D3 with OBS by mediation analysis (all $P < 0.05$).

Conclusions: In summary, our study revealed relationship between the serum 25(OH)D3 and OBS in patients with TB, which provided a piece of evidence that vitamin D can assist the treatment of TB.

© 2024 The Author(s). Published by Elsevier España, S.L.U. on behalf of Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Correlación entre la 25-hidroxivitamina D3 sérica y la puntuación del equilibrio oxidativo en pacientes con infección tuberculosa

RESUMEN

Palabras clave:

Infección tuberculosa

25-Hidroxivitamina D3 sérica

Puntuación del equilibrio oxidativo

Estudio transversal

Antecedentes: Estudios previos han informado sobre el desequilibrio entre prooxidantes y antioxidantes en la tuberculosis (TB), implicado en la progresión de la enfermedad. El objetivo de este estudio fue identificar un indicador clave de estrés oxidativo asociado al equilibrio oxidativo en pacientes con TB.

Métodos: Un total de 339 participantes con TB fueron incluidos en este estudio. La puntuación del equilibrio oxidativo (OBS) se calculó a partir de factores dietéticos y de estilo de vida. Se utilizaron análisis de regresión de tendencias, modelos aditivos generalizados, regresión polinómica y análisis de efecto umbral para examinar la relación entre la 25-hidroxivitamina D3 [25(OH)D3] sérica y la OBS. Además, se exploró la correlación entre la 25(OH)D3 sérica y los componentes de la OBS mediante análisis de Pearson, y se aplicó un análisis de mediación para investigar la función de los componentes de la OBS.

Resultados: El análisis de regresión lineal mostró que, entre los 6 indicadores de estrés oxidativo, solo la 25(OH)D3 sérica estaba relacionada de manera independiente con la OBS. Se observó una asociación positiva entre ambos (p para la tendencia $< 0,05$) en una relación no lineal (todas las $p < 0,05$). Una concentración de 25(OH)D3 de 51,9 nmol/L se identificó como un punto de inflexión clave mediante análisis

* Corresponding author.

E-mail address: 13588878702@163.com (Q.-s. Cai).

de efecto umbral, encontrándose la asociación únicamente cuando la concentración era $<51,9$ nmol/L. Además, se reveló que la 25(OH)D3 se correlacionaba principalmente con 2 componentes de la OBS: el IMC y la riboflavina. Asimismo, el análisis de mediación mostró que el IMC y la riboflavina mediaban la asociación entre la 25(OH)D3 sérica y la OBS (todas las $p < 0,05$).

Conclusiones: En resumen, nuestro estudio reveló la relación entre la 25(OH)D3 sérica y la OBS en pacientes con TB, proporcionando evidencia de que la vitamina D podría ayudar en el tratamiento de la tuberculosis.

© 2024 Los Autores. Publicado por Elsevier España, S.L.U. en nombre de Sociedad Española de Enfermedades Infecciosas y Microbiología Clínica. Este es un artículo Open Access bajo la CC BY-NC-ND licencia (<http://creativecommons.org/licencias/by-nc-nd/4.0/>).

Introduction

Tuberculosis (TB) is an airborne infectious disease caused by microorganisms belonging to the mycobacterium TB complex. While predominantly affecting the lungs, mycobacterium TB has the potential to cause disease in various anatomical sites.¹ TB, the primary cause of mortality globally among adults due to an infectious disease, has been recognized as a critical public health crisis on a global scale for the past quarter century. In 2017, the World Health Organization (WHO) reported that approximately 10 million individuals contracted *Mycobacterium tuberculosis* (*M. tuberculosis*), with 8.7 million (87%) of cases occurring in 30 high-burden countries. Despite this high number of cases, only 6.4 million individuals were officially diagnosed and reported. It is estimated that 1.3 million people succumb to TB annually.² In 2015, the WHO released its End TB Strategy, outlining ambitious objectives to decrease global TB incidence by 90% and mortality rates by 95% by the year 2035.³ While public health interventions have resulted in the preservation of numerous lives, advancements in the containment of TB have been limited. The emergence of drug-resistant strains of TB poses a significant threat, with projections indicating that they may become the most lethal pathogens worldwide, accounting for a significant portion of fatalities attributed to antimicrobial resistance.^{4,5} Research has shown that HIV infection, diabetes, history of hospitalization, poor diet quality, and illiteracy^{6,7} were related factors for TB. However, these factors affect the outcome of TB infection by affecting the body's oxidative stress process.⁸ It follows that the involvement of oxidative stress may be critical for the infection of TB and its development.

Oxidative stress is a barrier to the development of many diseases, and excessive generation of harmful reactive oxygen and nitrogen compounds can cause its imbalance.⁹ Abnormal oxidative stress can damage cell membranes, DNA, oxidation of proteins, cell apoptosis process, and cause other adverse effects, thereby impacting the occurrence and development of various diseases.¹⁰ Oxidative stress status has a dual effect on the *M. tuberculosis*. For example, *M. tuberculosis* responds to reactive oxygen species (ROS) in a concentration-dependent manner. Exposure to low concentrations of ROS has no effect on bacterial cell viability but induces the expression of ROS-sensitive response genes, while high concentrations exposure of ROS have been found to be lethal to *M. tuberculosis* cells.¹¹ Overall, when the level of ROS produced by host cells is low, *M. tuberculosis* triggers the release of DNA damage response genes, which initiate DNA repair mechanisms within the bacteria to counteract ROS-induced damage.¹¹ However, this counteraction mechanism of *M. tuberculosis* does not against high levels of ROS.¹² Therefore, appropriate oxidative stress status is critical for the inhibition of TB development.

However, single factors cannot reflect the level of oxidative stress in the body adequately. Therefore, researchers have proposed the concept of an oxidative balance score (OBS).¹³ The concept of "OBS" refers to a metric used to assess the overall balance between oxidative stress and antioxidant capacity in the body. It quantifies the equilibrium between reactive oxygen

species (ROS) production and the body's ability to neutralize them through antioxidant mechanisms. This score integrates various factors related to oxidative stress, such as levels of ROS, antioxidant enzymes, and oxidative damage markers, into a single numerical value, providing a comprehensive measure of the oxidative status of an individual.^{14,15} Most studies have shown that OBS is related to the occurrence and development of diseases. For instance, studies have shown that OBS is negatively correlated with sleep disorders and positively correlated with sleep duration.¹⁶ The results of Da-Hye Son et al. suggested that a healthy diet and lifestyle that increases the OBS may help prevent chronic kidney disease in adults.¹⁷ Higher OBS can reduce the risk of diabetes and heart disease.^{18,19} In TB patients, an imbalance of pro- and anti-oxidants has been observed.²⁰ The clinical value of OBS in various diseases has been gradually reported. However, no study reported the role of oxidative stress in the OBS in TB patients.

To the best of our knowledge, the relationship between oxidative stress and OBS in TB has not been unclear. In this study, we aimed to explore their association in patients with TB infection in U.S. adults based on the National Health and Nutrition Examination Survey (NHANES) database. In addition, we also revealed the key oxidative stress indicators and OBS components that affected the oxidative balance.

Materials and methods

Study population

NHANES is a series of nationwide population-based surveys conducted in the United States, collecting information on health and nutrition, which is released to the public every two years. Using a complex stratified sampling design, NHANES selects a representative sample of the non-institutionalized civilian population of the United States, including Mexican Americans, African Americans, and individuals of other races. The NHANES study protocol has been approved by the Institutional Review Board of the National Center for Health Statistics in the United States, and written informed consent has been obtained from all registered participants. The dataset used in this study is NHANES 2011–2012 as only this cycle has the record of TB. The components of the survey interviews used in this study include demographic characteristics, dietary data, and laboratory examination data.

The TB infection inclusion criteria were as follows: age >18 years old, had complete data of QuantiFERON-TB Gold In-Tube test (QFT-GIT). Individuals with indeterminate QFT-GIT results were excluded. A total of 9756 samples were obtained from the NHANES 2011–2012 cycle, and 7111 samples had the QFT-GIT data. After removing participants with age <18 years old, 5257 samples were retained. Then, the 51 pregnant women and 902 participants who missed the OBS data were excluded. A total of 4290 participants were further used to diagnose TB according to QFT-GIT. QFT-GIT was performed and interpreted according to the guidelines provided by the CDC for using IGRA. The three criteria were used to define positive QFT-GIT: first, Nil value must be ≤ 8.0 IU gamma

Table 1
Analysis of baseline characteristics and oxidative stress indicators related to OBS.

Variables	Total	Low-OBS	High-OBS	χ^2/Z	P-value
Age	57.000 [44.000,66.000]	60.000 [47.000,69.000]	54.000 [40.000,62.000]	3.352	<0.001
PIR	1.740 [0.930,3.620]	1.630 [0.890,2.820]	1.850 [1.020,4.340]	−2.402	0.016
Albumin	4.200 [4.000,4.500]	4.200 [4.000,4.400]	4.300 [4.100,4.500]	−3.055	0.002
GGT	20.000 [14.000,30.000]	22.000 [15.000,33.000]	18.000 [13.000,28.000]	3.019	0.003
Bilirubin	0.700 [0.500,0.800]	0.700 [0.500,0.900]	0.600 [0.500,0.800]	0.013	0.990
Uric acid	5.400 [4.600,6.400]	5.500 [4.800,6.500]	5.300 [4.400,6.200]	1.739	0.082
25(OH)D2	1.450 [1.450,1.450]	1.450 [1.450,1.450]	1.450 [1.450,1.450]	−0.105	0.886
25(OH)D3	55.390 [39.660,71.720]	52.690 [36.230,69.880]	58.280 [45.580,72.310]	−2.360	0.018
OBS lifestyle score	4.000 [3.000,5.000]	4.000 [3.000,5.000]	5.000 [4.000,6.000]	−5.344	<0.001
OBS dietary score	17.000 [12.000,22.000]	12.000 [7.000,15.000]	23.000 [20.000,25.000]	−16.696	<0.001
Sex, n (%)					
Male	232 (58.145)	127 (60.476)	105 (55.556)	0.990	0.320
Female	167 (41.855)	83 (39.524)	84 (44.444)		
Married status, n (%)					
Never marital	24 (11.707)	12 (12.245)	12 (11.215)	4.521	0.104
Marital/living with partner	44 (21.463)	27 (27.551)	17 (15.888)		
Widowed/divorced/separated	137 (66.829)	59 (60.204)	78 (72.897)		
Educational level, n (%)					
Less than high school	145 (36.709)	94 (44.976)	51 (27.419)	15.472	<0.001
High school or equivalent	81 (20.506)	43 (20.574)	38 (20.430)		
High school above	169 (42.785)	72 (34.450)	97 (52.151)		
Obesity, n (%)					
No	262 (66.329)	129 (61.429)	133 (71.892)	4.821	0.028
Yes	133 (33.671)	81 (38.571)	52 (28.108)		
Diabetes, n (%)					
No	314 (80.720)	160 (78.818)	154 (82.796)	0.987	0.320
Yes	75 (19.280)	43 (21.182)	32 (17.204)		
Asthma, n (%)					
No	359 (90.201)	194 (92.823)	165 (87.302)	3.423	0.064
Yes	39 (9.799)	15 (7.177)	24 (12.698)		
Hypertension, n (%)					
No	234 (58.794)	114 (54.545)	120 (63.492)	3.279	0.070
Yes	164 (41.206)	95 (45.455)	69 (36.508)		
Smoking, n (%)					
Never	205 (51.899)	98 (46.890)	107 (57.527)	8.795	0.012
Former	108 (27.342)	56 (26.794)	52 (27.957)		
Now	82 (20.759)	55 (26.316)	27 (14.516)		

Abbreviations: OBS: oxidative balance score, PIR: poverty income ratio, GGT: gamma-glutamyl transpeptidase, serum 25(OH)D2: 25-hydroxyvitamin D2, serum 25(OH)D3: serum 25-hydroxyvitamin D3, serum 25(OH)D: serum 25-hydroxyvitamin D.

interferon (IF)/ml, second, the TB antigen value minus Nil value must be ≥ 0.35 IU gamma interferon (IF)/ml, third, TB antigen value minus Nil value must be $\geq 25\%$ of the Nil value.²¹ Finally, 399 patients with positive QFT-GIT results were enrolled in our analysis.

Variables collection

We collected demographic characteristics including gender, age (years), marital status, education level, poverty status, smoking status, alcohol consumption, underlying diseases, body mass index (BMI, kg/m²), and oxidative stress indicators including Albumin (g/dL), Bilirubin (mg/dL), Uric acid (mg/dL), serum 25-hydroxyvitamin D2 (25(OH)D2, nmol/L), 25(OH)D3 (nmol/L), and gamma-glutamyl transpeptidase (GGT, U/L). OBS was assessed according to related dietary and lifestyle factors, including dietary fiber, carotenoids (α - and β -carotene), riboflavin, niacin, vitamin B6, total folate, vitamin B12, vitamin C, vitamin E, calcium, magnesium, zinc, copper, selenium, total fat intake, iron intake, BMI, physical activity, and contribution to oxidative stress by antioxidants.²² In the evaluation process, higher dietary antioxidant intake and more physical activity induce an increase in OBS, indicating a stronger antioxidant capacity. Conversely, increased total fat, iron, antioxidant intake, and BMI lead to a decrease in OBS, reflecting an increase in oxidative stress lev-

els. The final OBS is obtained by adding up the allocated scores for each component.

Statistical analyses

All statistical analyses used subsample weights following NHANES guidelines in R software version 4.2.2. The Chi-square test was used for the categorical variables and the *T*-test for continuous variables. The relationship between the indicators and OBS was analyzed via linear regression to obtain the indicators significantly related to OBS. We found that only serum 25(OH)D3 was significantly related to OBS. Therefore, the trend regression analysis, generalized additive model (GAM), polynomial regression, and threshold effect analysis, were used to further analyze the relationship between the serum 25(OH)D3 and OBS. In addition, the correlation between serum 25(OH)D3 and OBS components was explored via Pearson analysis, and mediation analysis was used to explore the function of OBS components that were significantly related to serum 25(OH)D3 in the relationship between serum 25(OH)D3 and OBS. All statistical tests conducted in this study were two-tailed, with the threshold for statistical significance set at *P* < 0.05.

Table 2

The association between the above significant indicators and OBS score by univariable and multivariable linear regression analysis in patients with TB.

Variables	Univariable			Multivariable		
	β	95% CI	P-value	β	95% CI	P-value
Age	−0.088	[−0.131, −0.044]	<0.001	−0.084	[−0.137, −0.032]	0.002
PIR	0.866	[0.411, 1.321]	<0.001	0.278	[−0.220, 0.776]	0.274
Albumin	4.660	[2.514, 6.806]	<0.001	2.336	[−0.087, 4.759]	0.060
GGT, (U/L)	−0.022	[−0.043, −0.001]	0.044	−0.018	[−0.040, 0.004]	0.111
25(OH)D3	0.038	[0.009, 0.067]	0.011	0.035	[0.005, 0.066]	0.023
Educational level						
Less than high school	Reference			Reference		
High school or equivalent	1.993	[0.118, 3.867]	0.037	1.010	[−0.919, 2.939]	0.306
Greater than high school	3.921	[2.391, 5.450]	<0.001	2.215	[0.430, 4.000]	0.016
Obesity						
No	Reference			Reference		
Yes	−2.319	[−3.791, −0.846]	0.002	−0.944	[−2.514, 0.626]	0.240
Smoking						
Never	Reference			Reference		
Former	−0.386	[−2.022, 1.249]	0.643	0.952	[−0.767, 2.671]	0.279
Now	−2.954	[−4.751, −1.157]	0.001	−2.364	[−4.220, −0.508]	0.013

Abbreviations: OBS: oxidative balance score, GGT: gamma-glutamyl transpeptidase, CI: confidence interval, serum 25(OH)D3: serum 25-hydroxyvitamin D.

Table 3

Trend regression analysis was between serum 25(OH)D3 and OBS score.

Model	β [95%CI]	P-value	P for trend
Model 1			
Serum 25(OH)D3 as hierarchy variable			<0.001
Q1	Reference		
Q2	2.537 [0.574, 4.500]	0.011	
Q3	2.476 [0.514, 4.439]	0.013	
Q4	2.587 [0.625, 4.550]	0.010	
Model 2			
Serum 25(OH)D3 as hierarchy variable			<0.001
Q1	Reference		
Q2	3.326 [1.343, 5.308]	0.001	
Q3	2.923 [0.972, 4.875]	0.003	
Q4	3.787 [1.758, 5.817]	<0.001	
Model 3			
Serum 25(OH)D3 as hierarchy variable			<0.001
Q1	Reference		
Q2	1.951 [−0.951, 4.854]	0.188	
Q3	1.831 [−0.948, 4.610]	0.197	
Q4	3.505 [0.383, 6.626]	0.028	

Model 1: no adjustment.

Model 2: adjusted for age and PIR.

Model 3: adjusted for age, PIR, educational level, obesity, and smoking.

Abbreviations: OBS: oxidative balance score, CI: confidence interval, serum 25(OH)D3: serum 25-hydroxyvitamin D3.

Results

The baseline characteristics of participants

A total of 399 QFT-GIT positive patients were enrolled in this study. Characteristics of TB patients grouped by the median value of OBS are presented in Table 1. Eight characteristics including age, PIR, albumin, GGT, serum 25(OH)D3, educational level, obesity, and smoking status all showed significant differences between the 2 groups (all $P < 0.05$). These 8 characteristics were enrolled in our further analysis.

Screening the most critical characteristics associated with OBS

We further performed the univariable linear regression to analyze the relationship between the 8 characteristics and OBS. The results showed that all variables had significant relationships with OBS (Table 2). Therefore, those variables were further enrolled in

multivariable linear regression to further screen the independent critical characteristics associated with OBS. The results showed that among 3 oxidative stress indicators, only serum 25(OH)D3 and OBS had a significant independent relationship ($\beta = 0.035$, $P = 0.023$). These results all suggested a significant association between 25(OH)D3 and OBS.

Further, we explored the association changes between serum 25(OH)D3 and OBS by trend regression analysis after setting the serum 25(OH)D3 as a hierarchy variable (Table 3). The results illustrated that with the increase of serum 25(OH)D3 level, the OBS was increased. The same results were observed among 3 models (all P for trend < 0.05). However, we found that the changes of β value were not stable from Q1 to Q4 groups, and its changes appeared a fluctuant phenomenon in Q3 groups among 3 models. Therefore, we further visualized the relationship between serum 25(OH)D3 and OBS by GAM analysis. The results showed that there was a non-linear correlation of serum 25(OH)D3 with OBS (Fig. 1A, $P < 0.01$). Consistent results are obtained by polynomial regression analysis




Fig. 1. The relationship between the serum 25(OH)D3 and OBS score adjusted for age, PIR, educational level, obesity, and smoking. (A) Generalized additive model (GAM). (B) Polynomial regression.

Table 4
Threshold effect analysis of OBS on serum 25(OH)D3.

		Crude		Adjusted	
		β [95%CI]	P-value	β [95%CI]	P-value
Fitting by standard linear model	OBS	0.038 [0.009,0.067]	0.011	0.040 [0.010,0.071]	0.010
Fitting by two-piecewise linear model	In turning point (K)	K = 51.92		K = 51.9	
	<K effect 1	0.107 [0.031,0.183]	0.006	0.120 [0.044,0.195]	0.002
	>K effect 2	0.005 [-0.040,0.049]	0.843	0.001 [-0.045,0.047]	0.977
	Log-likelihood ratio		0.054		0.024

Crude: without adjustment.
Adjusted: adjusting for age, PIR, educational level, smoking, diabetes, hypertension.
Abbreviations: OBS: oxidative balance score, CI: confidence interval, serum 25(OH)D3: serum 25-hydroxyvitamin D3.

(Fig. 1B). Their non-linear correlation may support the fluctuation of β value.

Then, the potential inflection point influencing their association was explored through the threshold effect analysis. The results indicated that serum 25(OH)D3 was significantly related to OBS when it <51.92 (nmol/L) in the crude model, while serum 25(OH)D3 was not related to OBS when it >51.92 (nmol/L) (Table 4). A similar inflection point (51.9) was also found in the adjusted model.

The effect of serum 25(OH)D3 on the composition of OBS

We found that the serum 25(OH)D3 was significantly related to OBS via the above results. Therefore, we further explored the relationship between the serum 25(OH)D3 and 20 components of OBS. The Pearson analysis indicated that the association of serum 25(OH)D3 with BMI and riboflavin was significant ($P < 0.05$, Table 5).

Next, we explored the effect of the BMI and riboflavin on the relationship between the serum 25(OH)D3 and OBS via mediation analysis. The results indicated that BMI and riboflavin had mediating effects on the association of serum 25(OH)D3 with OBS (all $P < 0.05$ for the indirect effect, Table 6).

Discussion

A previous study has reported the imbalance of pro- and anti-oxidants in TB.²⁰ In this study, we further explored the important oxidative stress indicators that correlated with the oxidative balance in TB. Among 6 oxidative stress indexes, only serum 25(OH)D3 showed a significant association with the OBS in patients with TB infection. To delve deeper into this relationship, the threshold effect analysis was performed and its results indicated that

Table 5
Correlation between 25(OH)D3 and OBS components.

OBS components	Coefficient	P-value
Total fat, g	−0.042	0.404
Alpha carotene, mcg	0.055	0.275
Beta carotene, mcg	0.095	0.062
Riboflavin, mg	0.109	0.031
Niacin, mg	0.048	0.344
Vitamin B6, mg	0.043	0.400
Total folate, mcg	−0.017	0.738
Vitamin B12, mcg	0.044	0.384
Vitamin C, mg	−0.021	0.683
Vitamin E ATE, mg	0.024	0.630
Calcium, mg	0.070	0.170
Magnesium, mg	0.060	0.236
Iron, mg	0.005	0.927
Zinc, mg	0.026	0.603
Copper, mg	0.038	0.453
Selenium, mcg	0.023	0.655
Alcohol, g	−0.028	0.586
BMI, kg/m ²	−0.175	<0.001

Abbreviations: OBS: oxidative balance score, GGT: gamma-glutamyl transpeptidase, BMI: body mass index, serum 25(OH)D3: serum 25-hydroxyvitamin D3.

serum 25(OH)D3 was positively related to OBS when serum 25(OH)D3 <51.9 nmol/L in the patients with TB. To our knowledge, this is the first study to explore the association between serum 25(OH)D3 and OBS in patients with TB. Our study initially suggested the importance of 25(OH)D3 on positive regulation of oxidative balance in TB.

Currently, few studies reported their potential association. However, studies have indicated that there was a slight correlation between the levels of serum 25(OH)D3 and the outcomes of TB, and higher concentrations of serum 25(OH)D3 was associated with higher cure rates.²³ According to the positive association

Table 6

Mediation analysis by setting BMI and riboflavin as a mediator.

	Coef	SE	P-value	CI [2.5%]	CI [97.5%]	Sig
<i>Path 1: BMI as mediator</i>						
BMI ~ 25(OH)D3	−0.054	0.016	0.001	−0.085	−0.023	Yes
OBS ~ BMI	−0.134	0.052	0.011	−0.237	−0.031	Yes
Total	0.046	0.015	0.003	0.016	0.076	Yes
Direct	0.040	0.016	0.011	0.009	0.071	Yes
Indirect	0.006	0.003	0.032	0.001	0.015	Yes
<i>Path 1: Riboflavin as mediator</i>						
Riboflavin ~ 25(OH)D3	0.005	0.002	0.016	0.001	0.009	Yes
OBS ~ riboflavin	4.991	0.322	<0.001	4.357	5.625	Yes
Total	0.047	0.015	0.003	0.016	0.077	Yes
Direct	0.023	0.012	0.056	−0.001	0.047	No
Indirect	0.023	0.010	0.024	0.003	0.046	Yes

Abbreviations: OBS: oxidative balance score, BMI: body mass index, CI: confidence interval, serum 25(OH)D3: serum 25-hydroxyvitamin D3.

between serum 25(OH)D3 and OBS in this study, we speculated that the increased cure rates associated with higher serum 25(OH)D3 level may be related to the increase of OBS. OBS is a scoring index based on the calculation of dietary and lifestyle-promoting and antioxidant components,²² and always used to assess systemic oxidative stress status. Higher OBS scores implied exposure to more antioxidants.²⁴ Therefore, higher serum 25(OH)D3 level may refer to a stronger antioxidant environment in TB patients. A study reported that the epigallocatechin gallate (a phenolic antioxidant) in green and black tea can decrease the risk of contracting TB.²⁵ Ginger was an effective antioxidant supplement along with anti-TB therapy, as it possesses strong free radical scavenging property.²⁶ These knowledges may partially explain the favorable effect of serum 25(OH)D3 on the outcome of TB patients by inducing antioxidant environment and scavenging free radical.

Besides the regulation on the oxidative stress environment in TB, serum 25(OH)D3 can also mediate immune responses against *M. tuberculosis*, inducing the production of methyl ethyl ketone and β -defensin 2 (antimicrobial peptides), recruiting monocytes, neutrophils, and T cells to the site of infection.²⁷ Granulysin in T cells is a cationic, amphiphilic, low-molecular-weight antimicrobial peptide that functions to damage cell walls, exhibiting antimicrobial activity against various microbial pathogens, promoting the permeabilization of mycobacteria, and reducing growth in a perforin-dependent manner,^{28,29} thereby inhibiting the progression of TB and consequently improving the OBS of patients with TB.

Further, we identified 2 key OBS components associated with the serum 25(OH)D3, namely BMI and riboflavin. The serum 25(OH)D3 showed negative correlation with pro-oxidant factor BMI but displayed positive correlation with antioxidant factor riboflavin. Existing studies have demonstrated an association between BMI and vitamin D levels.^{30,31} For example, in patients with Ulcerative Colitis, an increase in BMI was associated with a 12% increased likelihood of developing serum 25(OH)D3 deficiency, as indicated by Pallav and ESPEN guidelines.³² According to the research by Vimalaswarns and Pereira-Santos, there was a direct relationship between obesity and 25(OH)D3 deficiency.^{33,34} The specific mechanism needs further study in the future. Our results further supported the establishment of antioxidant environment by serum 25(OH)D3. In addition, we also found that BMI and riboflavin mediated the association of serum 25(OH)D3 with OBS, which highlighted the importance of BMI and riboflavin on the oxidative balance in TB.

This study revealed the relationship between the serum 25(OH)D3 and the OBS in patients with TB for the first time but also had some limitations. First, due to all the components used in the OBS being equally weighted, they may not accurately represent the actual biological contributions. Second, the result was

not an exact causal relationship because of cross-sectional analysis. Third, TB patients diagnosed by other methods may be omitted. Fourth, the population was collected from the NHANES database, which did not represent the global population. Fifth, this study only included important indicators that we collected from the literature, and there may be omissions of indicators that affect serum 25(OH)D3 levels, which may lead to biased results.

Conclusions

In summary, serum 25(OH)D3 was significantly related to OBS in the patients with TB, and their association was especially significant when serum 25(OH)D3 level < 51.9 (nmol/L). Moreover, BMI and riboflavin were the key OBS components correlated with serum 25(OH)D3, and they significantly mediated the association of serum 25(OH)D3 with OBS in patients with TB.

Authors' contributions

Yan Yong contributed to the conception and design. Yan Yong and Li-hong Zhou, Sheng-ya Yang contributed to the collection and assembly of data. Yan Yong, Xiao-qin Ran and Qing-shan Cai analyzed and interpreted the data. All authors wrote and approved the final manuscript.

Funding

Not applicable.

Conflict of interest

The authors declare no conflict of interest.

Data availability statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Acknowledgements

Not applicable.

References

- Pai M, Behr MA, Dowdy D, Dheda K, Divangahi M, Boehme CC, et al. Tuberculosis. *Nat Rev Dis Primers*. 2016;2:16076.
- (GTB) GTP. Global tuberculosis report 2023: global report; 2023.
- Tuberculosis Programme (GTB) TVAC. Global tuberculosis report 2022: global report; 2022. p. 68.
- Furin J, Cox H, Pai M. Tuberculosis. *Lancet*. 2019;393:1642–56.

5. Nathavitharana RR, Friedland JS. A tale of two global emergencies: tuberculosis control efforts can learn from the Ebola outbreak. *Eur Respir J*. 2015;46:293–6.
6. Ji F, Yang Y, Xu L, Cai J, Ni M, Wang Q, et al. Poor diet quality evaluated with the China Healthy Diet Index in Chinese tuberculosis patients. *J Hum Nutr Diet*. 2022;35:1192–201.
7. Gilmour B, Xu Z, Bai L, Alene KA, Clements ACA. Risk factors associated with unsuccessful tuberculosis treatment outcomes in Hunan Province, China. *Trop Med Int Health*. 2022;27:290–9.
8. Shastri MD, Shukla SD, Chong WC, Dua K, Peterson GM, Patel RP, et al. Role of oxidative stress in the pathology and management of human tuberculosis. *Oxid Med Cell Longev*. 2018;2018, 7695364.
9. Sies H. Oxidative stress: a concept in redox biology and medicine. *Redox Biol*. 2015;4:180–3.
10. Hutcheson R, Rocić P. The metabolic syndrome, oxidative stress, environment, and cardiovascular disease: the great exploration. *Exp Diabetes Res*. 2012;2012, 271028.
11. Voskuil MI, Bartek IL, Visconti K, Schoolnik GK. The response of mycobacterium tuberculosis to reactive oxygen and nitrogen species. *Front Microbiol*. 2011;2:105.
12. Wu Y, Gulbins E, Grassme H. Crosstalk between sphingomyelinases and reactive oxygen species in mycobacterial infection. *Antioxid Redox Signal*. 2018;28:935–48.
13. Ellsworth DL, Costantino NS, Blackburn HL, Engler RJ, Kashani M, Vernalis MN. Lifestyle modification interventions differing in intensity and dietary stringency improve insulin resistance through changes in lipoprotein profiles. *Obes Sci Pract*. 2016;2:282–92.
14. Xu Z, Xue Y, Wen H, Chen C. Association of oxidative balance score and lung health from the National Health and Nutrition Examination Survey 2007–2012. *Front Nutr*. 2022;9, 961950.
15. Zhang W, Peng SF, Chen L, Chen HM, Cheng XE, Tang YH. Association between the oxidative balance score and telomere length from the National Health and Nutrition Examination Survey 1999–2002. *Oxid Med Cell Longev*. 2022;2022, 1345071.
16. Lei X, Xu Z, Chen W. Association of oxidative balance score with sleep quality: NHANES 2007–2014. *J Affect Disord*. 2023;339:435–42.
17. Son DH, Lee HS, Seol SY, Lee YJ, Lee JH. Association between the oxidative balance score and incident chronic kidney disease in adults. *Antioxidants (Basel)*. 2023;12:335.
18. Kwon YJ, Park HM, Lee JH. Inverse association between oxidative balance score and incident type 2 diabetes mellitus. *Nutrients*. 2023;15.
19. Ilori TO, Wang X, Huang M, Gutierrez OM, Narayan KM, Goodman M, et al. Oxidative balance score and the risk of end-stage renal disease and cardiovascular disease. *Am J Nephrol*. 2017;45:338–45.
20. Rajopadhye SH, Mukherjee SR, Chowdhary AS, Dandekar SP. Oxidative stress markers in tuberculosis and HIV/TB co-infection. *J Clin Diagn Res*. 2017;11:BC24–8.
21. Hua HB, Wang HJ. Monocyte-to-lymphocyte ratio is significantly associated with positive QuantiFERON-TB Gold-In-Tube and adult survival: an observational study. *Sci Rep*. 2022;12:20406.
22. Goodman M, Bostick RM, Dash C, Flanders WD, Mandel JS. Hypothesis: oxidative stress score as a combined measure of pro-oxidant and antioxidant exposures. *Ann Epidemiol*. 2007;17:394–9.
23. Wang M, Kong W, He B, Li Z, Song H, Shi P, et al. Vitamin D and the promoter methylation of its metabolic pathway genes in association with the risk and prognosis of tuberculosis. *Clin Epigenet*. 2018;10:118.
24. Liu X, Liu X, Wang Y, Zeng B, Zhu B, Dai F. Association between depression and oxidative balance score: National Health and Nutrition Examination Survey (NHANES) 2005–2018. *J Affect Disord*. 2023;337:57–65.
25. Maiolini M, Gause S, Taylor J, Steakin T, Shipp G, Lamichhane P, et al. The war against tuberculosis: a review of natural compounds and their derivatives. *Molecules*. 2020;25.
26. Kulkarni RA, Deshpande AR. Anti-inflammatory and antioxidant effect of ginger in tuberculosis. *J Complement Integr Med*. 2016;13:201–6.
27. Rivas-Santiago B, Serrano CJ, Enciso-Moreno JA. Susceptibility to infectious diseases based on antimicrobial peptide production. *Infect Immun*. 2009;77:4690–5.
28. Ernst WA, Thoma-Uszynski S, Teitelbaum R, Ko C, Hanson DA, Clayberger C, et al. Granulysin, a T cell product, kills bacteria by altering membrane permeability. *J Immunol*. 2000;165:7102–8.
29. Gansert JL, Kiessler V, Engele M, Wittke F, Rollinghoff M, Krensky AM, et al. Human NKT cells express granulysin and exhibit antimycobacterial activity. *J Immunol*. 2003;170:3154–61.
30. Shin JH, Kim CR, Park KH, Hwang JH, Kim SH. Predicting clinical outcomes using phase angle as assessed by bioelectrical impedance analysis in maintenance hemodialysis patients. *Nutrition*. 2017;41:7–13.
31. Tajika M, Matsuura A, Nakamura T, Suzuki T, Sawaki A, Kato T, et al. Risk factors for vitamin D deficiency in patients with Crohn's disease. *J Gastroenterol*. 2004;39:527–33.
32. do Amaral Paes TC, de Oliveira KCC, de Carvalho Padilha P, Peres WAF. Phase angle assessment in critically ill cancer patients: relationship with the nutritional status, prognostic factors and death. *J Crit Care*. 2018;44:430–5.
33. Pereira-Santos M, Costa PR, Assis AM, Santos CA, Santos DB. Obesity and vitamin D deficiency: a systematic review and meta-analysis. *Obes Rev*. 2015;16:341–9.
34. Malden S, Gillespie J, Hughes A, Gibson AM, Farooq A, Martin A, et al. Obesity in young children and its relationship with diagnosis of asthma, vitamin D deficiency, iron deficiency, specific allergies and flat-footedness: a systematic review and meta-analysis. *Obes Rev*. 2021;22, e13129.