



ORIGINAL ARTICLE

Influence of enlarged waist circumference and hypertriglyceridemia in the severity of acute pancreatitis: A retrospective study

Jean Félix Piñerúa-Gonsálvez^{a,*}, María Lourdes Ruiz-Rebollo^a,
Rosanna del Carmen Zambrano-Infantino^b, María Antonella Rizzo-Rodríguez^a,
Luis Fernández-Salazar^a

^a Department of Gastroenterology, Hospital Clínico Universitario de Valladolid, Valladolid, Spain

^b Department of Nuclear Medicine, Hospital Clínico Universitario de Valladolid, Valladolid, Spain

Received 13 November 2022; accepted 30 January 2023

Available online 24 February 2023



KEYWORDS

Acute pancreatitis;
Hypertriglyceridemia;
Waist circumference;
Organ failure;
Local complications

Abstract

Introduction: Acute pancreatitis is a frequent inflammatory gastrointestinal disorder with high mortality rates in severe forms. An early evaluation of its severity is key to identify high-risk patients. This study assessed the influence of waist circumference together with hypertriglyceridemia on the severity of acute pancreatitis.

Methods: A retrospective study was performed, which included patients admitted with acute pancreatitis from March 2014 to March 2021. Patients were classified into four phenotype groups according to their waist circumference and triglyceride levels: normal waist circumference and normal triglycerides; normal waist circumference and elevated triglycerides; enlarged waist circumference and normal triglycerides; and enlarged waist circumference and triglycerides, namely hypertriglyceridemic waist (HTGW) phenotype. Clinical outcomes were compared among the groups.

Results: 407 patients were included. Systemic inflammatory response syndrome (SIRS) and intensive care unit admission were most frequent among patients in the HTGW phenotype group, at 44.9% and 8.2%, respectively. The incidence of local complications was higher in the normal waist circumference with elevated triglycerides group (27%). On multivariable analysis, an enlarged waist circumference was related to an increase of 4% and 2% in the likelihood of developing organ failure and SIRS, respectively. Hypertriglyceridemia was an independent risk factor for both organ failure and local complications.

* Corresponding author.

E-mail address: jeanfelixmedicina@hotmail.com (J.F. Piñerúa-Gonsálvez).

PALABRAS CLAVE

Pancreatitis aguda;
Hipertrigliceridemia;
Circunferencia de
cintura;
Fallo orgánico;
Complicaciones
locales

Conclusions: HTGW phenotype was significant related to developing of SIRS. It seems that an enlarged waist circumference has a greater role than hypertriglyceridemia in the development of SIRS. Obesity and hypertriglyceridemia were both independent risk factors for organ failure. Patients with hypertriglyceridemia were more likely to develop local complications.
© 2023 Elsevier España, S.L.U. All rights reserved.

Influencia del aumento de la circunferencia de la cintura y la hipertrigliceridemia en la gravedad de la pancreatitis aguda: un estudio retrospectivo

Resumen

Introducción: La pancreatitis aguda es una patología frecuente con altas tasas de mortalidad en sus formas graves. Este estudio evaluó la influencia de la circunferencia de la cintura (CC) junto con la hipertrigliceridemia en la gravedad de la pancreatitis aguda.

Métodos: Se realizó un estudio retrospectivo que incluyó pacientes con pancreatitis aguda desde 2014 hasta 2021. Los pacientes se clasificaron en cuatro grupos fenotípicos según su CC y los niveles de triglicéridos: CC normal y triglicéridos normales, CC normal y triglicéridos elevados, CC aumentada y triglicéridos normales, y CC aumentada y triglicéridos elevados, es decir, el fenotipo cintura hipertrigliceridémica (HTGW).

Resultados: Se incluyeron 407 pacientes. El síndrome de respuesta inflamatoria sistémica (SIRS) y la admisión a la unidad de cuidados intensivos fueron más frecuentes entre los pacientes con fenotipo HTGW, en 44,9 y 8,2%, respectivamente. La incidencia de complicaciones locales fue mayor en el grupo de CC normal con triglicéridos elevados (27%). En el análisis multivariable, una CC aumentada se relacionó con un aumento de 4 y 2% en la probabilidad de desarrollar fallo orgánico y SIRS, respectivamente. La hipertrigliceridemia fue un factor de riesgo tanto para el fallo orgánico como para las complicaciones locales.

Conclusiones: El fenotipo HTGW se relacionó con el desarrollo de SIRS. Parece que una CC aumentada tiene un papel más importante que la hipertrigliceridemia en el desarrollo de SIRS. La obesidad y la hipertrigliceridemia fueron factores de riesgo independientes para el fallo orgánico. Los pacientes con hipertrigliceridemia tenían más probabilidades de desarrollar complicaciones locales.

© 2023 Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Acute pancreatitis (AP) is one of the most frequent inflammatory gastrointestinal disorders worldwide. The global pooled incidence is reported to be 34 cases per 100,000 people per year, with North America and the Western Pacific having the highest rates.^{1,2} Most of the patients admitted have a mild disease and a short hospital stay.³ Nevertheless, in severe cases, the mortality rate rises to 30–50%, mainly due to its association with systemic inflammatory response syndrome (SIRS) and multiple organ failure.^{4,5} As a result, an accurate and timely assessment of the AP severity is critical for identifying high-risk patients, which improves outcomes and lowers the mortality rate.

Obesity (defined as a body mass index [BMI] > 30 kg/m²) is a probed independent risk factor for AP, which is associated with a complicated course, especially abdominal obesity (characterized by a large waist circumference [WC]). This can be explained by the fact that visceral adipose tissue leads to a hypersecretion of proinflammatory mediators such as interleukin 6 and tumor necrosis factor, increasing the inflammatory response when pancreatitis is present.^{6,7} In

addition, hypertriglyceridemia (HTG), which is one of the most frequent lipometabolic disorders, is an independently significant risk factor for the development of organ failure in AP.^{8–11}

The concept of hypertriglyceridemic waist (HTGW) phenotype was formulated by Lemieux et al. in 2000, which includes HTG and a large WC.¹² This phenotype has been used to identify the characteristics of excessive visceral adipose tissue. Furthermore, these comorbidities have been associated with metabolic syndrome, cardiovascular disease, diabetes, and hypertension. In recent years, interest in the relationship between HTGW and AP has increased.^{4,5} This study was aimed to assess the influence of HTG and an enlarged WC on the severity of AP.

Materials and methods**Study design**

We performed a retrospective study from a prospective database of patients diagnosed with AP who were admitted

from March 2014 to March 2021 at the Gastroenterology Service of *Hospital Clínico Universitario de Valladolid*, Spain.

The exclusion criteria included age <18 years, recurrent pancreatitis, previous diagnosis of chronic pancreatitis, history of surgery involving the pancreas, history of cancer, pregnancy, and incomplete data in the electronic medical record.

The study was conducted in accordance with the Declaration of Helsinki and was approved by the Ethics Committee of *Hospital Clínico Universitario de Valladolid* under the number PI-22-2850.

Data collection

Data were obtained from electronic medical records at *Hospital Universitario de Valladolid*. Information collected included patient clinical demographic data, etiology, severity, length of stay, local complications, organ failure, systemic complications, SIRS, intensive care unit (ICU) admission, interventional procedure need, and mortality.

Study definitions and classifications

According to the revision of the Atlanta classification, the diagnosis of AP required at least two of the following features: abdominal pain consistent with AP, biochemical evidence of pancreatitis (amylase or lipase at least three times greater than the upper limit of normal), and/or characteristic findings on contrast-enhanced computed tomography (CECT), magnetic resonance imaging (MRI) or transabdominal ultrasonography.¹³

The severity of pancreatitis was classified as mild (absence of organ failure and the absence of local or systemic complications), moderately severe (presence of transient organ failure and/or local or systemic complications within 48 h), and severe (presence of persistent organ failure involving one or more organs and lasting >48 h).^{4,13}

Organ failure is defined as a score ≥ 2 points for one of three organs (respiratory, cardiovascular or renal system) of the modified Marshall scoring.¹⁴ Systemic complication was defined as an exacerbation of a pre-existing co-morbidity such as heart failure, coronary artery disease, chronic lung disease, or chronic kidney disease precipitated by AP.¹³

SIRS was defined as the presence of two or more of the following criteria: temperature >38 or <36 °C, heart rate >90 beats/min, respiratory rate >20 breaths/minute or partial pressure of CO₂ <32 mmHg, leukocyte count $>12,000$ or <4000 /microliters or $>10\%$ immature forms.^{13,15}

Large WC was defined as ≥ 102.0 for males and ≥ 88.0 cm for females.¹⁶ This measurement was made by using a measuring tape at the midpoint between the lowest rib margin and iliac crest after a normal exhale.

HTG was defined as serum triglycerides (TG) ≥ 1.70 mmol/L.^{16,17} As the measurement of TG levels is not routinely performed in the emergency department, this measurement was carried out 24 h after the admission to hospital. The waist triglyceride index (WTI) was calculated by using the formula: $WTI = WC \text{ (cm)} \times TG \text{ (mmol/L)}$.¹⁸

Heavy alcohol drinking was defined as a consumption of >4 standard drinks/day in men or >3 standard drinks/day in women, or >14 standard drinks/week in men and >7 standard drinks/week in females.¹⁹

Definition of phenotypes

According to WC and TG, participants were divided into four phenotype groups: NWNT (normal WC + normal TG); NWET (normal WC + elevated TG); EWNT (enlarged WC + normal TG); HTGW (enlarged WC + elevated TG).^{4,5,17,20}

Statistical analysis

In the case of a normal distribution, continuous variables were expressed as mean $\pm$ standard deviation (SD) while they were expressed as the median and interquartile range in a nonnormal distribution. Categorical variables as frequencies and proportions. For continuous variables, phenotype groups were compared by using one-way analysis of variance (ANOVA) with Bonferroni post hoc correction or Kruskal–Wallis test according to the result normality test. For categorical data, we use the Chi-square or 2-tailed Fisher's exact test. Univariable and multivariable logistic regressions were used to calculate odds ratios (ORs) and 95% confidence intervals (95% CIs) for predictors of organ failure, SIRS and local complications. The following variables were included: age, hypertension, diabetes, heavy alcohol consumption, smoking, BMI, high levels of TG, WC and WTI. Predictors that were determined to be significant by univariable analysis remained in the adjusted model. Statistical analyses were calculated using IBM® SPSS® Statistics 21.0.

Results

Among the 635 patients diagnosed with AP throughout the study period, 407 (64.0%) met the inclusion criteria (Fig. 1). Two hundred seven (56.5%) were female. The mean age was 68.3 ± 17.0 years. Regarding lifestyle habits, 16% were smokers and 27.8% drank alcohol. Among the previous chronic diseases, 53.1% had hypertension and 15.7% had diabetes. The most frequent cause of AP among the different phenotypes was biliary (62.4%). Baseline patient and admission characteristics are summarized in Table 1.

Clinical outcome comparison among phenotype groups is shown in Table 2. Regarding severity of AP, 75.9% presented with mild acute pancreatitis (MAP), 17.9% with moderately severe acute pancreatitis (MSAP), and 6.1% with severe acute pancreatitis (SAP). There was a significant difference in the severity of AP among the phenotype groups. MSAP (32.7%) and SAP (10.2%) were most frequent in the HTGW group. The overall incidence of organ failure was 13.8%, being higher in the HTGW group (34.7%). Forty-eight patients (11.8%) developed local complications, and their incidence was higher in the NWET group. Overall, SIRS developed in 116 (28.5%) patients and was most common among patients in the HTGW group (44.9%). ICU care was required most frequently among patients in the HTGW group (8.2%). There were no significant differences among the groups in terms of systemic complications ($P=0.23$), interventional proce-




Figure 1 Flow diagram of patient selection. AP: acute pancreatitis; CP: chronic pancreatitis; NWNT: normal waist circumference + normal triglycerides; NWET: normal waist circumference + elevated triglycerides; EWNT: enlarged waist circumference + normal triglycerides; HTGW, enlarged waist circumference + triglycerides.

Table 1 Patient baseline and clinical characteristics at admission in four phenotypes groups.

Variables	NWNT (n = 128)	NWET (n = 37)	EWNT (n = 193)	HTGW (n = 49)	P value ^a
Age (years), mean ($\pm$ SD)	68.3 ($\pm$ 18.8)	58.7 ($\pm$ 17.6) ^a	70.8 ($\pm$ 15.9) ^b	65.5 (15.0)	<0.001
Sex, n (%) ^{ab}					<0.001
Male	71 (55.5)	28 (75.7)	61 (31.6)	17 (34.7)	
Female	57 (44.5)	9 (24.3)	132 (68.4)	32 (65.3)	
BMI, mean ($\pm$ SD)	24.4 ($\pm$ 4.1)	25.8 ($\pm$ 3.6)	29.8 ($\pm$ 5.3) ^{ab}	29.9 ($\pm$ 4.2) ^{ab}	<0.001
WC (cm), mean ($\pm$ SD)	84.5 ($\pm$ 10.6)	89.7 ($\pm$ 9.3)	107.9 ($\pm$ 12.3) ^{ab}	108.3 ($\pm$ 10.1) ^{ab}	<0.001
TG (mmol/L), median (IQR)	0.93 (0.7–1.2)	2.22 (1.8–2.7) ^a	1.02 (0.7–1.3) ^b	2.22 (1.8–2.6) ^{ac}	<0.001
WTI (cm mmol/L), mean ($\pm$ SD)	81.3 ($\pm$ 31.9)	465.5 ($\pm$ 1069.1) ^a	113.4 ($\pm$ 41.8) ^b	276.5 ($\pm$ 139.0) ^{ac}	<0.001
Etiologies, n (%)					<0.001
Biliary	69 (53.9)	19 (51.4)	140 (72.5)	26 (53.1)	
Alcohol	13 (10.2)	10 (27)	8 (4.1)	0 (0)	
ERCP	3 (2.3)	1 (2.7)	3 (1.6)	5 (10.2)	
Idiopathic	30 (23.4)	3 (8.1)	35 (18.1)	12 (24.5)	
Others	13 (10.2)	4 (10.8)	7 (3.6)	6 (12.2)	
Ranson score, median (IQR)	1 (1–2)	1 (1–2)	1 (1–3)	1 (1–2)	0.029
BISAP score $\geq$ 3, n (%)	11 (8.6)	2 (5.4)	21 (10.9)	3 (6.1)	0.691
Diabetes mellitus, n (%)	19 (14.8)	4 (10.8)	30 (15.5)	11 (22.4)	0.489
Hypertension, n (%)	56 (43.8)	15 (40.5)	76 (39.4) ^{ab}	21 (42.9)	0.009
Heavy alcohol drinker, n (%)	39 (30.5)	17 (45.9)	46 (23.8) ^b	7 (14.3) ^{ab}	0.006
Smoker, n (%)	23 (18.0)	18 (48.6) ^a	18 (9.3) ^{ab}	6 (12.2) ^b	<0.001

Abbreviations: NWNT, normal waist circumference + normal triglycerides; NWET, normal waist circumference + elevated triglycerides; EWNT, enlarged waist circumference + normal triglycerides; HTGW, enlarged waist circumference + triglycerides; BMI, body mass index; WC, waist circumference; TG, triglycerides; WTI, waist triglyceride index; IQR, interquartile range; BISAP, Bedside Index of Severity in Acute Pancreatitis.

The significance for pairwise comparisons was calculated using the Bonferroni adjustment. ^a $P < 0.05$, compared with NWNT; ^b $P < 0.05$, compared with the NWET; ^c $P < 0.05$, compared with EWNT.

^a Comparison among the four groups. Bold indicates statistical significance.

Table 2 Comparison of clinical outcomes in four phenotypes groups.

Variables	NWNT (n = 128)	NWET (n = 37)	EWNT (n = 193)	HTGW (n = 49)	P value ^a
Severity, n (%)					0.003
MAP	108 (84.4)	25 (67.6)	148 (76.7)	28 (57.1)	
MSAP	12 (9.4)	11 (29.7)	34 (17.6)	16 (32.7)	
SAP	8 (6.3)	1 (2.7)	11 (5.7)	45 (10.2)	
SIRS, n (%)	25 (19.5)	9 (24.3)	60 (31.1)	22 (44.9) ^a	0.006
Organ failure, n (%)	11 (8.6)	2 (5.4)	26 (13.5)	17 (34.7) ^{abc}	<0.001
Local complication, n (%)	7 (5.5)	10 (27) ^a	24 (12.4)	7 (14.3)	0.004
Systemic complications, n (%)	22 (17.3)	9 (24.3)	47 (24.6)	15 (30.6)	0.239
ICU admission, n (%)	0	1 (2.7)	7 (3.6) ^a	4 (8.2) ^a	0.015
Interventional procedure need, n (%)	7 (5.5)	4 (10.8)	18 (9.3)	9 (18.4)	0.069
Length of stay (days), mean (±SD)	7.6 (±11.1)	12.6 (±17.8)	9.2 (±13.9)	13 (±19.5)	0.076
Mortality, n (%)	5 (3.9)	0	5 (2.6)	2 (4.1)	0.684

Abbreviations: NWNT, normal waist circumference + normal triglycerides; NWET, normal waist circumference + elevated triglycerides; EWNT, enlarged waist circumference + normal triglycerides; HTGW, enlarged waist circumference + triglycerides; MAP, mild acute pancreatitis; MSAP, moderately severe acute pancreatitis; SAP, severe acute pancreatitis; SIRS, systemic inflammatory response syndrome; ICU, intensive care unit.

The significance for pairwise comparisons was calculated using the Bonferroni adjustment. ^a $P < 0.05$, compared with NWNT; ^b $P < 0.05$, compared with the NWET; ^c $P < 0.05$, compared with EWNT.

^a Comparison among the four groups. Bold indicates statistical significance.

dures need ($P = 0.07$), length of stay ($P = 0.07$) or mortality ($P = 0.68$).

Univariable regression analysis showed that age, hypertension, diabetes mellitus, BMI, WC and HTG were associated with organ failure, and all of them except diabetes and HTG were related to SIRS. On multivariable logistic regression analysis, WC remained a significant predictor, increasing by 4% and 2% the likelihood of developing organ failure (OR 1.04; 95% CI 1.01–1.06) and SIRS (OR 1.02; 95% CI 1.01–1.04), respectively, for each 1-cm increase in WC. BMI was found to be a risk factor for SIRS (OR 1.06; 95% CI 1.01–1.12). HTG (OR 3.10; 95% CI 1.52–6.30), age (OR 1.04; 95% CI 1.01–1.07) and hypertension (OR 2.31; 95% CI 1.03–5.17) were also predictors of organ failure. The results are summarized in Tables 3 and 4.

With regard to local complications, HTG (OR 2.31; 95% CI 1.20–4.46) as well as alcoholism (OR 2.40; 95% CI 1.28–4.49) remained independent risk factors for local complications on the multivariable logistic regression analysis (Table 5).

Discussion

In our research from a prospective database, it was investigated the effects of WC and HTG on AP severity. The present study demonstrated that the isolated HTG phenotype exhibited the most significant association with local complications. On the other hand, a significantly high incidence of SIRS was linked with both the HTGW and EWNT groups. Nevertheless, this relationship was more significant for the HTGW group. In addition, ICU need was most frequent in the HTGW (8.2%).

Several studies have reported that HTG is related to a high incidence of SAP. This can be explained by the fact that in the setting of AP, high levels of TG are hydrolyzed into free fatty acids by pancreatic lipase, which exceeds the capacity of the regulatory mechanisms. The excess of

free fatty acids causes direct tissue injury (lipotoxicity) and pancreatic capillary damage. In addition, an upregulation of cytokines induced by high levels of TG leads to an intense inflammatory response causing multiple organ failure.^{8,21–23}

The current study found that the incidence of SIRS was higher in the HTGW, which is in line with the study from Ding et al.⁴ However, TG levels had no statistical significance in the multivariable analysis, which suggests that the enlarged WC is what could be related to SIRS in this group of patients instead of high levels of TG. On the other hand, we found that patients with HTG had higher incidences of organ failure and local complications. These findings are consistent with those of Hidalgo et al. and Kiss et al., who reported that high levels of TG were independently associated with pancreatic necrosis and organ failure.^{24,25}

This study has shown that the enlarged WC remained a significant predictor of organ failure and SIRS in the adjusted model. These findings were consistent with two previous studies conducted at Instituto Nacional de Ciencias Médicas y Nutrición “Salvador Zubiran” in Mexico, which demonstrated that abdominal adiposity according to WC is associated with an increased risk of multiple organ failure.^{7,26} WC is the most accurate surrogate marker of visceral adiposity and an easy index to obtain.²⁷ Low levels of adiponectin, which inhibits macrophage production of tumor necrosis factor- α and interleukin-6, have been linked to visceral adiposity. Consequently, the high levels of these cytokines lead to an amplified inflammatory response, increasing the severity of AP.⁶

There are several limitations in the current study that need to be considered. First, there might be potential confounders that were not measured but may influence the present associations. Second, this study was carried out in a single center, and most of the patients came from the same geographical area. Therefore, caution should be taken at the moment of generalizing the present results to other popula-

Table 3 Univariable and multivariable analysis of proposed predictors of organ failure in patients with acute pancreatitis.

Variables	Univariable analysis		Multivariable analysis	
	OR (95% CI)	P value	OR (95% CI)	P value
Age	1.042 (1.020–1.064)	<0.001	1.043 (1.015–1.071)	0.003
Hypertension	4.898 (2.395–10.013)	<0.001	2.319 (1.039–5.173)	0.040
Diabetes mellitus	2.525 (1.312–4.860)	0.006	1.427 (0.695–2.930)	0.333
Heavy alcohol drinker	1.245 (0.642–2.417)	0.517		
Smoker	1.684 (0.690–4.108)	0.252		
BMI	1.065 (1.014–1.118)	0.011	1.034 (0.956–1.116)	0.407
WC	1.037 (1.018–1.058)	<0.001	1.040 (1.017–1.064)	0.001
HTG (≥ 1.70 mmol/L)	2.177 (1.178–4.022)	0.013	3.101 (1.526–6.303)	0.002
WTI	1.000 (0.999–1.001)	0.846		

Bold indicates statistical significance.

Abbreviations: BMI, body mass index; WC, waist circumference; HTG, hypertriglyceridemia; WTI, waist triglyceride index.

Table 4 Univariable and multivariable analysis of proposed predictors of SIRS in patients with acute pancreatitis.

Variables	Univariable analysis		Multivariable analysis	
	OR (95% CI)	P value	OR (95% CI)	P value
Age	1.016 (1.002–1.030)	0.021	1.012 (0.996–1.028)	0.148
Hypertension	1.670 (1.076–2.593)	0.022	1.261 (0.752–2.116)	0.379
Diabetes mellitus	1.637 (0.935–2.865)	0.084		
Heavy alcohol drinker	1.293 (0.784–2.134)	0.314		
Smoker	1.265 (0.687–2.330)	0.450		
BMI	1.026 (1.012–1.042)	<0.001	1.067 (1.010–1.127)	<0.001
WC	1.026 (1.012–1.042)	<0.001	1.026 (1.011–1.042)	0.001
HTG (≥ 1.70 mmol/L)	1.565 (0.944–2.594)	0.082		
WTI	1.001 (1.000–1.003)	0.141		

Bold indicates statistical significance.

Abbreviations: BMI, body mass index; WC, waist circumference; HTG, hypertriglyceridemia; WTI, waist triglyceride index.

Table 5 Univariable and multivariable analysis of proposed predictors of local complications in patients with acute pancreatitis.

Variables	Univariable analysis		Multivariable analysis	
	OR (95% CI)	P value	OR (95% CI)	P value
Age	0.992 (0.975–1.009)	0.359		
Hypertension	1.550 (0.834–2.881)	0.166		
Diabetes mellitus	1.349 (0.548–3.319)	0.515		
Heavy alcohol drinker	2.395 (1.290–4.448)	0.006	2.406 (1.287–4.496)	0.006
Smoker	1.249 (0.573–2.721)	0.576		
BMI	0.992 (0.937–1.051)	0.794		
WC	1.014 (0.994–1.033)	0.173		
HTG (≥ 1.70 mmol/L)	2.305 (1.207–4.402)	0.011	2.317 (1.203–4.463)	0.012
WTI	1.001 (1.000–1.002)	0.116		

Bold indicates statistical significance.

Abbreviations: BMI, body mass index; WC, waist circumference; HTG, hypertriglyceridemia; WTI, waist triglyceride index.

tions. Third, TG levels were measured 24 h after admission since this measurement is not routinely performed at emergency room. This delay in the measurement might lead to bias because most patients with AP are treated by fasting during the first hours of hospitalization, resulting in a fall in TG levels. This fact has been observed in previous studies.^{25,28–30} Due to those limitations, we suggest carrying

out multicenter, prospective studies in order to corroborate the results of the present study.

In summary, our retrospective study found that HTGW and isolated enlarged WC phenotypes were significantly related to the development of SIRS. It seems that an enlarged WC has a greater role than HTG in the development of SIRS in these patients. The need for admission to the ICU was higher in the

HTGW phenotype. Moreover, an enlarged WC and HTG were independent risk factors for organ failure. Patients with HTG were more likely to develop local complications. The previous findings suggest that the measurement of WC and TG levels seems to be helpful in identifying high-risk patients with AP on admission.

Authors' contributions

Involved in the conception, analysis of data and draft of the manuscript: JFIG and RCZI. Involved in the acquisition of data: MLRR and MARR. Revised manuscript critically for important intellectual content: MLRR and LFS. All authors contributed to the article and approved the submitted version.

Funding

None declared.

Conflict of interest

No potential conflict of interest was reported by the authors.

References

- Petrov MS, Yadav D. Global epidemiology and holistic prevention of pancreatitis. *Nat Rev Gastroenterol Hepatol*. 2019;16:175–84.
- Xiao AY, Tan MLY, Wu LM, Asrani VM, Windsor JA, Yadav D, et al. Global incidence and mortality of pancreatic diseases: a systematic review, meta-analysis, and meta-regression of population-based cohort studies. *Lancet Gastroenterol Hepatol*. 2016;1:45–55.
- Forsmark CE, Vege SS, Wilcox CM. Acute pancreatitis. *N Engl J Med*. 2016;375:1972–81.
- Ding Y, Zhang M, Wang L, Yin T, Wang N, Wu J, et al. Association of the hypertriglyceridemic waist phenotype and severity of acute pancreatitis. *Lipids Health Dis*. 2019;18:93.
- Yang X, He J, Ma S, Wang T, Zhu Q, Cao F, et al. The role of comorbid hypertriglyceridemia and abdominal obesity in the severity of acute pancreatitis: a retrospective study. *Lipids Health Dis*. 2021;20:171.
- Natu A, Stevens T, Kang L, Yasinow S, Mansoor E, Lopez R, et al. Visceral adiposity predicts severity of acute pancreatitis. *Pancreas*. 2017;46:776–81.
- Duarte-Rojo A, Sosa-Lozano LA, Saul A, Herrera-Caceres JO, Hernandez-Cardenas C, Vazquez-Lamadrid J, et al. Methods for measuring abdominal obesity in the prediction of severe acute pancreatitis, and their correlation with abdominal fat areas assessed by computed tomography. *Aliment Pharmacol Ther*. 2010;32:244–53.
- Nawaz H, Koutroumpakis E, Easler J, Slivka A, Whitcomb DC, Singh VP, et al. Elevated serum triglycerides are independently associated with persistent organ failure in acute pancreatitis. *Am J Gastroenterol*. 2015;110:1497–503.
- Pascual I, Sanahuja A, Garcia N, Vazquez P, Moreno O, Tosca J, et al. Association of elevated serum triglyceride levels with a more severe course of acute pancreatitis: cohort analysis of 1457 patients. *Pancreatol*. 2019;19:623–9.
- Ruiz-Rebollo ML, Munoz-Moreno MF. Clinical relationship between serum triglycerides and acute pancreatitis. *Pancreatol*. 2020;20:785–6.
- Niknam R, Moradi J, Jahanshahi KA, Mahmoudi L, Eftehadi F. Association between metabolic syndrome and its components with severity of acute pancreatitis. *Diabetes Metab Syndr Obes*. 2020;13:1289–96.
- Lemieux I, Pascot A, Couillard C, Lamarche B, Tchernof A, Almeras N, et al. Hypertriglyceridemic waist: a marker of the atherogenic metabolic triad (hyperinsulinemia; hyper-apolipoprotein B; small, dense LDL) in men? *Circulation*. 2000;102:179–84.
- Banks PA, Bollen TL, Dervenis C, Gooszen HG, Johnson CD, Sarr MG, et al. Classification of acute pancreatitis – 2012: revision of the Atlanta classification and definitions by international consensus. *Gut*. 2013;62:102–11.
- Marshall JC, Cook DJ, Christou NV, Bernard GR, Sprung CL, Sibbald WJ. Multiple organ dysfunction score: a reliable descriptor of a complex clinical outcome. *Crit Care Med*. 1995;23:1638–52.
- Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest*. 1992;101:1644–55.
- Grundy SM, Brewer HB Jr, Cleeman JI, Smith SC Jr, Lenfant C, American Heart A, et al. Definition of metabolic syndrome: report of the National Heart Lung, and Blood Institute/American Heart Association conference on scientific issues related to definition. *Circulation*. 2004;109:433–8.
- Zhou M, Li F, Tang H, Wu S, Meng L, Dong Y, et al. The hypertriglyceridemic waist phenotype is associated with fatty liver and glycometabolic profiles in overweight and obese adults: a cross-sectional study. *Sci Rep*. 2022;12:2410.
- Yu X, Wang D, Xiao W, Shi X, She Q, Sun H, et al. Relationship between fatty pancreas and hypertriglyceridemic waist phenotype: a cross-sectional study. *Sci Rep*. 2020;10:21937.
- National Institute on Alcohol Abuse and Alcoholism. Drinking levels defined. National Institute on Alcohol Abuse and Alcoholism; 2023. Available from: <https://www.niaaa.nih.gov/alcohol-health/overview-alcohol-consumption/moderate-binge-drinking>
- Amini M, Esmailzadeh A, Sadeghi M, Mehvarifar N, Amini M, Zare M. The association of hypertriglyceridemic waist phenotype with type 2 diabetes mellitus among individuals with first relative history of diabetes. *J Res Med Sci*. 2011;16:156–64.
- Yu S, Wu D, Jin K, Yin L, Fu Y, Liu D, et al. Low serum ionized calcium, elevated high-sensitivity c-reactive protein neutrophil-lymphocyte ratio, and body mass index (BMI) are risk factors for severe acute pancreatitis in patients with hypertriglyceridemia pancreatitis. *Med Sci Monit*. 2019;25:6097–103.
- Kimura W, Mossner J. Role of hypertriglyceridemia in the pathogenesis of experimental acute pancreatitis in rats. *Int J Pancreatol*. 1996;20:177–84.
- Chen L, Huang Y, Yu H, Pan K, Zhang Z, Man Y, et al. The association of parameters of body composition and laboratory markers with the severity of hypertriglyceridemia-induced pancreatitis. *Lipids Health Dis*. 2021;20:9.
- Hidalgo NJ, Pando E, Alberti P, Vidal L, Mata R, Fernandez N, et al. Elevated serum triglyceride levels in acute pancreatitis: a parameter to be measured and considered early. *World J Surg*. 2022;46:1758–67.
- Kiss L, Fur G, Matrai P, Hegyi P, Ivany E, Cazacu IM, et al. The effect of serum triglyceride concentration on the outcome of acute pancreatitis: systematic review and meta-analysis. *Sci Rep*. 2018;8:14096.
- Mery CM, Rubio V, Duarte-Rojo A, Suazo-Barahona J, Pelaez-Luna M, Milke P, et al. Android fat distribution as predictor of severity in acute pancreatitis. *Pancreatol*. 2002;2:543–9.
- Borrueal S, Molto JF, Alpanes M, Fernandez-Duran E, Alvarez-Blasco F, Luque-Ramirez M, et al. Surrogate markers of visceral

- adiposity in young adults: waist circumference and body mass index are more accurate than waist hip ratio, model of adipose distribution and visceral adiposity index. *PLoS One*. 2014;9:e114112.
28. Carr RA, Rejowski BJ, Cote GA, Pitt HA, Zyromski NJ. Systematic review of hypertriglyceridemia-induced acute pancreatitis: a more virulent etiology? *Pancreatology*. 2016;16: 469–76.
 29. Lloret Linares C, Pelletier AL, Czernichow S, Vergnaud AC, Bonnefont-Rousselot D, Levy P, et al. Acute pancreatitis in a cohort of 129 patients referred for severe hypertriglyceridemia. *Pancreas*. 2008;37:13–22.
 30. Dominguez-Munoz JE, Malfertheiner P, Ditschuneit HH, Blanco-Chavez J, Uhl W, Buchler M, et al. Hyperlipidemia in acute pancreatitis relationship with etiology, onset, and severity of the disease. *Int J Pancreatol*. 1991;10:261–7.