



ELSEVIER  
DOYMA



## ACTUALIZACIÓN

# Resonancia magnética artrografía de cadera con énfasis en el atrapamiento femoroacetabular

A. Kassarjian<sup>a,\*</sup>, L. Cerezal<sup>b</sup> y E. Llopis<sup>c</sup>

<sup>a</sup>División de Radiología Musculoesquelética, Massachusetts General Hospital,  
Profesor adjunto de Radiología, Harvard Medical School, Boston, Estados Unidos,  
Consultor de Radiología, Corades, S.L., Madrid, España

<sup>b</sup>Servicio de Radiología. Clínica Mompía, Santander, España

<sup>c</sup>Servicio de Radiología. Hospital de la Ribera, Alzira, Valencia, España

Recibido el 21 de diciembre de 2007; aceptado el 26 de marzo de 2008

MR arthrography of the hip with emphasis on femoroacetabular impingement

### Abstract

Hip pain is a common complaint in patients of all ages. Recent advances in imaging and treatment are changing the approach to the evaluation and management of hip pain. Abnormal femoral and acetabular morphology and lesions of the acetabular labrum and cartilage are increasingly recognized as crucial in the development of degenerative changes. In addition, femoroacetabular impingement is increasingly recognized as an etiologic factor in hip pain. This article discusses techniques for MR arthrography of the hip, normal anatomy at hip MR arthrography, common intra-articular pathologies in patients with hip pain, and imaging findings of femoroacetabular impingement.

© 2007 SERAM. Published by Elsevier España, S.L. All rights reserved.























