




REVISTA COLOMBIANA DE PSIQUIATRÍA

www.elsevier.es/rcp



Original Article

Prevalence of Anxiety Symptoms Among Health Care Workers in Colombia During the COVID-19 Pandemic



Joaquín O. Ruiz-Villa^{a,*}, Sergio A. Ochoa-Orozco^b, Andrés Gaviria-Mendoza^{c,d}, Juan D. Castrillón-Spitia^c, Luis F. Echeverri-Cataño^e, Jorge E. Machado-Alba^f

^a Anesthesiologist in Chief, Clínica los Rosales, Pereira, Colombia

^b Psychiatrist in Chief, Grupo de Investigación Hospital Mental Universitario de Risaralda (HOMERIS), Pereira, Colombia

^c Epidemiologist, Grupo de Investigación en Biomedicina, Fundación Universitaria Visión de las Américas, Pereira, Colombia

^d Epidemiologist, Grupo de Investigación en Farmacoepidemiología y Farmacovigilancia, Universidad Tecnológica de Pereira-Audifarma S.A. Pereira, Colombia

^e General Practitioner, Grupo de Investigación en Farmacoepidemiología y Farmacovigilancia, Universidad Tecnológica de Pereira-Audifarma S.A. Pereira, Colombia

^f MD, PhD Pharmacology, Grupo de Investigación en Farmacoepidemiología y Farmacovigilancia, Universidad Tecnológica de Pereira-Audifarma S.A. Pereira, Colombia

ARTICLE INFO

Article history:

Received 8 November 2021

Accepted 16 January 2023

Available online 31 January 2023

Keywords:

Coronavirus

Healthcare workers

Anxiety

Mental health

Colombia

ABSTRACT

Objective: This study aimed to determine the prevalence of anxiety symptoms in a Colombian HCW sample during the COVID-19 pandemic.

Methods: A cross-sectional study was carried out by means of an online survey (May–June 2020). Respondents were HCWs in Colombia reached by a nonprobability sample. Zung's self-rating anxiety scale allowed the estimation of prevalence and classification of anxiety symptoms.

Results: A total of 568 HCWs answered the questionnaire, 66.0% were women, the mean age was 38.6 ± 11.4 years. 28.9% presented with anxiety symptoms, of whom 9.2% were moderate–severe. Characteristics such as living with relatives at higher risk of mortality from COVID-19 infection (OR: 1.90; 95% CI: 1.308–2.762), female sex (OR: 2.16; 95% CI: 1.422–3.277), and personal history of psychiatric illness (OR: 3.41; 95% CI: 2.08–5.57) were associated with higher levels of anxiety. Access to sufficient personal protective equipment (OR: 0.45; 95% CI: 0.318–0.903) and age >40 years (OR: 0.53; 95% CI: 0.358–0.789) were associated with lower anxiety levels.

Conclusions: Anxious symptoms are common in the population of HCWs faced with patient care during the COVID-19 pandemic. Different strategies are required to intervene with subgroups at risk of developing higher levels of anxiety during the pandemic.

© 2023 Asociación Colombiana de Psiquiatría. Published by Elsevier España, S.L.U. All rights reserved.

* Corresponding author.

E-mail address: jorruizvilla@gmail.com (J.O. Ruiz-Villa).

<https://doi.org/10.1016/j.rcp.2023.01.004>

0034-7450/© 2023 Asociación Colombiana de Psiquiatría. Published by Elsevier España, S.L.U. All rights reserved.

Prevalencia de síntomas de ansiedad en personal de salud en Colombia durante la pandemia por COVID-19

R E S U M E N

Palabras clave:

Coronavirus
Personal de salud
Ansiedad
Salud mental
Colombia

Objetivo: Determinar la prevalencia de síntomas de ansiedad en una muestra de personal de salud (PDS) colombianos durante la pandemia por COVID-19.

Métodos: Se llevó a cabo un estudio de corte transversal mediante una encuesta en línea (mayo a junio de 2020). Los encuestados fueron PDS en Colombia reclutados mediante una muestra no probabilística. La escala de autoevaluación de ansiedad de Zung permitió la estimación de la prevalencia y la clasificación de los síntomas de ansiedad.

Resultados: Un total de 568 PDS respondieron el cuestionario, el 66,0% fueron mujeres y la edad promedio fue de $38,6 \pm 11,4$ años. El 28,9% presentaron síntomas de ansiedad, de los cuales el 9,2% fueron moderados-severos. Características como vivir con familiares con mayor riesgo de mortalidad por infección por COVID-19 (OR: 1,90; IC 95%: 1,308-2,762), sexo femenino (OR: 2,16; IC 95%: 1,422-3,277) y la presencia de historia personal de patología psiquiátrica (OR: 3,41; IC 95%: 2,08-5,57) se asociaron con mayores niveles de ansiedad. El acceso a elementos de protección personal suficientes (OR: 0,45; IC 95%: 0,318-0,903) y las edades >40 años (OR: 0,53; IC 95%: 0,358-0,789) se correlacionaron con menores niveles de ansiedad.

Conclusiones: Los síntomas ansiosos son comunes en la población de PDS enfrentados al cuidado de pacientes durante la pandemia por COVID-19. Se requieren diferentes estrategias para intervenir los subgrupos en riesgo de desarrollar mayores niveles de ansiedad durante la pandemia.

© 2023 Asociación Colombiana de Psiquiatría. Publicado por Elsevier España, S.L.U.
Todos los derechos reservados.

Introduction

Novel coronavirus infection described in December 2019 at Wuhan (COVID-19) started as a pneumonia of an unknown virus circumscribed to a region of China and then grew to a pandemic.¹⁻³ Some studies carried out during previous epidemics have associated quarantine duration with higher incidence of negative emotions, affecting people's mental health and leading to an aftermath featuring characteristics such as post-traumatic stress disorder (PTSD), avoidant behavior, and anger.⁴⁻⁶ Similar studies have also showed that isolation times longer than 10 days had a positive correlation with more severe symptoms of PTSD than shorter isolation times.^{4,7,8}

During the initial stages of the COVID-19 pandemic, Chinese health care workers (HCWs) suffered an increase in the prevalence of anxiety symptoms. The symptoms (such as anxiety, sleep disorders, or exacerbations of previously diagnosed psychiatric pathologies) appear to be stronger among those HCWs in charge of the care of patients with a diagnosis or suspicion of COVID-19 infection.⁹⁻¹⁵

There are differences in perceived stress, anxiety levels, depression symptoms, and changes in mental health status among HCWs in several countries, with different results having been found among professions, specialties, gender, and other variables.¹⁶⁻¹⁸ Some governmental and scientific initiatives have exposed the need to raise awareness about

this phenomenon and the possible management that it may entail.^{12,13,15}

COVID-19 is probably the worst pandemic faced by Latin America in the last century.¹⁹ HCWs in this region have had several challenges related to this pandemic: lack of evidence-based knowledge about the new virus and changing protocols; constantly crowded consultation and emergency services facilities, which interfere with social distancing; the need for personal protective equipment (PPE); and rejection and attacks from the general population caused by the fear of being infective agents.²⁰

Colombia first faced a national quarantine in March 2020, after the first COVID-19 case and related death were reported in the country. Previous pandemics have shown that quarantine produces in HCWs fear of contagion, concern about the safety of coworkers and peers in the health care field, loneliness, and demanding expectations, which could result in anger, anxiety, and stress related to the uncertainty of the event.²⁰

In Colombia, previous studies have reported changes in perceived stress,²¹ sleep disorders,²² and psychological distress²³ associated with COVID-19 infection in the general population. Currently there is a lack of evidence, limited to few studies assessing HCW population about perceived anxiety levels in Colombia during the COVID-19 pandemic. The aim of this study was to determine the prevalence of anxiety symptoms in a sample of Colombian HCWs during the early stage of the COVID-19 pandemic and quarantine.

Methods

Study design and population

We carried out a cross-sectional study through a self-administered online survey among a nonprobability sample of HCWs in Colombia. The survey remained online from May 20 to June 20, 2020.

Sample size calculation and sampling strategy

We calculated sample size with an expected frequency of 50%, an acceptable error rate of 5%, and a design effect of one for a population survey, with a resulting sample size of 384 participants.

Sampling was divided into two strategies. The first one was a virtual snowball method, by posting the survey in social media groups and contacts (WhatsApp and Telegram), asking respondents to answer and share the link with coworkers and colleagues. The other method was to send a predefined e-mail to renowned providers in charge of the direction or leadership of scientific communities (e.g., the Colombian Society of Anesthesiology and the Colombian Society of Plastic Surgery), asking for permission to share the posted link with their associate databases.

Estimation of prevalence and classification of anxiety

To assess anxiety among the survey respondents, we used the Zung self-rating anxiety scale (SAS) designed by William W.K. Zung in 1971²⁴ and posteriorly validated in Colombia by De La Ossa et al.²⁵

The Zung SAS deploys a questionnaire of 20 items, each of which tests psychological or somatic anxiety-related symptoms using a 4-point ordinal scale based on frequency of presentation. Those items are presented in a positive or negative form (e.g., “I feel calm and can sit still easily” vs “I feel weak and get tired easily”). Respondents are asked to base their answers on experiences lived in a given period (usually the previous week, and with the present study, the previous month). The scores from the 20 items are then added together.²⁴

According to the norms for the Zung SAS, raw scores (ranging from 20 to 80) are converted to an index score (ranging from 25 to 100).²⁴ We used the original 1971 cutoff value of 50 points in the index score, considering it the value that produces fewer misdiagnoses compared with the cutoff value recommended later by Zung in 1980.²⁴ The results with a positive value in the Zung index score (above 50 points) were reclassified according to the severity degree in three subgroups – mild, moderate, and severe anxiety – with cutoff points of 50–59, 60–69, and ≥ 70 , correspondingly.²⁶

Data analysis

Statistical analysis of the results was done using the software SPSS v26.0. Continuous variables are shown with measures of central tendency and their dispersion. Categorical data

are shown in frequencies and proportions. The chi-squared test was used to identify possible relations between categorical variables, as well as testing for the presence of anxiety symptoms (yes/no). Significance was established at $p < 0.05$.

Ethical considerations

This study was endorsed by the bioethics committee of the Universidad Tecnológica de Pereira and was categorized as research “with minimal risk,” according to Resolution 8430 of 1993 of the Ministry of Health, which guides research in Colombia. Likewise, the study adhered to generally accepted ethical principles, confidentiality, and the Declaration of Helsinki. The patients accepted the informed consent to participate in the study as well as to receive the results of their SAS.

Results

A total of 568 participants from 47 municipalities (23 of 32 departments) answered the survey during the study observation window. Sixty-six percent ($n = 375$) were women. The mean age of the study population was 38.6 ± 11.4 years (median 36 years; interquartile range [IQR]: 29–46 years). Socioeconomic level (SEL) was classified in six categories from 1 to 6, where 1 and 2 correspond to low SEL (6.6%, $n = 37$), 3 and 4 to middle SEL (60.6%, $n = 344$), and 5 and 6 to high (32.5%, $n = 185$) (see [Table 1](#)).

Most of the HCWs from the survey population used their private vehicles to go to work ($n = 434$; 76.4%). The median of reported working time per week was 48 h (IQR: 36–50 h). Sixty-six percent ($n = 375$) of the health professionals were involved in the care of suspected or confirmed COVID-19 infection patients, and 82.9% stated they had received personal protective equipment (PPE) ($n = 471$) (see [Table 2](#)), a fact associated with a lower probability of having anxiety ($p < 0.001$; OR: 0.45; 95% CI 0.318–0.903) or, in cases of presenting symptoms of anxiety, having a lower severity ($p < 0.018$; OR: 0.54; 95% CI 0.318–0.903).

Of the respondents, 159 (28.0%) did not report financial dependents, and 87 (21.5%) reported four or more financial dependents. When asked about people with whom they live, 186 (33.0%) reported living with people older than 60 years, 205 (36.0%) lived with people younger than 18 years, and 196 (35.0%) lived with people who presented a known risk factor of increased mortality from COVID-19 infection. For these categories of living arrangements, the participants isolated themselves from their relatives in 29.0%, 20.5%, and 27.6% of cases, respectively (see [Table 2](#)).

Zung anxiety scale and anxiety prevalence

The median raw Zung anxiety scale score was 34.0 points (IQR: 30–41 points), and the median index score was 42.5 (IQR: 37.5–51.3 points). In [Table 2](#) we describe the index score according to anxiety severity classification. Of the total population, 28.9% (95% CI 25.1–32.6%) presented with anxiety symptoms ($n = 164$), and from those, 112 participants (19.7%)

Table 1 – Sociodemographic information of the respondents. Total sample and subgroups according to anxiety level.

Variable	Total (n = 568)		Anxiety severity classification*							
	n	%	No anxiety (n = 404)		Mild (n = 112)		Moderate (n = 42)		Severe (n = 10)	
			n	%	n	%	n	%	n	%
Sex										
Male	193	34.0	156	80.8	29	15.0	6	3.1	2	1.0
Female	375	66.0	248	66.1	83	22.1	36	9.6	8	2.1
Age – mean (SD)	38.6 (11.4)		39.8 (11.9)		36.2 (9.9)		35.2 (8.8)		32.8 (9.2)	
≤40 years	355	62.5	236	66.5	81	22.8	31	8.7	7	2.0
≥40 years	213	37.5	168	78.9	31	14.6	11	5.2	3	1.4
Socioeconomic level										
Low	37	6.5	23	62.2	6	16.2	3	8.1	5	13.5
Middle	344	60.8	235	68.3	78	22.7	28	8.1	3	0.9
High	185	32.7	144	77.8	28	15.1	11	5.9	2	1.1
Educational level										
Technician	51	9.0	33	64.7	12	23.5	5	9.8	1	2.0
Undergraduate degree	263	46.3	165	62.7	68	25.9	24	9.1	6	2.3
Specialty degree	220	38.7	179	81.4	29	13.2	10	4.5	2	0.9
Unknown	34	6.0	27	79.4	3	8.8	3	8.8	1	2.9
City										
Bogotá	148	26.1	109	73.6	29	19.6	9	6.1	1	0.7
Pereira	111	19.6	93	83.8	14	12.6	4	3.6	0	0.0
Unknown	75	13.2	48	64.0	18	24.0	7	9.3	2	2.7
Cali	55	9.7	38	69.1	13	23.6	3	5.5	1	1.8
Medellin	37	6.5	26	70.3	7	18.9	4	10.8	0	0.0
La Virginia	27	4.8	21	77.8	4	14.8	1	3.7	1	3.7
Cartago	15	2.6	10	66.7	4	26.7	0	0.0	1	6.7
Apartadó	12	2.1	5	41.7	5	41.7	2	16.7	0	0.0
Pácora	10	1.8	6	60.0	1	10.0	2	20.0	1	10.0
Barranquilla	10	1.8	7	70.0	1	10.0	1	10.0	1	10.0
Other cities	67	11.8	40	59.7	16	23.9	9	13.4	2	3.0
Occupation										
General practitioner	156	27.5	93	59.6	42	26.9	17	10.9	4	2.6
Anesthesiologist	72	12.7	62	86.1	9	12.5	0	0.0	1	1.4
Nurse	60	10.6	40	66.7	13	21.7	6	10.0	1	1.7
Nurse assistant	49	8.6	31	63.3	12	24.5	5	10.2	1	2.0
Pediatrician	35	6.2	27	77.1	4	11.4	3	8.6	1	2.9
Dentist	27	4.8	15	55.6	10	37.0	1	3.7	1	3.7
Ophtalmologist	20	3.5	16	80.0	4	20.0	0	0.0	0	0.0
Occupational physician	13	2.3	9	69.2	3	23.1	1	7.7	0	0.0
Surgical technologist	10	1.8	9	90.0	1	10.0	0	0.0	0	0.0
Other ^a	126	22.2	102	81.0	14	11.1	9	7.1	1	0.8

^a Other occupations include: epidemiologist, infectologist, gynecologist, family physician, internist, intensivist, orthopedist, bacteriologist, psychiatrist, dermatologist, general surgeon, respiratory therapist, otorhinolaryngologist, neurologist, neonatologist, neurosurgeon, gastroenterologist, radiology technician, pharmacy manager, pharmaceutical chemist, optometrist, nutritionist, rheumatologist, pneumologist, maxillofacial surgeon, gastrointestinal surgeon, cardiovascular surgeon. SD: standard deviation.

* Percentages of anxiety severity classification are shown considering the total of each row, not columns.

had mild, 42 (7.4%) moderate, and 10 (1.8%) severe anxiety. Women had a higher anxiety proportion than men (33.9% vs 19.2%, $p < 0.001$, OR: 2.16; 95% CI: 1.422–3.277). Regarding anxiety severity classification according to sex, women had higher figures compared to men in mild (22.1% vs 15.0%), moderate (9.6% vs 3.1%), and severe (2.1% vs 1.0%) anxiety.

Those with a personal history of psychiatric illness had a greater probability of presenting anxiety symptoms during the COVID-19 pandemic ($p < 0.001$, OR: 3.41; 95% CI: 2.080–5.570).

We also found that anxiety symptoms were more frequent in the younger group (≤ 40 years, 33.5%, $n = 119$) compared with their older counterparts (> 40 years, 21.1%, $n = 45$, $p < 0.005$, OR: 0.53; 95% CI: 0.358–0.789).

Those participants living with relatives with risk factors of increased mortality from COVID-19 infection had a higher proportion of anxiety compared with those who did not live with relatives with risk factors of increased mortality (37.8% vs 24.2%, $p < 0.001$, OR: 1.90; 95% CI: 1.308–2.762). Although anxiety proportions in the groups of HCWs living with younger

Table 2 – Miscellaneous factors explored in the respondents.

Variable	Anxiety severity classification							
	No anxiety (n = 404)		Mild (n = 112)		Moderate (n = 42)		Severe (n = 10)	
	n	%	n	%	n	%	n	%
<i>Impact on income</i>								
Decreased	226	55.9	72	64.3	29	69.0	4	40.0
Was not modified	175	43.3	40	35.7	12	28.6	6	60.0
Raised	3	0.7	0	0.0	1	2.4	0	0.0
<i>COVID patient in last month</i>								
No	125	30.9	22	19.6	6	14.3	3	30.0
Yes	258	63.9	78	69.6	32	76.2	7	70.0
Don't know	21	5.2	12	10.7	4	9.5	0	0.0
<i>Personal protective elements</i>								
No	47	11.6	25	22.3	10	23.8	3	30.0
Yes	346	85.6	86	76.8	32	76.2	7	70.0
Not applicable	11	2.7	1	0.9	0	0.0	0	0.0
<i>Work overload (> 48 h/week)</i>								
No	292	72.5	71	63.4	33	78.6	6	60.0
Yes	111	27.5	41	36.6	9	21.4	4	40.0
<i>Lives with people over 60 years of age</i>								
No	276	68.3	72	64.3	28	66.7	6	60.0
Yes, with personal isolation	36	8.9	12	10.7	4	9.5	2	20.0
Yes, without personal isolation	92	22.8	28	25.0	10	23.8	2	20.0
<i>Lives with children under 18 years of age</i>								
No	263	65.1	71	63.4	23	54.8	6	60.0
Yes, with personal isolation	25	6.2	11	9.8	4	9.5	2	20.0
Yes, without personal isolation	116	28.7	30	26.8	15	35.7	2	20.0
<i>Lives with people with risk factors for COVID-19</i>								
No	282	69.8	62	55.4	23	54.8	5	50.0
Yes, with personal isolation	33	8.2	13	11.6	6	14.3	2	20.0
Yes, without personal isolation	89	22.0	37	33.0	13	31.0	3	30.0
<i>Number of financial dependents (including participant)</i>								
One (1)	110	27.2	37	33.0	9	21.4	3	30.0
Two (2)	107	26.5	27	24.1	12	28.6	2	20.0
Three (3)	100	24.8	30	26.8	14	33.3	1	10.0
Four or more (4+)	87	21.5	18	16.1	7	16.7	4	40.0
<i>Opposition, anger, and disappointment over mandatory isolation</i>								
Never	187	46.3	20	17.9	6	14.3	0	0.0
Sometimes	197	48.8	61	54.5	23	54.8	4	40.0
Many times	18	4.5	30	26.8	10	23.8	3	30.0
Always	2	0.5	1	0.9	3	7.1	3	30.0
<i>Mental and physical stress caused by being separated from loved ones</i>								
Never	113	28.0	11	9.8	5	11.9	1	10.0
Sometimes	209	51.7	30	26.8	9	21.4	1	10.0
Many times	74	18.3	56	50.0	21	50.0	3	30.0
Always	8	2.0	15	13.4	7	16.7	5	50.0
<i>History of psychiatric illness</i>								
No	368	91.1	85	75.9	30	71.4	8	80.0
Yes	36	8.9	27	24.1	12	28.6	2	20.0
Zung anxiety index score median (IQR)		40.0 (7.5)		53.8 (3.8)		62.5 (3.7)		75.0 (7.8)

IQR: interquartile range.

(<18 years) or older (>65 years) relatives were slightly higher than those who did not, there was not a statistically significant difference. Higher SEL was associated with a lower prevalence of anxiety (22.2% vs 32.3%, $p < 0.013$, OR: 0.59; 95% CI: 0.397–0.898).

Discussion

With this analysis we were able to establish the prevalence of anxiety in a sample of Colombian HCWs during the COVID-19 pandemic through an online survey. These findings can be

helpful to HCWs, their employers, and health care authorities to create policies that mitigate risk.

Despite the fact that COVID-19-associated anxiety in HCWs presents a novel situation, some studies, mainly in the Chinese population, have found an anxiety prevalence ranging from 12.5% to 44.6%.^{9,14,27,28} Our results fit within the range of prevalence previously reported. Another study performed in Colombia (Medellín) found anxiety symptoms in 50.1% of the HCW included, on a similar date.²⁹ The difference would be explained by the difference between the instruments used to assess anxiety. They used GAD-7, which includes cognitive and emotional anxiety symptoms characteristic of genal anxiety disorder (potentially more prevalent at that time for HCW), whereas SAS evaluates cognitive, emotional, autonomic, and physical anxiety symptoms in a broader spectrum.²⁴

Many factors would influence the prevalence of anxiety in HCWs, explaining the differences among the studies. Our data were collected between May and June 2020, when Colombia was going through the third month of quarantine. Despite the fact that Colombia was reporting increasing new daily COVID-19 cases, this period represented an early stage of the outbreak. It is necessary to consider that the number of cases and the possibility of exposure to infection would affect the anxiety in HCWs, as has been proposed in other studies.³⁰ Even though this was an early stage of the pandemic, knowledge about the strategies to face it was scarce, and uncertainty was high among the general population, including HCWs. These conditions, even in the absence of previous data about the prevalence of anxiety symptoms in HCWs in Colombia, could possibly be an independent cause of higher levels of anxiety.

As we did, some other studies have found a higher COVID-19-related anxiety prevalence in women working in health services.^{14,15,27} These sex-based differences are expected because the prevalence of anxiety disorders is higher in women potentially because of genetic, neurodevelopmental, environmental, and neurobiological factors.³¹ In this study, an increased anxiety prevalence was observed in HCWs with preexisting psychiatric diagnoses. This finding was also seen in other previous studies in HCWs.²⁷ There is evidence in the general population showing that people with preexisting anxiety or mood-related disorders would exhibit increased COVID-19-related stress.³² There are some limitations in our study with respect to this issue, because we did not ask respondents to specify the mental health previous diagnosis, and this variable was based on self-report rather than clinical diagnostic evaluation.

A study performed on frontline medical staff members found that their sources of anxiety included their own and their family's safety. In our study, having family members with risk factors for increased COVID-19 mortality was associated with higher anxiety, a key factor because 38% of the population lived with people with these conditions. There is an under-recognized problem that would increase HCWs' anxiety: fear, avoidance, and stigma. A study in the Canadian general population found that people had these negative feelings regarding HCWs during the COVID-19 pandemic.³³ This situation would reinforce the belief that HCWs are dangerous for their families.

HCWs who received PPE had lower anxiety prevalence. This finding has been described in other studies performed in

similar populations,³⁴ where researchers found a relationship between lack of PPE and increased COVID-19-related stress in HCWs.³⁵ A worrying finding was the lack of PPE in almost 20% of HCWs in our study, something explored in Latin America, where this population lacks sufficient PPE and feels limited support from human resources and public officials.³⁶

A systematic review assessed the COVID-19 pandemic's impact on mental health in the general population. The review, which included 19 studies in a qualitative synthesis of the literature, found among the included studies a correlation between younger age groups (≤ 40 years), lower educational level, poor self-rated health, high loneliness, female sex, divorced/widowed status, quarantine status, worry about being infected, history of mental health issues/medical problems, presence of chronic illness, living in urban areas, and specific physical symptoms.³⁷

In China, the government and members of society typically compliment medical personnel for their dedication to fight COVID-19, trying to generate in HCWs feelings of being "honored and proud".³⁰ In Colombia, the government has defined an "economic recognition" with regard to HCWs dealing with COVID-19. Considering the influence of HCWs' financial situation on the development of anxiety symptoms, interventions are needed to handle the anxiety burden in this specific population; such interventions would include better labor and self-care conditions.

Stuijzand et al. discussed different alternatives to address the problem of mental health illnesses in HCWs during pandemics, including the COVID-19 pandemic. One consideration includes dividing the strategies according to the temporality of the outbreak during a pandemic's evolution. Stuijzand et al. placed special emphasis on training of HCWs, planning and assignation of roles for staff and physical resources, and provisioning of sufficient PPE before the disease outbreak. Another strategy highlighted by these researchers was the importance of educational campaigns alerting HCWs about mental health and symptoms that warrant attention and professional help during disease outbreaks. Finally, after a disease outbreak, surveillance should be carried out to detect mental health and occupational problems over one year later.¹⁷

No specific interventions have been shown to mitigate COVID-19-related anxiety in HCWs. Some relevant aspects to improve HCW wellness include the following³⁴: immediate and individualized access to mental health resources, short- and long-term individualized wellness and mental health interventions, individual and organizational strategies to optimize wellness of HCWs (nutrition, exercise, mindfulness, sleep quality), quality and accessible PPE, limitation in the infection risk (through strategies such as telehealth), reduced stigma on mental health symptoms, and development of new HCW community groups to reduce feelings of isolation.

The complexity of these interventions requires a commitment from governmental institutions: the possibility of the intervention being executed not by the government but by health personnel, health care provider institutions (IPS, from its acronym in Spanish), occupational risk management entities (ARL, from its acronym in Spanish), and health insurance companies (EPS, from its acronym in Spanish). In the absence of clear guidelines and a joint effort, implementation

of programs aimed to improve mental health in HCWs will be difficult.

We used a nonrandomly distributed questionnaire to perform the research, causing selection bias in our study. Additionally, levels of anxiety might have modified the participation rate in our study, whether higher levels of anxiety raised or lowered participation rate is unknown. Our sample included a wide variety of HCWs (frontline personnel, medical staff, nurses, administrative staff), and this made it difficult to compare results between specific occupations, which reduced the statistical significance. Likewise, the distribution of our population does not represent the expected distribution of healthcare workers in Colombia, limiting the generalizability of our results. We used the SAS to assess the anxiety prevalence in our population. Even though the SAS is a well-known and validated scale, it is not specific for the evaluation of COVID-related anxiety (or anxiety from any stressor), and many troubles regarding its interpretation have been reported. The original SAS cut-off selected for this study allowed to exclude potential false positives, and to identify participants with more severe anxiety symptoms as positives, reducing the probability of misdiagnosis pursued in a research context. In a clinical context, the modified SAS cut-off could be most appropriate because the false negatives may represent not treated patients.²⁶ We did not include other important variables such as the amount and quality of sleep in the population. Beyond the declared limitations, to our knowledge, this is the first published study to assess anxiety symptoms among HCWs in Colombia during the COVID-19 pandemic, which increases awareness about a problem that is usually disregarded.

Conclusions

From the results discussed above, it can be concluded that almost a third of the Colombian HCW population presented anxiety symptoms, and a positive relationship was found with some demographic variables like sex (greater prevalence for women), history of previous psychiatric illness, being a general practitioner, and cohabiting with people with risk factors of increased mortality from COVID-19 infection. Conversely, subgroups including anesthesiologists, those with access to PPE, older age groups (>40 years), and those at a higher socioeconomic level had fewer anxiety symptoms compared with the rest of the population.

We consider that the proper distribution of PPE and the creation of psychological work groups focused on populations like women and younger professionals could lead to a positive impact on anxiety prevalence among the HCW population.

There is a need to raise awareness about tired and stressed caregivers among HCWs. Timely identification of these characteristics will allow interventions, which would affect not only the health of HCWs but also that of those who are under their medical supervision.

Conflicts of interest

The authors have no conflicts of interest to declare.

REFERENCES

1. Singhal T. A review of coronavirus disease-2019 (COVID-19). *Indian J Pediatr.* 2020;87:281-6.
2. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet.* 2020;395:507-13.
3. Zhai P, Ding Y, Wu X, Long J, Zhong Y, Li Y. The epidemiology, diagnosis and treatment of COVID-19. *Int J Antimicrob Agents.* 2020;55:105955.
4. Hull HF. SARS control and psychological effects of quarantine, Toronto, Canada. *Emerg Infect Dis.* 2005;11:354 [author reply 354-5].
5. Marjanovic Z, Greenglass ER, Coffey S. The relevance of psychosocial variables and working conditions in predicting nurses' coping strategies during the SARS crisis: an online questionnaire survey. *Int J Nurs Stud.* 2007;44:991-8.
6. Reynolds DL, Garay JR, Deamond SL, Moran MK, Gold W, Styra R. Understanding, compliance and psychological impact of the SARS quarantine experience. *Epidemiol Infect.* 2008;136:997-1007.
7. Xiao H, Zhang Y, Kong D, Li S, Yang N. Social capital and sleep quality in individuals who self-isolated for 14 days during the coronavirus disease 2019 (COVID-19) outbreak in January 2020 in China. *Med Sci Monit Int Med J Exp Clin Res.* 2020;26:e923921.
8. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet (London, England).* 2020;395:912-20.
9. Tan BYQ, Chew NWS, Lee GKH, Jing M, Goh Y, Yeo LLL, et al. Psychological impact of the COVID-19 pandemic on health care workers in Singapore. *Ann Intern Med.* 2020;173:317-20.
10. Wu W, Zhang Y, Wang P, Zhang L, Wang G, Lei G, et al. Psychological stress of medical staffs during outbreak of COVID-19 and adjustment strategy. *J Med Virol.* 2020;92:1962-70.
11. Xiao H, Zhang Y, Kong D, Li S, Yang N. The effects of social support on sleep quality of medical staff treating patients with coronavirus disease 2019 (COVID-19) in January and February 2020 in China. *Med Sci Monit Int Med J Exp Clin Res.* 2020;26:e923549.
12. Ripp J, Peccoraro L, Charney D. Attending to the emotional well-being of the health care workforce in a New York City Health System during the COVID-19 pandemic. *Acad Med.* 2020;95:1136-9.
13. Li W, Yang Y, Liu Z-H, Zhao Y-J, Zhang Q, Zhang L, et al. Progression of mental health services during the COVID-19 outbreak in China. *Int J Biol Sci.* 2020;16:1732-8.
14. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Netw open.* 2020;3:e203976.
15. Kang L, Ma S, Chen M, Yang J, Wang Y, Li R, et al. Impact on mental health and perceptions of psychological care among medical and nursing staff in Wuhan during the 2019 novel coronavirus disease outbreak: a cross-sectional study. *Brain Behav Immun.* 2020;87:11-7.
16. Moreno C, Wykes T, Galderisi S, Nordentoft M, Crossley N, Jones N, et al. How mental health care should change as a consequence of the COVID-19 pandemic. *Lancet Psychiatry.* 2020;7:813-24.
17. Stuijzand S, Deforges C, Sandoz V, Sajin C-T, Jaques C, Elmers J, et al. Psychological impact of an epidemic/pandemic on the mental health of healthcare professionals: a rapid review. *BMC Public Health.* 2020;20:1230.

18. Evanoff BA, Strickland JR, Dale AM, Hayibor L, Page E, Duncan JG, et al. Work-related and personal factors associated with mental well-being during the COVID-19 response: survey of health care and other workers. *J Med Internet Res*. 2020;22:e21366.
19. Rivarola Puntigliano A. Pandemics and multiple crises in Latin America. *Lat Am Policy*. 2020;11:313-9.
20. Abuabara YC. Ataque al personal de la salud durante la pandemia de Covid-19 en Latinoamérica. *Acta Médica Colomb*. 2020;45:1-15.
<http://actamedicacolombiana.com/ojs/index.php/actamed/article/view/1975>
21. Pedrozo-Pupo JC, Pedrozo-Cortés MJ, Campo-Arias A. Perceived stress associated with COVID-19 epidemic in Colombia: an online survey. *Cad Saude Publica*. 2020;36, e00090520.
22. Monterrosa-Castro Á, Monterrosa-Blanco A, González-Sequeda A. Perceived loneliness and severe sleep disorders in adult women during the Covid-19 quarantine: a cross-sectional study in Colombia. *J Prim Care Community Health*. 2021;12, 21501327211025170.
23. Caballero-Domínguez CC, De Luque-Salcedo JG, Campo-Arias A. Social capital and psychological distress during Colombian coronavirus disease lockdown. *J Community Psychol*. 2021;49:691-702.
24. Dunstan DA, Scott N. Norms for Zung's self-rating anxiety scale. *BMC Psychiatry*. 2020;20:90.
25. De La Ossa S, Martínez Y, Herazo E, Campo A. Study of internal consistency and factor structure of three versions of the Zung's rating instrument for anxiety disorders. *Colomb Med*. 1969;40:71-84.
<http://www.redalyc.org/src/inicio/ArtPdfRed.jsp?iCve=28340109>
26. Dunstan DA, Scott N. Assigning clinical significance and symptom severity using the Zung scales: levels of misclassification arising from confusion between index and raw scores. *Depress Res Treat*. 2018;2018:9250972.
27. Zhu Z, Xu S, Wang H, Liu Z, Wu J, Li G, et al. COVID-19 in Wuhan: immediate psychological impact on 5062 health workers. *medRxiv*. 2020,
<http://dx.doi.org/10.1101/2020.02.20.20025338>.
<https://www.medrxiv.org/content/10.1101/2020.02.20.20025338v2>
28. Liu CY, Yang YZ, Zhang XM, Xu X, Dou QL, Zhang WW, et al. The prevalence and influencing factors in anxiety in medical workers fighting COVID-19 in China: a cross-sectional survey [Internet]. *Epidemiol Infect*. 2020;148,
<http://dx.doi.org/10.1017/S0950268820001107>.
29. Restrepo-Martínez M, Escobar M, Marín LA, Restrepo D. Prevalence and clinical characteristics of depression and anxiety symptoms in staff at a health institution in Medellín during the COVID-19 pandemic. *Rev Colomb Psiquiatr (English Ed.)*. 2021, <http://dx.doi.org/10.1016/j.rcp.2021.02.001>. S0034-7450(21)00037-8. Advance online publication.
30. Huang L, Wang Y, Liu J, Ye P, Chen X, Xu H, et al. Factors influencing anxiety of health care workers in the Radiology Department with high exposure risk to COVID-19. *Med Sci Monit Int Med J Exp Clin Res*. 2020;26:e926008.
31. Jalnapurkar I, Allen M, Pigott A. Sex differences in anxiety disorders: a review. *Psychiatry Depress Anxiety*. 2018;4:1-9.
<http://www.heraldopenaccess.us/openaccess/sex-differences-in-anxiety-disorders-a-review>
32. Asmundson GJG, Paluszczek MM, Landry CA, Rachor GS, McKay D, Taylor S. Do pre-existing anxiety-related and mood disorders differentially impact COVID-19 stress responses and coping? *J Anxiety Disord*. 2020;74:102271.
33. Taylor S, Landry CA, Rachor GS, Paluszczek MM, Asmundson GJG. Fear and avoidance of healthcare workers: an important, under-recognized form of stigmatization during the COVID-19 pandemic. *J Anxiety Disord*. 2020;75:102289.
34. Shreffler J, Petrey J, Huecker M. The impact of COVID-19 on Healthcare worker wellness: a scoping review. *West J Emerg Med*. 2020;21:1059-66.
35. Zhang SX, Liu J, Afshar Jahanshahi A, Nawaser K, Yousefi A, Li J, et al. At the height of the storm: healthcare staff's health conditions and job satisfaction and their associated predictors during the epidemic peak of COVID-19. *Brain Behav Immun*. 2020;87:144-6.
36. Delgado D, Wyss Quintana F, Perez G, Sosa Liprandi A, Ponte-Negretti C, Mendoza I, et al. Personal safety during the COVID-19 pandemic: realities and perspectives of healthcare workers in Latin America. *Int J Environ Res Public Health*. 2020;17:2798.
37. Xiong J, Lipsitz O, Nasri F, Lui LMW, Gill H, Phan L, et al. Impact of COVID-19 pandemic on mental health in the general population: a systematic review. *J Affect Disord*. 2020;277:55-64.