




Asociación
Colombiana de
Reumatología®

Revista Colombiana de REUMATOLOGÍA

www.elsevier.es/rcreuma



Case Report

Retrobulbar optic neuritis in a patient with enteropathic spondyloarthritis treated with tumour necrosis factor inhibitors

Ben Ayed Hiba^{a,b,*}, Fazaa Alia^{a,b}, Ouenniche Kmar^{a,b}, Miladi Saoussen^{a,b}, Sellami Mariem^{a,b}, Souabni Leila^{a,b}, Kassab Selma^{a,b}, Chekili Selma^{a,b}, Ben Abdelghani Kaouther^{a,b}, Laatar Ahmed^{a,b}

^a Rheumatology Department, University Hospital Center Mongi Slim, La Marsa, Tunisia

^b Faculté de Médecine de Tunis, Tunis El Manar University, Tunis, Tunisia

ARTICLE INFO

Article history:

Received 14 December 2021

Accepted 7 February 2022

Available online 15 mayo 2022

Keywords:

Optic neuritis

Crohn's disease

Spondyloarthritis

ABSTRACT

Optic neuritis is a rare ocular manifestation of inflammatory bowel diseases and spondyloarthritis. The use of tumour necrosis alpha inhibitors to treat these conditions remains possible. We report a case of a 62-year-old-woman with a 17 year-history of axial and peripheral spondyloarthritis associated with Crohn's disease, treated with tumour necrosis alpha inhibitors, who developed an asymmetric retrobulbar optic neuritis that promptly responded to a high dose of steroids.

© 2022 Asociación Colombiana de Reumatología. Published by Elsevier España, S.L.U. All rights reserved.

Neuritis óptica retrobulbar en un paciente con espondiloartritis enteropática tratado con inhibidores del factor de necrosis tumoral

RESUMEN

La neuritis óptica es una manifestación ocular poco común de las enfermedades inflamatorias intestinales y de la espondiloartritis. El uso de fármacos contra el factor de necrosis tumoral (anti-TNF) para tratar estas patologías sigue siendo posible. Se presenta el caso de una mujer de 62 años con un historial médico de espondiloartritis axial y periférica asociada con la enfermedad de Crohn, tratada con anti-TNF y que padece una neuritis óptica asimétrica. La paciente alcanza una respuesta favorable a la terapia basada en los esteroides.

© 2022 Asociación Colombiana de Reumatología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Neuritis óptica

Enfermedad de Crohn

Espondiloartritis

* Corresponding author.

E-mail address: benayedhiba3@gmail.com (B.A. Hiba).

<https://doi.org/10.1016/j.rcreu.2022.02.004>

0121-8123/© 2022 Asociación Colombiana de Reumatología. Published by Elsevier España, S.L.U. All rights reserved.

Introduction

Retrobulbar optic neuritis (RBON) is an inflammatory condition of the optic nerve in which the disease process occurs behind the lamina cribrosa.¹ It is frequently associated with demyelinating diseases as Multiple Sclerosis (MS),² autoimmune disorders and infections.³ However, many other rare causes have been identified. We report the case of a patient with enteropathic spondyloarthritis treated with tumor necrosis factor alpha inhibitors (TNF alpha inhibitors) presenting a RBON.

Case presentation

Our patient is a 62-year-old-woman with a family medical history of MS, a 17 year-history of axial and peripheral spondyloarthritis associated with Crohn's disease (CD), with no previous neurologic or ocular manifestation. She was initially treated with a 20 mg/week of oral Methotrexate for two years, which failed to bring her symptoms under control. Then, she received Infliximab (5 mg/kg). She became asymptomatic after Infliximab-induction therapy (weeks 0, 2, and 6). The treatment was discontinued after the eightieth dose for secondary non-response. Then, she was treated with Etanercept (50 mg/week), discontinued after 4 years for lymphopenia, and lastly with 40 mg/2 weeks of Adalimumab. One year after the switch to Adalimumab, anti-nuclear, anti-double stranded, and anti-histone antibodies were positive, consistent with a drug induced lupus. Adalimumab was discontinued for 6 months then reintroduced after an immunologic remission.

Two years later she presented with an occipital headache, central blind spots in both eyes, and pain with her left eye movement too. At the time of presentation, the neurologic physical examination found a quadripyramidal syndrome, a 10° convergent strabismus in the left eye and a normal pupillary reflex. Extra-ocular movements were full with no overreactions. The visual acuity was 20/20 in both eyes. Anterior segment examination was unremarkable and no optic disc swelling was observed. Formal visual field showed the presence of caecocentral scotoma in both eyes.

Laboratory tests demonstrated normal levels for complete blood count and no vitamin B12 deficiency was detected. The Neuromyelitis optica (NMO) antibodies were negative. A visual evoked potential showed prolonged latency consistent with bilateral demyelinating RBON. MRI of the brain was otherwise normal.

The diagnosis of an asymmetric bilateral RBON was made. Both anti-TNF therapy and inflammatory diseases (CD and spondyloarthritis) were incriminated. The patient received 1 g intravenous Methylprednisolone per day on 3 successive days with a dramatic improvement of blind spots and eye pain. A registration of the case was recorded in the national pharmacovigilance database under the number 482/2012 on April 29, 2019. According to the French method,⁴ the imputability score was C2S1 I1. The decision was to discontinue Adalimumab.

An ophthalmic and a neurologic follow-up were recommended. No signs of relapse were noted during follow-up visits.

Discussion

The frequency of ophthalmic involvement in CD is 6.8%.^{5,6} It is more frequently seen in the first follow-up year, during activity of the bowel disease, in colonic localisation and in the case of coexistence of articular manifestations (enteropathic arthritis).⁷

Episcleritis and acute anterior uveitis are the most frequent ocular complications.⁷ The latter is also reported as the most common extra-articular manifestation of Spondyloarthritis, affecting more than 20% of the patients.⁸ Many other ophthalmic manifestations have been described in relation to inflammatory bowel disease (IBD) such as glaucoma, keratitis and dry eyes.⁷ However, the reported incidence of posterior segment involvement ranges between less than 1% and 30%, depending upon the series.^{8,9} Optic neuritis (ON), which is an inflammation or demyelination or a degeneration of the optic nerve, may be present in up to 4% of adult IBD patients.^{9,10} Recently, a Taiwanese population-based study aimed to evaluate possible factors associated with ON in patients with IBD¹¹: IBD patients who were not with concomitant ON on the index date were included. All the patients were followed up until the development of ON or the end of the study period (the mean follow-up time was of 7 years). A four-fold matched group was selected using age, sex and year of index date for comparison. At the study period conclusion, eight (0.18%) and five (0.003%) patients with and without IBD, respectively, had developed ON ($p=0.001$). Among the eight IBD patients who developed ON only one patient was diagnosed with Crohn's disease, and the male gender was slightly dominant. Adjusted HRs showed that patients aged between 30 and 39 years, with comorbidities including neuromyelitis optica, acute disseminated encephalomyelitis systemic lupus erythematosus and with a higher Charlson Comorbidity Index, had a significantly higher risk of developing ON (all $p < 0.005$).¹¹

To our knowledge, only 6 documented cases of ON with CD were reported in the literature (Table 1).¹² However, the co-existence of ankylosing spondylitis (AS) and ON were only reported in three cases (Table 2).¹³

On the other hand, a possible association between IBD and MS was hypothesized and an approximately 3-fold increased risk of MS in IBD patients was suggested.¹³ A retrospective study performed to examine the relation between the two diseases in the era before TNF alpha inhibitors showed a small but significant association.¹² Patients with CD and ulcerative colitis were 54% and 75% more likely than community controls to have been diagnosed with MS, ON or other demyelinating conditions.¹⁴ ON was recorded in 6 out of 7988 CD patients (0.08%) and in 17 out of 12,185 ulcerative colitis patients (0.14%), in comparison with 50 out of 80,666 controls (0.06%).¹²

TNF alpha inhibitors which are commonly used for the treatment of IBD and refractory rheumatic diseases are incriminated in increasing the risk of developing MS and ON. By 2001, the Food and Drug Administration had received more than 20 reports of MS or other demyelinating conditions in patients treated with these medications.¹² In 2004, clinician and patient warnings were updated on 3 major anti-TNF α therapies: Etanercept, Adalimumab, and Infliximab.¹² The mechanism of this complication is not yet well elucidated. A

Table 1 – Cases of co-existence of Crohn's disease and optic neuritis in the absence of anti-TNF therapy reported in the literature.

Authors Year	Patient characteristics	Presenting symptoms	Treatment	Systemic steroid response
Ernst, B et al. ⁷ 1991	A 25-year-old woman	Blurred Visual acuity in the right eye	Pulse methylprednisolone	Yes
Van de Scheur et al. ¹⁶ 2002	A 39-year-old man	Bilateral visual loss	2 intermittent courses of systemic steroids for 3 weeks in doses of up to 60 mg	Yes
Sung H et al. ¹⁷ 2006	A 22-year-old man	Loss of visual acuity	High dose steroid treatment	Yes
Taxiarchis F et al. ⁸ 2009	A 32-year-old man	Not mentioned	Intravenous methylprednisolone (1 g/day for 3 days)	Not mentioned
-Barabinoet et al. ² 2011	An 11-year-old boy	Bilateral visual loss	Intravenous methylprednisolone (30 mg/kg/day for 3 days) followed by an oral prednisone taper	Yes
-McClelland et al. ¹⁸ 2012	A 42-year-old man	Painless loss of the entire visual field in the left eye	Intravenous methylprednisolone (1 g/day for 1 week) followed by an oral prednisone taper and corticosteroid eye drops	Yes

Table 2 – Cases of co-existence of ankylosing spondylitis and optic neuritis in the literature.

Authors Year	Patient characteristics	Presenting symptoms	Treatment	Systemic steroid response
Menon et al. ¹⁹ 2001	A 20-year-old man	Sudden decrease in vision in the right eye	Intravenous methylprednisolone (1 g for 3 days) followed by 60 mg oral prednisolone/day for 10 days	2 attacks of optic neuritis in the left eye after one year
Y-S Chou et al. ⁴ 2011	A 31-year-old woman	Sudden pain and decrease in vision in the right eye	Pulse steroid therapy for 3 days	No obvious visual field defect on the third day
Zhao et al. ²⁰ 2015	A 31-year-old man	Sudden decrease in vision in both eyes		Improvement of the Visual acuity in the right eye after a follow-up period of 2 years/No improvement of the visual acuity in the left eye after a follow-up period of 2 years

hypothesis suggests that systemically administered anti-TNF agents may inhibit the apoptosis of self-reactive T cells but fail to penetrate and reach the central nervous system, inducing an autoimmune demyelinating process.¹⁵ The other hypothesis suggests an activation of infecting microorganisms which may result in a demyelinating process.¹⁵

In our case, the incrimination of anti-TNF α therapy remains possible, but it is less likely to be the sole cause of the RBON, since one of the most common criterion for causality, which is the absence of alternate cause, is lacking.⁴

Conclusion

ON is rarely the ocular manifestation of AS or CD. To date, the reported cases of demyelination after anti-TNF therapy

suggest a possible causal relationship. However, ON can also be coincidental. Further research should be conducted to elucidate the mechanism of this complication which is a serious adverse event.

Ethical considerations

The institution 's ethics committee was not obtained since this article does not contain personal information that allows to identify the patients. The research complies with the current regulations on bioethical research.

Conflict of interest

None.

Acknowledgements

None.

REFERENCES

- Barabino AV, Gandullia P, Calvi A, Vignola S, Arrigo S, Marco RD. Sudden blindness in a child with Crohn's disease. *World J Gastroenterol*. 2011;17:4344–6, <http://dx.doi.org/10.3748/wjg.v17.i38.4344>.
- Hong D, Bosc C, Chiambaretta F. Progression of nerve fiber layer defects in retrobulbar optic neuritis by the macular ganglion cell complex. *J Fr Ophtalmol*. 2017;40:777–87, <http://dx.doi.org/10.1016/j.jfo.2017.06.003>. French.
- Chou YS, Lu DW, Chen JT. Ankylosing spondylitis presented as unilateral optic neuritis in a young woman. *Ocul Immunol Inflamm*. 2011;19:115–7, <http://dx.doi.org/10.3109/09273948.2010.530732>.
- Moore N, Berdai D, Blin P, Droz C. Pharmacovigilance: the next chapter. *Therapie*. 2019;74:557–67, <http://dx.doi.org/10.1016/j.therap.2019.09.004>.
- Lanna CCD, Ferrari M, de LA, Rocha SL, Nascimento E, de Carvalho MAP, et al. A cross-sectional study of 130 Brazilian patients with Crohn's disease and ulcerative colitis: analysis of articular and ophthalmologic manifestations. *J Clin Rheumatol*. 2008;27:503–9, <http://dx.doi.org/10.1007/s10067-007-0797-5>.
- Mendoza JL, Lana R, Taxonera C, Alba C, Izquierdo S, Díaz-Rubio M. [Extraintestinal manifestations in inflammatory bowel disease: differences between Crohn's disease and ulcerative colitis]. *Med Clin (Barc)*. 2005;125:297–300, <http://dx.doi.org/10.1157/13078423>. Spanish.
- Ernst BB, Lowder CY, Meisler DM, Gutman FA. Posterior segment manifestations of inflammatory bowel disease. *Ophthalmology*. 1991;98:1272–80, [http://dx.doi.org/10.1016/s0161-6420\(91\)32143-2](http://dx.doi.org/10.1016/s0161-6420(91)32143-2).
- Felekis T, Katsanos K, Kitsanou M, Trakos N, Theopistos V, Christodoulou D, et al. Spectrum and frequency of ophthalmologic manifestations in patients with inflammatory bowel disease: a prospective single-center study. *Inflamm Bowel Dis*. 2009;15:29–34, <http://dx.doi.org/10.1002/ibd.20584>.
- Ghanchi FD, Rembacken BJ. Inflammatory bowel disease and the eye. *Surv Ophthalmol*. 2003;48:663–76, <http://dx.doi.org/10.1016/j.survophthal.2003.08.004>.
- Knox DL, Schachat AP, Mustonen E. Primary, secondary and coincidental ocular complications of Crohn's disease. *Ophthalmology*. 1984;91:163–73, [http://dx.doi.org/10.1016/s0161-6420\(84\)34322-6](http://dx.doi.org/10.1016/s0161-6420(84)34322-6).
- Hsieh YH, Chung CH, Sun CA, Chen PH, Chen YH, Liang CM, et al. Association between optic neuritis and inflammatory bowel disease: a population-based study. *J Clin Med*. 2021;10:688, <http://dx.doi.org/10.3390/jcm10040688>.
- Gupta G, Gelfand JM, Lewis JD. Increased risk for demyelinating diseases in patients with inflammatory bowel disease. *Gastroenterology*. 2005;129:819–26, <http://dx.doi.org/10.1053/j.gastro.2005.06.022>.
- Gouveia EB, Elmann D, Morales MSdÁ. Ankylosing spondylitis and uveitis: overview. *Rev Bras Reumatol*. 2012;52:742–56.
- Loftus EV. Inflammatory Bowel disease extending its reach. *Gastroenterology*. 2005;129:1117–20, <http://dx.doi.org/10.1053/j.gastro.2005.07.042>.
- Robinson WH, Genovese MC, Moreland LW. Demyelinating and neurologic events reported in association with tumor necrosis factor alpha antagonism: by what mechanisms could tumor necrosis factor alpha antagonists improve rheumatoid arthritis but exacerbate multiple sclerosis? *Arthritis Rheum*. 2001;44:1977–83, [http://dx.doi.org/10.1002/1529-0131\(200109\)44:9<1977::AID-ART345>3.0.CO;2-6](http://dx.doi.org/10.1002/1529-0131(200109)44:9<1977::AID-ART345>3.0.CO;2-6).
- van de Scheur MR, van der Waal RI, van Bodegraven AA, Völker-Dieben HJ, Starink TM, van der Waal I. Cheilitis granulomatosa and optic neuropathy as rare extraintestinal manifestations of Crohn's disease. *J Clin Gastroenterol*. 2002;34:557–9, <http://dx.doi.org/10.1097/00004836-200205000-00016>.
- Han SH, Lee OY, Yang SY, Jun DW, Lee HL, Jeon YC, et al. A case of optic neuritis associated with Crohn's disease. *Korean J Gastroenterol*. 2006;48:42–5. Korean.
- McClelland C, Zaveri M, Walsh R, Fleisher J, Galetta S. Optic perineuritis as the presenting feature of Crohn disease. *J Neuroophthalmol*. 2012;32:345–7, <http://dx.doi.org/10.1097/WNO.0b013e31825e8e2e>.
- Menon V, Khokhar S. Ankylosing spondylitis in a case of recurrent optic neuritis. *J Neuroophthalmol*. 2001;21:235, <http://dx.doi.org/10.1097/00041327-200109000-00020>.
- Zhao S, Xu Q-G, Zhu J, Peng C-X, Li X-M, Zhou H-F, et al. Acute bilateral optic neuritis in active ankylosing spondylitis. *Chin Med J (Engl)*. 2015;128:2821–2, <http://dx.doi.org/10.4103/0366-6999.167366>.